



## Metric Dimension of Irregular Convex Triangular Networks

S. Prabhu, D. Sagaya Rani Jeba, M. Arulperumjothi & Sandi Klavžar

To cite this article: S. Prabhu, D. Sagaya Rani Jeba, M. Arulperumjothi & Sandi Klavžar (2024) Metric Dimension of Irregular Convex Triangular Networks, AKCE International Journal of Graphs and Combinatorics, 21:3, 225-231, DOI: [10.1080/09728600.2023.2280799](https://doi.org/10.1080/09728600.2023.2280799)

To link to this article: <https://doi.org/10.1080/09728600.2023.2280799>



© 2023 The Author(s). Published with license by Taylor & Francis Group, LLC



Published online: 07 Dec 2023.



Submit your article to this journal [↗](#)



Article views: 395



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)

# Metric Dimension of Irregular Convex Triangular Networks

S. Prabhu<sup>a</sup>, D. Sagaya Rani Jeba<sup>b</sup>, M. Arulperumjothi<sup>c</sup>, and Sandi Klavžar<sup>d,e,f</sup>

<sup>a</sup>Department of Mathematics, Rajalakshmi Engineering College, Chennai, India; <sup>b</sup>Department of Mathematics, Panimalar Engineering College, Chennai, India; <sup>c</sup>Department of Mathematics, St. Joseph's College of Engineering, Chennai, India; <sup>d</sup>Faculty of Mathematics and Physics, University of Ljubljana, Slovenia; <sup>e</sup>Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia; <sup>f</sup>Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia

## ABSTRACT

Irregular convex triangular networks consist of the interior of a 6-sided convex polygon drawn on the infinite triangular network. Formal description of these applicable networks is provided. In the main result it is proved that the metric dimension of an irregular convex triangular network is either 2 or 3 and the determination of which value is the right one is specified. Metric dimensions of various graph classes and interconnection networks are also collected.

## ARTICLE HISTORY

Received 18 October 2023  
Accepted 3 November 2023

## KEYWORDS

Graph networks; metric dimension; irregular convex triangular network

**AMS SUBJECT CLASSIFICATION**  
05C12; 05C90

## 1. Introduction

Network science has risen as a multidisciplinary effort to estimate the structural features of complex systems due to the requirement to define interconnections mathematically [5]. Due to the development of big data, the concept of a network (a collective of components, alias “nodes,” linked together by the connections, alias “edges,” among them) has emerged as a practical way to represent a wide variety of empirical knowledge in physical, biological, and social phenomena [37]. Network science acquired its basic principles via graph theory, mathematically preoccupied with pairwise relationships between features [33], and was inspired by landmark works in sociology and economics [9, 53]. Graph theory and network science, on the other hand, have gone their own ways since then, with so little convergence in their academic buildings and research community. This subject is discussed in [12, 47]. The challenge of coming up with a unique representation for each vertex in a connected graph arises in many contexts including robot navigation [27], coin weighing problems [49], connected joins in graphs [44], hierarchical data structures [38], network discovery and verification [7] and mastermind game strategy [18].

The tessellation of the plane into triangles is the foundation of hexagonal networks. Networks with a regular square separation are termed mesh networks, and those with a regular hexagonal partition are called honeycomb networks. There is a degree of inconsistency in the choice of the name, which arises from the duality between graphs that correspond to hexagonal networks and honeycomb networks. These networks are applied in many areas, say in mathematical chemistry, in image processing, in

wireless networks, in computer graphics, and in interconnection networks, cf. [39]. As a generalization of the planar hexagonal networks, higher dimensional hexagonal networks have been defined in [14, 39], and the honeycomb design has been presented in [50]. For a network with hexagonal nodes, an addressing method for the processors and accompanying routing and broadcasting algorithms have been presented [13]. Cellular networks have employed hexagonal networks for user tracking and connection rerouting [39].

In this paper we consider the metric dimension of irregular convex triangular networks and proceed as follows. In the next section we recall the metric dimension and make a small survey of known results related to it, where we concentrate on determining the metric dimension of graph classes. In Section 3 we introduce the networks of interest in this paper and prepare the mathematical formalities on them that we need for the following. Then, in Section 4 we determine their metric dimension.

## 2. The metric dimension

Visual identification of a distinctive landmark by a robot moving in Euclidean space offers information about the landmark's direction and helps the robot to pinpoint its position. However, a graph lacks any discernible direction or transparency. Moreover, it is generally accepted that a robot moving along a graph can gauge how far apart landmarks are as it travels them. Clearly, if the robot knows the distances to a big enough selection of landmarks, its position on the graph may be computed uniquely. A metric basis is a set of landmarks that uniquely determines the robot's position, and the metric dimension of the graph

[27] is the lowest number of landmarks. As a result, a graph-theoretic understanding of this problem is to construct codes for the vertices of a graph in such a way that each vertex has a unique code. The recent publications [1, 25, 29, 31, 32, 41, 52] all reflect on this.

Let  $G = (V(G), E(G))$  be a connected graph. The distance between vertices  $u$  and  $v$  is the length of a shortest  $u, v$ -path and denoted by  $d_G(u, v)$ . If  $G$  will be clear from the context, we may also write  $d(u, v)$ . For an ordered set of vertices  $R = \{r_1, r_2, \dots, r_l\}$  and a vertex  $u \in V(G)$ , the representation of  $u$  with respect to  $R$  is the vector  $(d_G(u, r_1), d_G(u, r_2), \dots, d_G(u, r_l))$ . If the vertices of  $G$  have pairwise different representations with respect to  $R$ , then  $R$  is a *resolving set* for  $G$ . That is,  $R$  is a resolving set if for each pair of different vertices  $x, y \in V(G)$  there exists a vertex  $r \in R$  such that  $d_G(x, r) \neq d_G(y, r)$ . The *metric dimension*  $\dim(G)$  of  $G$  is the cardinality of a smallest resolving set for  $G$ , such a set is called a *metric basis*.

These concepts were first suggested by Slater [47, 48], who was inspired by applications to the placement of a small number of sonar detecting devices in a network such that the position of each vertex in the network may be uniquely identified in terms of its distance from the set of devices. Harary and Melter [23] independently introduced this concept. Two decades later, Khuller et al. [27] also (independently) found these ideas and coined the term metric dimension. While attempting to establish a capability for enormous data sets of chemical graphs, these notions were rediscovered once more in [12, 26].

Determination of the metric dimension is NP-hard for general graphs [27], and, moreover, the problem remains NP-hard on bipartite graphs [34]. This makes it interesting to determine the metric dimension for specific classes of graphs, and so far this has been very intensively investigated. In Table 1 we have

**Table 1.** Metric dimension of various classes.

Structure	Reference
Trees, multi-dimensional grids	Khuller et al. [27]
Line graphs, para-line graphs	Klein et al. [28]
Torus networks	Manuel et al. [35]
Some families of graphs	Cáceres et al. [11]
Cayley digraphs	Fehr et al. [15]
Unicyclic graphs	Poisson et al. [40]
Petersen graphs	Shao et al. [45]
Circulant graphs	Rajan et al. [19]; Imran et al. [24]
Harary graphs	Rajan et al. [19]
Illiac networks	Rajan et al. [43]
Jahangir graph	Tomescu et al. [51]
Enhanced hypercubes	Rajan et al. [42]
Random graphs	Bollobas et al. [8]
Benes networks	Manuel et al. [34]
Honeycomb networks	Manuel et al. [36]
Distance-regular graphs	Guo et al. [21]
Infinite graphs	Cáceres et al. [10]
Hypercubes	Beardon [6]
Wheel-related graphs	Siddiqui et al. [46]
Bilinear forms graphs	Feng et al. [16]
Grassmann graphs	Bailey et al. [4]
Symplectic dual polar graphs	Guo et al. [20]
Permutation graphs	Hallaway et al. [22]
Kneser and Johnson graphs	Bailey et al. [3]
$TiO_2[m, n]$	Prabhu et al. [41]
Chain graphs	Fernau et al. [17]
Convex polytopes	Imran et al. [25]
Incidence graphs	Bailey [2]

gathered the work on the metric dimension of various graph classes and interconnection networks.

To conclude this section we recall the following result needed later on.

**Theorem 2.1.** [27] If  $\{s, t\}$  is a metric basis of a graph  $G$ , then  $\deg_G(s) \leq 3$ ,  $\deg_G(t) \leq 3$ , and there is exactly one shortest  $s, t$ -path  $P$ . Moreover, if  $w$  is an internal vertex of  $P$ , then  $\deg_G(w) \leq 5$ .

### 3. Irregular convex triangular networks

In this section we introduce irregular convex triangular networks and provide their formal description. Intuitively, they consist from the interior of a 6-sided convex polygon drawn on the infinite triangular network.

If  $r < s$  are integers, then we will denote the set  $\{r, r + 1, \dots, s\}$  by  $\llbracket r, s \rrbracket$ . For given parameters  $n, m, p$ , and  $q$ , let  $\{H_0, H_1, \dots, H_{2n+m-p-q}\}$ ,  $\{A_0, A_1, \dots, A_n\}$  and  $\{O_0, O_1, \dots, O_{m+n}\}$  be three respective sets of lines corresponding to horizontal, acute, and obtuse lines fulfilling the following conditions:

1.  $\angle H_i A_j = \pi/3$  for  $i \in \llbracket 0, m + 2n - p - q \rrbracket$  and  $j \in \llbracket 0, n \rrbracket$ ;
2.  $\angle H_i O_k = 2\pi/3$  for  $i \in \llbracket 0, m + 2n - p - q \rrbracket$  and  $k \in \llbracket 0, m + n \rrbracket$ ;
3.  $\angle A_j O_k = \pi/3$  for  $j \in \llbracket 0, n \rrbracket$  and  $k \in \llbracket 0, m + n \rrbracket$ .

Assuming the above conditions, a *convex triangular grid*  $\Delta_{2n+m-p-q, n, m+n}$  is a graph derived by treating all triple intersection points as vertices and the line joining all triple intersection points as edges. (Many non-isomorphic convex triangular grids have the same parameters [30].) If the boundary of the triangular grid  $\Delta_{h, a, o}$  is a trapezium, we say that the grid is a *triangular trapezium*. In this case we have  $a = o$  in  $\Delta_{h, a, o}$ . The height and the length of the base are respectively denoted by  $h - 1$  and  $a - 1$ , the graph is denoted by  $T_{h-1, a-1}$ . This triangular trapezium is reduced to a *triangular triangle*  $T_{h-1}$  when  $h = a$ . By  $I_{2n-1}$ ,  $n \geq 1$ , we denote a chain of  $(2n - 1)$ -triangles. It is easy to observe that  $I_{2n-1} \cong T_{1, n}$ . In  $\Delta_{h, a, o}$ , if the boundary of  $\Delta_{h, a, o}$  forms a parallelogram, then  $\Delta_{h, a, o}$  is a *triangular parallelogram*. For this case we have  $h - 1 = |o - a|$  and  $\Delta_{h, a, o} \cong \Delta_{h, o, a}$ . For a triangular parallelogram we always assume that  $a < o$  and denote  $\Delta_{h, a, h+a-1}$  by  $P_{h-1, a-1}$ . Also for  $n \geq 1$ , we have  $P_{1, n} \cong I_{2n}$ . Several examples of these networks can be seen in Figure 1.

Now, an *irregular convex triangular hexagon*  $H_{p, q, m, n}$  is formed by merging the base of  $T_{n-q, n}$  and  $T_{n-p, n}$  with two bases of  $P_{m, n}$ , where  $p \in \llbracket 0, n \rrbracket$ ,  $q \in \llbracket 0, n \rrbracket$ , and  $m \geq 0$ . By symmetry we have  $H_{p, q, m, n} \cong H_{q, p, m, n}$ , hence we may assume in the rest that always  $p \leq q$  holds. Also note that  $T_{n, n-q} \cong T_{n-q, n} \cong H_{q, 0, 0, n}$ ;  $P_{m, n} \cong P_{n, m} \cong H_{0, 0, m, n}$ ;  $H_n \cong H_{p, p, 0, 2p}$ ;  $I_{2n-1} \cong H_{1, 0, 0, n}$  and  $I_{2n} \cong H_{0, 0, 1, n}$ . The structure and the construction of  $H_{p, q, m, n}$  are illustrated in Figure 2.

To complete the description of the irregular convex triangular networks, we introduce the coordinates of its vertices in a natural way. That is, based on the horizontal, acute, and obtuse lines, a vertex of  $H_{p, q, m, n}$  is assigned the natural coordinates  $(h, a, o)$ , see Figure 3.




Figure 1. (a)  $P_{4,5}$ ; (b)  $T_5$ ; (c)  $T_{3,6}$ ; (d)  $H_3$ ; (e)  $H_{3,4,3,6}$ ; (f) Convex parts of  $H_{3,4,3,6}$ .




Figure 2. Construction of irregular convex triangular networks.

#### 4. The metric dimension of irregular convex hexagonal networks

The main result of this paper reads as follows.

**Theorem 4.1.** If  $H_{p,q,m,n}$  is an irregular convex hexagonal net, then

$$\dim(H_{p,q,m,n}) = \begin{cases} 2; & m \geq p, \\ 3; & m < p. \end{cases}$$

In the rest of the section we prove **Theorem 4.1**. We first show:

**Lemma 4.2.** If  $m < p$ , then  $\dim(H_{p,q,m,n}) > 2$ .

**Proof.** Suppose on the contrary that  $\dim(H_{p,q,m,n}) = 2$  and let  $\{u, v\}$  be a resolving set of  $H_{p,q,m,n}$ . By **Theorem 2.1**,  $\deg_G(u) \leq 3$ ,  $\deg_G(v) \leq 3$ , and there is a unique shortest  $u, v$ -path. Therefore, we need to consider the following six cases.

**Case 1:**  $u = (0, n - p, n + m)$  and  $v = (0, n, m + n - p)$ . We have  $d(u, (p + 1, n - p - 1, m + n - p)) = d(u, (p + 1, n - p, m + n - p - 1))$  and  $d(v, (p + 1, n - p - 1, m + n - p)) = d(v, (p + 1, n - p, m + n - p - 1))$ , a contradiction.

**Case 2:**  $u = (0, n, m + n - p)$  and  $v = (n + m - p, n, 0)$ . We have  $d(u, (n - p + m, p - m - 1, n - p + m + 1)) = d(u, (n - p + m + 1, p - m - 1, n - p + m))$  and  $d(v, (n - p + m, p - m - 1, n - p + m + 1)) = d(v, (n - p + m + 1, p - m - 1, n - p + m))$ , a contradiction.

**Case 3:**  $u = (n + m - p, n, 0)$  and  $v = (2n + m - p - q, q, 0)$ . We have  $d(u, (n - p + m, q - 1, n - q + 1)) = d(u, (n - p + m - 1, q, n - q + 1))$  and  $d(v, (n - p + m, q - 1, n - q + 1)) = d(v, (n - p + m - 1, q, n - q + 1))$ , a contradiction.

**Case 4:**  $u = (2n + m - p - q, q, 0)$  and  $v = (2n + m - p - q, 0, q)$ . We have  $d(u, (2n + m - p - 2q - 1, q, q + 1)) = d(u, (2n + m - p - 2q - 1, q + 1, q))$  and  $d(v, (2n + m - p - 2q - 1, q, q + 1)) = d(v, (2n + m - p - 2q - 1, q + 1, q))$ , a contradiction.

**Case 5:**  $u = (2n + m - p - q, 0, q)$  and  $v = (n - p, 0, m + n)$ . Then we have  $d(u, (n - p, m + n - q + 1, q - 1)) = d(u, (n - p - 1, m + n - q + 1, q))$  and  $d(v, (n - p, m + n - q + 1, q - 1)) = d(v, (n - p - 1, m + n - q + 1, q))$ , a contradiction.

**Case 6:** Let  $u = (n - p, 0, m + n)$  and  $v = (0, n - p, n + m)$ . We have,  $d(u, (n - p, n - p + 1, m + p - 1)) = d(u, (n - p + 1, n - p, m + p - 1))$  and  $d(v, (n - p, n - p + 1, m + p - 1)) = d(v, (n - p + 1, n - p, m + p - 1))$ , a contradiction.

We conclude that  $\dim(H_{p,q,m,n}) > 2$ . □

If  $u$  is a vertex of a  $H_{p,q,m,n}$  and  $r$  is a positive integer, then let  $N_r(u)$  denote the set of vertices of  $H_{p,q,m,n}$  which are at distance  $r$  from  $u$ . In addition, we will use the following convention. If  $N_r(u)$  consists of consecutive vertices from a horizontal line, then this will be denoted by  $N_r(u) = P_H$ . If needed, we will




Figure 3. (a) Addressing scheme; (b) Coordinates of corner vertices of  $H_{p,q,m,n}$ .

also identify  $P_H$  with the set of vertices in the horizontal line that are at distance  $r$  from  $u$ . We will use analogues conventions  $P_A$  and  $P_O$  when  $N_r(u)$  consists of consecutive vertices from an acute or an obtuse line, respectively. Moreover, if  $N_r(u)$  consists of, say, some consecutive vertices from a horizontal line followed by some consecutive vertices from an obtuse line, this will be briefly denoted by  $N_r(u) = P_H \cup P_O$ .

**Lemma 4.3.** In any  $H_{p,q,m,n}$  we have

1.  $N_r((0, n-p, n+m)) \in \{P_H \cup P_O, P_H, P_O\}$ .
2.  $N_r((0, n, m+n-p)) \in \{P_H \cup P_A, P_H\}$ .
3.  $N_r((m+n-p, n, 0)) \in \{P_A \cup P_O, P_O\}$ .

**Proof.** Suppose first that  $n-p-q \geq 0$ . Then

$$N_r((0, n-p, n+m)) = \begin{cases} P_H \cup P_O; & r \in \llbracket 1, m+n \rrbracket, \\ P_H; & r \in \llbracket 1+m+n, m+2n-q-p \rrbracket, \end{cases}$$

where

$$P_H = \begin{cases} \{(r, n-p-r+i, m+n-i) : i \in \llbracket 0, r \rrbracket\}; \\ \quad r \in \llbracket 1, n-p \rrbracket, \\ \{(r, i, m+2n-i-p-r) : i \in \llbracket 0, n-p \rrbracket\}; \\ \quad r \in \llbracket 1+n-p, m+n \rrbracket, \\ \{(r, i, m+2n-i-p-r) : i \in \llbracket 0, m+2n-r-p \rrbracket\}; \\ \quad r \in \llbracket 1+n+m, m+2n-q-p \rrbracket, \end{cases}$$

and

$$P_O = \begin{cases} \{(r-i, n+i-p, m+n-r) : i \in \llbracket 1, r \rrbracket\}; \\ \quad r \in \llbracket 1, p \rrbracket, \\ \{(r-i, n+i-p, m+n-r) : i \in \llbracket 1, p \rrbracket\}; \\ \quad r \in \llbracket 1+p, m+n \rrbracket. \end{cases}$$

If  $n-p-q < 0$ , then

$$N_r((0, n-p, n+m)) = \begin{cases} P_H \cup P_O; & r \in \llbracket 1, m+2n-p-q \rrbracket, \\ P_O; & r \in \llbracket 1+m+2n-p-q, m+n \rrbracket, \end{cases}$$

where

$$P_H = \begin{cases} \{(r, n+i-p-r, m+n-i) : i \in \llbracket 0, r \rrbracket\}; \\ \quad r \in \llbracket 1, n-p \rrbracket, \\ \{(r, i, m+2n-i-p-r) : i \in \llbracket 0, n-p \rrbracket\}; \\ \quad r \in \llbracket 1+n-p, m+2n-p-q \rrbracket, \end{cases}$$

and

$$P_O = \begin{cases} \{(r-i, n+i-p, m+n-r) : i \in \llbracket 1, r \rrbracket\}; \\ \quad r \in \llbracket 1, p \rrbracket, \\ \{(r-i, n+i-p, m+n-r) : i \in \llbracket 1, p \rrbracket\}; \\ \quad r \in \llbracket 1+p, m+2n-p-q \rrbracket, \\ \{(r-i, n+i-p, m+n-r) : i \in \llbracket r-m-2n+p+q, p \rrbracket\}; \\ \quad r \in \llbracket 1+m+2n-p-q, m+n \rrbracket. \end{cases}$$

For  $m-p \geq 0$  we have

$$N_r((0, n, m+n-p)) = \begin{cases} P_H \cup P_A; & r \in \llbracket 1, n \rrbracket, \\ P_H; & r \in \llbracket 1+n, m+2n-p-q \rrbracket, \end{cases}$$

where

$$P_H = \begin{cases} \{(r, n-i, m+n+i-p-r) : i \in \llbracket 0, r \rrbracket\}; \\ \quad r \in \llbracket 1, n \rrbracket, \\ \{(r, n-i, m+n+i-p-r) : i \in \llbracket 0, n \rrbracket\}; \\ \quad r \in \llbracket 1+n, m+n-p \rrbracket, \\ \{(r, m+2n-i-p-r, i) : i \in \llbracket 0, m+2n-p-r \rrbracket\}; \\ \quad r \in \llbracket 1+m+n-p, m+2n-p-q \rrbracket, \end{cases}$$

and

$$P_A = \begin{cases} \{(r-i, n-r, m+n+i-p) : i \in \llbracket 1, r \rrbracket\}; \\ \quad r \in \llbracket 1, p \rrbracket, \\ \{(r-i, n-r, m+n+i-p) : i \in \llbracket 1, p \rrbracket\}; \\ \quad r \in \llbracket 1+p, n \rrbracket. \end{cases}$$

If  $m-p < 0$ , then

$$N_r((0, n, m+n-p)) = \begin{cases} P_H \cup P_A; & r \in \llbracket 1, n \rrbracket, \\ P_H; & r \in \llbracket 1+n, m+2n-p-q \rrbracket, \end{cases}$$

where

$$P_H = \begin{cases} \{(r, n - i, m + n + i - p - r) : i \in \llbracket 0, r \rrbracket\}; \\ r \in \llbracket 1, m + n - p \rrbracket, \\ \{(r, m + 2n - i - p - r, i) : i \in \llbracket 0, m + 2n - p - r \rrbracket\}; \\ r \in \llbracket 1 + m + n - p, m + 2n - p - q \rrbracket, \end{cases}$$

and

$$P_A = \begin{cases} \{(r - i, n - r, m + n + i - p) : i \in \llbracket 1, r \rrbracket\}; \\ r \in \llbracket 1, p \rrbracket, \\ \{(r - i, n - r, m + n + i - p) : i \in \llbracket 1, p \rrbracket\}; \\ r \in \llbracket 1 + p, n \rrbracket. \end{cases}$$

If  $m - p \geq 0$ , then we have

$$N_r((m + n - p, n, 0)) = \begin{cases} P_A \cup P_O; & r \in \llbracket 1, n \rrbracket, \\ P_O; & r \in \llbracket 1 + n, m + n \rrbracket, \end{cases}$$

where

$$P_A = \begin{cases} \{(m + n - i - p + r, n - r, i) : i \in \llbracket 0, r \rrbracket\}; \\ r \in \llbracket 1, n - p \rrbracket, \\ \{(m + 2n - i - p - q, n - r, r - n + i + q) : \\ i \in \llbracket 0, n - q \rrbracket\}; \\ r \in \llbracket 1 + n - q, n \rrbracket, \end{cases}$$

and

$$P_O = \begin{cases} \{(m + n - p - i, n + i - r, r) : i \in \llbracket 1, r \rrbracket\}; \\ r \in \llbracket 1, n \rrbracket, \\ \{(1 + m + 2n - i - p - r, i - 1, r) : i \in \llbracket 1, 1 + n \rrbracket\}; \\ r \in \llbracket 1 + n, m + n - p \rrbracket, \\ \{(1 + m + 2n - i - p - r, i - 1, r) : \\ i \in \llbracket 1, 1 + m + 2n - p - r \rrbracket\}; \\ r \in \llbracket 1 + m + n - p, m + n \rrbracket. \end{cases}$$

Finally, if  $m - p < 0$ , then

$$N_r((m + n - p, n, 0)) = \begin{cases} P_A \cup P_O; & r \in \llbracket 1, n \rrbracket, \\ P_O; & r \in \llbracket 1 + n, m + n \rrbracket, \end{cases}$$

where

$$P_A = \begin{cases} \{(m + n - i - p + r, n - r, i) : i \in \llbracket 0, r \rrbracket\}; \\ r \in \llbracket 1, n - q \rrbracket, \\ \{(m + 2n - i - p - q, n - r, r - n + i + q) : \\ i \in \llbracket 0, n - q \rrbracket\}; \\ r \in \llbracket 1 + n - q, n \rrbracket, \end{cases}$$

and

$$P_O = \begin{cases} \{(m + n - i - p, n + i - r, r) : i \in \llbracket 1, r \rrbracket\}; \\ r \in \llbracket 1, m + n - p \rrbracket, \\ \{(m + n - p - i, n + i - r, r) : i \in \llbracket 1, m + n - q \rrbracket\}; \\ r \in \llbracket 1 + m + n - q, n \rrbracket, \\ \{(1 + m + 2n - i - p - r, i - 1, r) : \\ i \in \llbracket 1, 1 + m + 2n - p - r \rrbracket\}; \\ r \in \llbracket 1 + n, m + n \rrbracket. \end{cases}$$

**Lemma 4.4.** In any  $H_{p,q,m,n}$ , if  $r_1, r_2 \in \llbracket 1, m + 2n - p - q \rrbracket$ , then  $N_{r_1}((0, n - p, n + m)) \cap N_{r_2}((0, n, m + n - p))$  is either empty, or a vertex, or a line segment of a horizontal line.

**Proof.** From the proof of Lemma 4.3 we recall that if  $n - q - p \geq 0$ , then

$$N_r((0, n - p, n + m)) = \begin{cases} P_H \cup P_O; & r \in \llbracket 1, m + n \rrbracket, \\ P_H; & r \in \llbracket 1 + m + n, m + 2n - p - q \rrbracket, \end{cases}$$

and

$$N_r((0, n, m + n - p)) = \begin{cases} P_H \cup P_A; & r \in \llbracket 1, n \rrbracket, \\ P_H; & r \in \llbracket 1 + n, m + 2n - p - q \rrbracket. \end{cases}$$

Moreover, if  $n - q - p < 0$ , then

$$N_r((0, n - p, n + m)) = \begin{cases} P_H \cup P_O; & r \in \llbracket 1, m + 2n - p - q \rrbracket, \\ P_O; & r \in \llbracket 1 + m + 2n - p - q, m + n \rrbracket, \end{cases}$$

and

$$N_r((0, n, m + n - p)) = \begin{cases} P_H \cup P_A; & r \in \llbracket 1, n \rrbracket, \\ P_H; & r \in \llbracket 1 + n, m + 2n - p - q \rrbracket. \end{cases}$$

The conclusion then follows from these facts.  $\square$

**Lemma 4.5.** In any  $H_{p,q,m,n}$ , if  $r_1 \in \llbracket 1, 2n + m - p - q \rrbracket$ ,  $r_2 \in \llbracket 1, n + m \rrbracket$ , and  $m - p \geq 0$ , then  $|N_{r_1}((0, n, m + n - p)) \cap N_{r_2}((m + n - p, n, 0))| \in \{0, 1\}$ .

**Proof.** From the proof of Lemma 4.3 we recall that if  $m - p \geq 0$ , then

$$N_r((0, n, m + n - p)) = \begin{cases} P_H \cup P_A; & r \in \llbracket 1, n \rrbracket, \\ P_H; & r \in \llbracket 1 + n, m + 2n - p - q \rrbracket, \end{cases}$$

and

$$N_r((m + n - p, n, 0)) = \begin{cases} P_A \cup P_O; & r \in \llbracket 1, n \rrbracket, \\ P_O; & r \in \llbracket 1 + n, m + n \rrbracket. \end{cases}$$

**Case 1:**  $n - p - q > 0$ .

Assume first that  $r_1, r_2 \in \llbracket 1, m + n - p \rrbracket$ . If  $r_1 + r_2 < m + n - p$ , then  $N_{r_1}((0, n, m + n - p)) \cap N_{r_2}((m + n - p, n, 0)) = \emptyset$ , and if  $r_1 + r_2 \geq m + n - p$ , then  $N_{r_1}((0, n, m + n - p)) \cap N_{r_2}((m + n - p, n, 0))$  is a singleton.

Assume second that  $r_1 \in \llbracket 1 + m + n - p, m + 2n - p - q \rrbracket$  and  $r_2 \in \llbracket 1, n \rrbracket$ . If  $r_1 - m - n + p - r_2 > 0$ , then  $N_{r_1}((0, n, m + n - p)) \cap N_{r_2}((m + n - p, n, 0)) = \emptyset$ , and if  $r_1 - m - n + p - r_2 \leq 0$ , then  $N_{r_1}((0, n, m + n - p)) \cap N_{r_2}((m + n - p, n, 0))$  is a singleton.

Assume finally that  $r_1 \in \llbracket 1, n \rrbracket$  and  $r_2 \in \llbracket 1 + m + n - p, m + n \rrbracket$ . Now, if  $r_2 - m - n + p - r_1 > 0$ , then  $N_{r_1}((0, n, m + n - p)) \cap N_{r_2}((m + n - p, n, 0)) = \emptyset$ , and if  $r_2 - m - n + p - r_1 \leq 0$ , then  $N_{r_1}((0, n, m + n - p)) \cap N_{r_2}((m + n - p, n, 0))$  is a singleton.

$\square$  **Case 2:**  $n - p - q \leq 0$ .

Assume first that  $r_1, r_2 \in \llbracket 1, n \rrbracket$ . If  $r_1 + r_2 < m + n - p$ , then  $N_{r_1}((0, n, m + n - p)) \cap N_{r_2}((m + n - p, n, 0)) = \emptyset$ , and if  $r_1 + r_2 \geq m + n - p$ , then  $N_{r_1}((0, n, m + n - p)) \cap N_{r_2}((m + n - p, n, 0))$  is a singleton.



Assume second that  $r_1 \in \llbracket 1 + n, m + 2n - p - q \rrbracket$  and  $r_2 \in \llbracket 1, n \rrbracket$ . If  $r_1 - m - n + p - r_2 > 0$ , then  $N_{r_1}((0, n, m + n - p)) \cap N_{r_2}((m + n - p, n, 0)) = \emptyset$ , and if  $r_1 - m - n + p - r_2 \leq 0$ , then  $N_{r_1}((0, n, m + n - p)) \cap N_{r_2}((m + n - p, n, 0))$  is singleton.

Assume finally that  $r_1 \in \llbracket 1, n \rrbracket$  and  $r_2 \in \llbracket 1 + n, m + n \rrbracket$ . If  $r_2 - n - r_1 > 0$ , then  $N_{r_1}((0, n, m + n - p)) \cap N_{r_2}((m + n - p, n, 0)) = \emptyset$ , and if  $r_2 - n - r_1 \leq 0$ , then  $N_{r_1}((0, n, m + n - p)) \cap N_{r_2}((m + n - p, n, 0))$  is a singleton.  $\square$

**Corollary 4.6.** If  $u = (h_1, a_1, o_1), v = (h_2, a_2, o_2)$  are vertices of  $H_{p,q,m,n}$  with  $h_1 \neq h_2, a_1 \neq a_2$ , and  $o_1 \neq o_2$ , then  $|N_{r_1}((0, n - p, n + m)) \cap N_{r_2}((0, n, m + n - p)) \cap \{u, v\}| \leq 1$ .

**Proof.** Suppose on the contrary that  $u$  and  $v$  are both in  $N_{r_1}((0, n - p, n + m)) \cap N_{r_2}((0, n, m + n - p))$ . Then Lemma 4.4 implies that  $N_{r_1}((0, n - p, n + m)) \cap N_{r_2}((0, n, m + n - p))$  is a line segment of a horizontal line. But then  $h_1 = h_2$ , a contradiction.  $\square$

To complete the proof of Theorem 4.1 it remains to demonstrate (having Lemma 4.2 in mind) that if  $m < p$ , then there exists a metric basis of cardinality 3, and if  $m \geq p$ , then there exists a metric basis of cardinality 2. We establish these facts in the concluding couple of lemmas.

**Lemma 4.7.** If  $m < p$ , then  $\{(0, n - p, n + m), (0, n, m + n - p), (m + n - p, n, 0)\}$  forms a metric basis of  $H_{p,q,m,n}$ .

**Proof.** Let  $u = (h_1, a_1, o_1)$  and  $v = (h_2, a_2, o_2)$  be two arbitrary vertices of  $H_{p,q,m,n}$ .

**Case 1:**  $h_1 = h_2$ .

Since in this case  $u$  and  $v$  lie in a segment of a horizontal line we infer that  $d(u, (m + n - p, n, 0)) \neq d(v, (m + n - p, n, 0))$ .

**Case 2:**  $a_1 = a_2$ .

Now we have  $d(u, (0, n - p, n + m)) \neq d(v, (0, n - p, n + m))$ .

**Case 3:**  $o_1 = o_2$ .

As  $u$  and  $v$  are in a segment of an obtuse line,  $d(u, (0, n, m + n - p)) \neq d(v, (0, n, m + n - p))$ .

**Case 4:**  $h_1 \neq h_2, a_1 \neq a_2$ , and  $o_1 \neq o_2$ .

Suppose that  $d(u, (0, n - p, n + m)) = d(v, (0, n - p, n + m)) = r_1$ . Then  $u, v \in N_{r_1}((0, n - p, n + m))$  and we claim that  $d(u, (0, n, m + n - p)) \neq d(v, (0, n, m + n - p))$ . Suppose that  $d(u, (0, n, m + n - p)) = d(v, (0, n, m + n - p)) = r_2$ . Then  $u, v \in N_{r_2}((0, n, m + n - p))$  which implies that  $\{u, v\} \subseteq N_{r_1}((0, n - p, n + m)) \cap N_{r_2}((0, n, m + n - p))$ . This contradiction to Corollary 4.6 yields  $d(u, (0, n, m + n - p)) \neq d(v, (0, n, m + n - p))$ .  $\square$

**Lemma 4.8.** If  $m \geq p$ , then  $\{(0, n, m + n - p), (m + n - p, n, 0)\}$  forms a metric basis of  $H_{p,q,m,n}$ .

**Proof.** Apply Lemma 4.5.  $\square$

## Funding

Sandi Klavžar was supported by the Slovenian Research Agency (ARRS) under the grants P1-0297, J1-2452, and N1-0285.

## References

- [1] Akhter, S., Farooq, R. (2019). Metric dimension of fullerene graphs. *Electron. J. Graph Theory Appl.* 7(1): 91–103.
- [2] Bailey, R. F. (2018). On the metric dimension of incidence graphs. *Discrete Math.* 341(6): 1613–1619.
- [3] Bailey, R. F., Cáceres, J., Garijo, D., González, A., Márquez, A., Meagher, K., Puertas, M. L. (2013). Resolving sets for Johnson and Kneser graphs. *Eur. J. Comb.* 34(4): 736–751.
- [4] Bailey, R. F., Meagher, K. (2011). On the metric dimension of Grassmann graphs. *Discrete Math. Theor. Comput. Sci.* 13(4): 97–104.
- [5] Barabási, A. L. (2013). Network science. *Philos. Trans. Soc. London Ser. A* 371: 20120375.
- [6] Beardon, A. F. (2013). Resolving the hypercube. *Discrete Appl. Math.* 161(13): 1882–1887.
- [7] Beerliova, Z., Eberhard, F., Erlebach, T., Hall, A., Hoffman, M., Mihalák, M. (2006). Network discovery and verification. *IEEE J. Sel. Areas Commun.* 24(12): 2168–2181.
- [8] Bollobas, B., Mitsche, D., Pralat, P. (2013). Metric dimension for random graphs. *Electron. J. Combin.* 20(4): Paper 1.
- [9] Borgatti, S. P., Mehra, A., Brass, D. J., Labianca, G. (2009). Network analysis in the social sciences. *Science* 323(5916): 892–895.
- [10] Cáceres, J., Hernando, C., Mora, M., Pelayo, I. M., Puertas, M. L. (2009). On the metric dimension of infinite graphs. *Electron. Notes Discrete Math.* 35: 15–20.
- [11] Cáceres, J., Hernando, C., Mora, M., Puertas, M. L., Pelayo, I. M., Seara, C., Wood, D. R. (2005). On the metric dimension of some families of graphs. *Electron. Notes Discrete Math.* 22: 129–133.
- [12] Chartrand, G., Eroh, L., Johnson, M. A., Oellermann, O. (2000). Resolvability in graphs and the metric dimension of a graph. *Discrete Appl. Math.* 105: 99–113.
- [13] Chen, M.-S., Shin, K. G., Kandlur, D. D. (1990). Addressing, routing, and broadcasting in hexagonal mesh multiprocessors. *IEEE Trans. Comput.* 39(1): 10–18.
- [14] Decayeux, C., Seme, D. (2005). 3D hexagonal network: modeling, topological properties, addressing scheme, and optimal routing algorithm. *IEEE Trans. Parallel Distrib. Syst.* 16(9): 87–884.
- [15] Fehr, M., Gosselin, S., Oellermann, O. R. (2006). The metric dimension of Cayley digraphs. *Discrete Math.* 306(1): 31–41.
- [16] Feng, M., Wang, K. (2012). On the metric dimension of bilinear forms graphs. *Discrete Math.* 312(6): 1266–1268.
- [17] Fernau, H., Heggernes, P., Hof, P., Meister, D., Saei, R. (2015). Computing the metric dimension of chain graphs. *Inf. Process. Lett.* 115(9): 671–676.
- [18] Goddard, W. (2004). Statistic mastermind revisited. *J. Combin. Math. Combin. Comput.* 51: 215–220.
- [19] Grigoriou, C., Manuel, P., Miller, M., Rajan, B., Stephen, S. (2014). On the metric dimension of circulant and Harary graphs. *Appl. Math. Comp.* 248: 47–54.
- [20] Guo, J., Wang, K., Li, F. (2013). Metric dimension of symplectic dual polar graphs and symmetric bilinear forms graphs. *Discrete Math.* 313(2): 186–188.
- [21] Guo, J., Wang, K., Li, F. (2013). Metric dimension of some distance-regular graphs. *J. Comb. Optim.* 26: 190–197.
- [22] Hallaway, M., Kang, C. X., Yi, E. (2014). On metric dimension of permutation graphs. *J. Comb. Optim.* 28(4): 814–826.
- [23] Harary, F., Melter, R. A. (1976). On the metric dimension of a graph. *Ars Combin.* 2: 191–195.
- [24] Imran, M., Baig, A. Q., Bokhary, S. A. U. H., Javaid, I. (2012). On the metric dimension of circulant graphs. *Appl. Math. Lett.* 25(3): 320–325.
- [25] Imran, M., Siddiqui, H. M. A. (2016). Computing the metric dimension of convex polytopes generated by wheel related graphs. *Acta Math. Hung.* 149(1): 10–30.
- [26] Johnson, M. A. (1993). Structure-activity maps for visualizing the graph variables arising in drug design. *J. Biopharm. Stat.* 3: 203–236.
- [27] Khuller, S., Ragavachari, B., Rosenfield, A. (1996). Landmarks in graphs. *Discrete Appl. Math.* 70(3): 217–229.
- [28] Klein, D. J., Yi, E. (2012). A comparison on metric dimension of graphs, line graphs, line graphs of the subdivision graphs. *European J. Pure Appl. Math.* 5(3): 302–316.

- [29] Laihonon, T. (2016). The metric dimension for resolving several objects. *Inf. Process. Lett.* 116(11): 694–700.
- [30] Lam, P. C. B., Shiu, W. C., Chan, W. H. (1997). On the bandwidth of convex triangulation meshes. *Discrete Math.* 173: 285–289.
- [31] Liu, J. B., Nadeem, M. F., Siddiqui, M. A., Nazir, W. (2019). Computing metric dimension of certain families of toeplitz graphs. *IEEE Access* 7: 126734–126741.
- [32] Liu, J., Zafari, A., Zarei, H., Farooq, R. (2020). Metric dimension, minimal doubly resolving sets, and the strong metric dimension for jellyfish graph and cocktail party graph. *Complexity* 2020: 9407456.
- [33] Lovász, L. (2012). *Large Networks and Graph Limits*. Providence: American Mathematical Society.
- [34] Manuel, P., Abd-El-Barr, M. I., Rajasingh, I., Rajan, B. (2008). An efficient representation of benes networks and its applications. *J. Discrete Algorithms* 6(1): 11–19.
- [35] Manuel, P., Rajan, B., Rajasingh, I., Monica, M. C. (2006). Landmarks in torus networks. *J. Discrete Math. Sci. Crypt.* 9(2): 263–271.
- [36] Manuel, P., Rajan, B., Rajasingh, I., Monica, M. C. (2008). On minimum metric dimension of honeycomb networks. *J. Discrete Algorithms* 6(1): 20–27.
- [37] Marx, V. (2013). The big challenges of big data. *Nature* 498(7453): 255–260.
- [38] Melter, R. A., Tomescu, I. (1984). Metric bases in digital geometry. *Comp. Vision Graph. Image Process.* 25: 113–121.
- [39] Nocetti, F. G., Stojmenović, I., Zhang, J. (2002). Addressing and routing in hexagonal networks with applications for tracking mobile users and connection rerouting in cellular networks. *IEEE Trans. Parallel Distrib. Syst.* 13(9): 963–971.
- [40] Poisson, C., Zhang, P. (2002). The metric dimension of unicyclic graphs. *J. Combin. Math. Combin. Comput.* 40: 17–32.
- [41] Prabhu, S., Flora, T., Arulperumjothi, M. (2018). On independent resolving number of  $TiO_2[m, n]$  nanotubes. *J. Int. Fuzzy Syst.* 35: 6421–6425.
- [42] Rajan, B., Rajasingh, I., Monica, M. C., Manuel, P. (2008). Metric dimension of enhanced hypercube networks. *J. Combin. Math. Combin. Comput.* 67: 5–15.
- [43] Rajan, B., Rajasingh, I., Venugopal, P., Monica, M. C. (2014). Minimum metric dimension of illiac networks. *Ars Combin.* 117: 95–103.
- [44] Sebö, A., Tannier, E. (2004). On metric generators of graphs. *Math. Oper. Res.* 29(2): 383–393.
- [45] Shao, Z., Sheikholeslami, S. M., Wu, P., Liu, J. (2018). The metric dimension of some generalized Petersen graphs. *Discrete Dyn. Nat. Soc.* 10: 4531958.
- [46] Siddiqui, H. M. A., Imran, M. (2014). Computing the metric dimension of wheel related graphs. *Appl. Math. Comp.* 242: 624–632.
- [47] Slater, P. J. (1975). Leaves of trees. *Cong. Numer.* 14: 549–559.
- [48] Slater, P. J. (1988). Dominating and reference sets in a graph. *J. Math. Phys. Sci.* 22(4): 445–455.
- [49] Söderberg, S., Shapiro, H. S. (1963). A combinatory detection problem. *Amer. Math. Monthly* 70: 1066–1070.
- [50] Stojmenović, I. (1997). Honeycomb networks: Topological properties and communication algorithms. *IEEE Trans. Parallel Distrib. Syst.* 8(10): 1036–1042.
- [51] Tomescu, I., Javaid, I. (2007). On the metric dimension of the Jahangir graph. *Bull. Math. Soc. Sci. Math. Roumanie* 50: 371–376.
- [52] Vetrik, T. (2017). The metric dimension of circulant graphs. *Can. Math. Bull.* 60(1): 206–216.
- [53] Wasserman, S., Faust, K. (1994). *Social Network Analysis: Methods and Applications*. Cambridge: Cambridge University Press.