

INDECOMPOSABLE MATRICES DEFINING PLANE CUBICS

ANITA BUCKLEY

ABSTRACT. In this article we find all (decomposable and indecomposable) 6×6 linear determinantal representations of Weierstrass cubics. As a corollary we verify the Kippenhahn conjecture for M_6 .

1. INTRODUCTION

Let C be an irreducible curve in $\mathbb{C}\mathbb{P}^2$ defined by a polynomial $F(x, y, z)$ of degree 3. Every smooth cubic can be brought by a projective change of coordinates [15] into a Weierstrass form

$$F(x, y, z) = yz^2 - x(x - y)(x - \lambda y) = 0,$$

or equivalently

$$F(x, y, z) = -yz^2 + x^3 + \alpha xy^2 + \beta y^3 = 0,$$

for some $\lambda \neq 0, 1$ and $\alpha, \beta \in \mathbb{C}$.

We consider the following question. For given C find a linear matrix

$$A(x, y, z) = x A_x + y A_y + z A_z$$

such that

$$\det A(x, y, z) = c F(x, y, z)^r,$$

where $A_x, A_y, A_z \in M_{3r}$ and $0 \neq c \in \mathbb{C}$. Here M_{3r} is the algebra of all $3r \times 3r$ matrices over \mathbb{C} .

We call A a *determinantal representation* of C of order r . Two determinantal representations A and A' are *equivalent* if there exist $X, Y \in \text{GL}_{3r}(\mathbb{C})$ such that

$$A' = XAY.$$

We study self-adjoint representations $A = A^*$ modulo unitary equivalence $Y = X^*$ and skew-symmetric representations $A = -A^t$ under $Y = X^t$ equivalence. Obviously, equivalent determinantal representations define the same curve. *Pfaffian representation* is a representation of order 2 with all 6×6 matrices being skew-symmetric. Study of pfaffian representations is strongly related to determinantal representations: every 3×3 determinantal representation A induces a *decomposable pfaffian representation*

$$\begin{bmatrix} 0 & A \\ -A^t & 0 \end{bmatrix}.$$

Note that the equivalence relation is well defined since

$$\begin{bmatrix} 0 & XAY \\ -(XAY)^t & 0 \end{bmatrix} = \begin{bmatrix} X & 0 \\ 0 & Y^t \end{bmatrix} \begin{bmatrix} 0 & A \\ -A^t & 0 \end{bmatrix} \begin{bmatrix} X^t & 0 \\ 0 & Y \end{bmatrix}.$$

The history of determinantal representations of order 1 goes back to the middle of the 19th century [10], [21]. Dickson [7] classified hypersurfaces that can be represented as linear determinants. In the last two decades determinantal representations

of hypersurfaces again became extremely popular due to their use in semidefinite programming [24]. Semidefinite programming feasibly uses convex sets determined by linear matrix inequalities

$$\left\{ (x_1, \dots, x_d) \in \mathbb{R}^d : A_0 + x_1 A_1 + \dots + x_d A_d \geq 0 \right\},$$

where A_0, \dots, A_d are real symmetric or complex hermitian matrices. A natural question in this framework is whether a positive integer power of the polynomial has a determinantal representations.

All modern treatments of determinantal representations of order 1 involve [6], the famous 1 – 1 correspondence between determinantal representations of hypersurfaces and line bundles (i.e., points on the Jacobian variety). Analogously, there is a one to one correspondence between linear pfaffian representations (up to equivalence) and rank 2 vector bundles (up to isomorphism) with certain properties. These well known results are summed up in Beauville [3].

Elliptic curves are of tame representation type according to Atiyah [1]. In particular, on a given cubic curve the number of indecomposable ACM bundles of rank r with trivial determinant equals to the number of r -torsion points. Recently, Ravindra and Tripathi [17] used these indecomposable vector bundles to prove the existence of indecomposable determinantal representations of order greater than two.

Vinnikov [22] explicitly parametrised 3×3 determinantal representations of a Weierstrass cubic by the affine points on the same cubic. In this paper we classify up to equivalence all linear pfaffian representations of a Weierstrass cubic: the decomposable ones are parametrized by the affine points on the cubic, additionally there are three indecomposable representations arising from nontrivial extensions of even theta characteristics (i.e., line bundles corresponding to 2-torsion points). The classification is based on Lancaster-Rodman canonical forms of matrix pairs [13]. More generally, any linear representation of order 2 is equivalent to either a block linear matrix or its cokernel is a degree 0 indecomposable rank 2 bundle of Atiyah.

The main result of this paper is Theorem 3.2 containing the construction of the three indecomposable pfaffian representations for a Weierstrass cubic. We outline similar constructions for indecomposable determinantal representations of order $r \geq 2$ corresponding to indecomposable vector bundles of rank r . These computations are an appendix to the paper of Ravindra and Tripathi [17]. As a corollary we verify the Kippenhahn conjecture for M_6 .

The author wishes to thank G. V. Ravindra for his guidance and his patience, and to I. Klep for pointing out the connection between our constructions and the conjecture of Kippenhahn.

2. DETERMINANTAL REPRESENTATIONS OF VINNIKOV

Vinnikov [22] found an explicit one to one correspondence between the 3×3 determinantal representations (up to equivalence) of C and the points on an affine piece of C :

Lemma 2.1 ([22]). *Every smooth cubic in \mathbb{P}^2 can be brought into the Weierstrass form*

$$F(x_0, x_1, x_2) = -x_1 x_2^2 + x_0^3 + \alpha x_0 x_1^2 + \beta x_1^3.$$

A complete set of determinantal representations of F is

$$(1) \quad x_0 \text{Id} + x_2 \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} + x_1 \begin{bmatrix} \frac{t}{2} & s & \alpha + \frac{3}{4}t^2 \\ 0 & -t & -s \\ -1 & 0 & \frac{t}{2} \end{bmatrix},$$

where $s^2 = t^3 + \alpha t + \beta$. Note that the last equation is exactly the affine part $F(t, 1, s)$.

We briefly repeat the proof as we will use similar ideas for 6×6 skew-symmetric determinantal representations.

Proof. Let $A(x_0, x_1, x_2) = x_0 A_0 + x_2 A_2 + x_1 A_1$ be a representation of $F(x_0, x_1, x_2)$. First we show that it is equivalent to a representation with

$$A_0 = \text{Id} \quad \text{and} \quad A_2 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}.$$

Observe that A_0 is invertible since $\det A(1, 0, 0) \neq 0$. We can multiply $A(x_0, x_1, x_2)$ by A_0^{-1} to obtain an equivalent representation with $A_0 = \text{Id}$. The characteristic polynomial of A_2 equals $\det A(x_0, 0, -1) = x_0^3$ which implies that A_2 is nilpotent. The nonzero term $x_1 x_2^2$ in F determines the order of nilpotency. We remark that this is a Lancaster–Rodman canonical form for real matrix pairs [13]. Further GL_3 action from left and right which preserves this canonical form (the first two matrices in the determinantal representation) reduces A_1 to the above. \square

The above is an implementation of the classic Cook–Thomas correspondence [6]:

Proposition 2.2 ([3], Proposition 3.1.). *Let \mathcal{L} be a line bundle of degree 0 on C with $H^0(C, \mathcal{L}) = 0$. Then there exists a 3×3 linear matrix A such that $F = \det A$ and*

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^2}(-2)^3 \xrightarrow{A} \mathcal{O}_{\mathbb{P}^2}(-1)^3 \rightarrow \mathcal{L} \rightarrow 0.$$

Conversely, let A be a 3×3 linear matrix with $F = \det A$. Then the cokernel of A is a line bundle \mathcal{L} on C of degree 0 with $H^0(C, \mathcal{L}) = 0$.

The explicit formula in Lemma 2.1 is due to the fact that the set of line bundles of degree 0 with no global sections equals to $JC \setminus \mathcal{O}_C$, where $JC = \text{Pic}^0 C$ is the Jacobian of C . Recall that JC equals to the curve itself. Therefore $JC \setminus \mathcal{O}_C$ can be parametrised by the affine points on the curve C .

Observe that if the cokernel of A is \mathcal{L} , the cokernel of A^t is \mathcal{L}^{-1} . This can also be seen directly from Lemma 2.1, just multiply (1) by anti-identity. Then

$$(2) \quad A = x_0 \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} + x_1 \begin{bmatrix} \alpha + \frac{3}{4}t^2 & s & \frac{t}{2} \\ -s & -t & 0 \\ \frac{t}{2} & 0 & -1 \end{bmatrix}$$

and A^t correspond to the inverse points (t, s) and $(t, -s)$ respectively on the affine part $s^2 = t^3 + \alpha t + \beta$ of $JC = C$. We can also conclude that the above A is symmetric if and only if $s = 0$. This implies that on C there are three symmetric determinantal representations corresponding to three 2-torsion points $\mathcal{L} \cong \mathcal{L}^{-1}$ on JC . In the affine coordinates these three points are $(t, 0)$, where $t^3 + \alpha t + \beta = 0$.

3. PFAFFIAN REPRESENTATIONS

In this section we will implement the one to one correspondence between linear pfaffian representations and rank 2 vector bundles with certain properties, applying similar methods as in the proof of Lemma 2.1. The correspondence is described in

Proposition 3.1 ([3], Proposition 5.1.). *Let \mathcal{E} be a rank 2 bundle on C with $\det \mathcal{E} = \mathcal{O}_C$ and $H^0(C, \mathcal{E}) = 0$. Then there exists a 6×6 linear skew-symmetric matrix A such that $F = \text{Pf } A$ and*

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^2}(-2)^6 \xrightarrow{A} \mathcal{O}_{\mathbb{P}^2}(-1)^6 \rightarrow \mathcal{E} \rightarrow 0.$$

Conversely, let A be a 6×6 linear skew-symmetric matrix with $F = \text{Pf } A$. Then the cokernel of A is a rank 2 bundle \mathcal{E} on C with the above properties .

Pfaffian representations are equivalent under the action

$$A \mapsto P \cdot A \cdot P^t,$$

where P is an invertible 6×6 constant matrix. By suitable choices of P it is possible to reduce the number of parameters in A . In other words, we will reduce the number of equivalent representations in each equivalence class. The proof of Theorem 3.2 outlines an algorithm for finding all pfaffian representations (up to equivalence) of

$$C = \{(x, y, z) \in \mathbb{P}^2 : yz^2 - x(x - y)(x - \lambda y) = 0\}.$$

For the sake of clearer notation we always write just the upper triangle of skew-symmetric matrices.

Theorem 3.2. *Let C be a smooth cubic in the Weierstrass form*

$$F(x, y, z) = yz^2 - x(x - y)(x - \lambda y).$$

A complete set of pfaffian representations of F consists up to equivalence of three indecomposable representations and for the whole affine curve of decomposable representations:

$$x \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ & 0 & 0 & 0 & 1 & 0 \\ & & 0 & 1 & 0 & 0 \\ & & & 0 & 0 & 0 \\ & & & & 0 & 0 \\ & & & & & 0 \end{bmatrix} + z \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ & 0 & 0 & 1 & 0 & 0 \\ & & 0 & 0 & 0 & 0 \\ & & & 0 & 0 & 0 \\ & & & & 0 & 0 \\ & & & & & 0 \end{bmatrix} + y \begin{bmatrix} 0 & 1 & 0 & \frac{3t^2 - 2t(1+\lambda) - (1-\lambda)^2}{4} & 0 & \frac{t-1-\lambda}{2} \\ & 0 & 0 & 0 & -t & 0 \\ & & 0 & \frac{t-1-\lambda}{2} & 0 & -1 \\ & & & 0 & 0 & 0 \\ & & & & 0 & 0 \\ & & & & & 0 \end{bmatrix} \quad \text{for } t = 0, 1, \lambda$$

and

$$x \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ & 0 & 0 & 0 & 1 & 0 \\ & & 0 & 1 & 0 & 0 \\ & & & 0 & 0 & 0 \\ & & & & 0 & 0 \\ & & & & & 0 \end{bmatrix} + z \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ & 0 & 0 & 1 & 0 & 0 \\ & & 0 & 0 & 0 & 0 \\ & & & 0 & 0 & 0 \\ & & & & 0 & 0 \\ & & & & & 0 \end{bmatrix} + y \begin{bmatrix} 0 & 0 & 0 & \frac{3t^2 - 2t(1+\lambda) - (1-\lambda)^2}{4} & s & \frac{t-1-\lambda}{2} \\ & 0 & 0 & -s & -t & 0 \\ & & 0 & \frac{t-1-\lambda}{2} & 0 & -1 \\ & & & 0 & 0 & 0 \\ & & & & 0 & 0 \\ & & & & & 0 \end{bmatrix},$$

where $s^2 = t(t-1)(t-\lambda)$. Note that the last equation is exactly the affine part $F(t, 1, s)$.

The proof will be based on Lancaster–Rodman canonical forms of matrix pairs [13]. Let $A = xA_x + zA_z + yA_y$ be a pfaffian representation of C . Observe that A_x is invertible and A_z nilpotent since C is defined by $\text{Pf } A$ and contains x^3 term and no z^3 term. Moreover, yz^2 determines the order of nilpotency. This determines the

unique skew-symmetric canonical form [13, Theorem 5.1] for the first two matrices. In other words, every pfaffian representation of C can be put into the following form

$$x \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} + z \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} + y \begin{bmatrix} 0 & c_{12} & c_{13} & c_{14} & c_{15} & c_{16} \\ 0 & c_{23} & c_{24} & c_{25} & c_{26} & 0 \\ 0 & c_{34} & c_{35} & c_{36} & 0 & 0 \\ 0 & c_{45} & c_{46} & 0 & 0 & 0 \\ 0 & c_{56} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Since Pf A defines the equation of C , we get

$$\begin{aligned} c_{36} &= -1, \\ c_{26} &= -c_{35}, \\ c_{25} &= -1 - \lambda - c_{16} - c_{34}, \\ c_{14} &= c_{16} + c_{16}^2 + c_{34} + c_{16}c_{34} + c_{34}^2 + 2c_{24}c_{35} + c_{16}c_{35}^2 - c_{34}c_{35}^2 - \\ &\quad c_{23}c_{45} - c_{13}c_{46} + c_{23}c_{35}c_{46} - c_{12}c_{56} + c_{13}c_{35}c_{56} + \lambda(1 + c_{16} + c_{34}), \\ c_{15} &= -c_{24} - c_{16}c_{35} + c_{34}c_{35} - c_{23}c_{46} - c_{13}c_{56}. \end{aligned}$$

There are 15–5 parameters c_{ij} left in the representation. Additionally, the coefficient at y^3 equals $c_{14}c_{26}c_{35} - c_{14}c_{25}c_{36} - c_{13}c_{26}c_{45} + c_{12}c_{36}c_{45} + c_{16}(c_{25}c_{34} - c_{24}c_{35} + c_{23}c_{45}) + c_{13}c_{25}c_{46} - c_{12}c_{35}c_{46} - c_{15}(c_{26}c_{34} - c_{24}c_{36} + c_{23}c_{46}) + c_{14}c_{23}c_{56} - c_{13}c_{24}c_{56} + c_{12}c_{34}c_{56} = 0$.

Lemma 3.3. *The action $A \mapsto P \cdot A \cdot P^t$ preserves the canonical form of the first two matrices in the representation if and only if P equals*

$$\begin{bmatrix} P_1 & P_2 \\ P_3 & P_1^{-1} + P_3P_1^{-1}P_2 \end{bmatrix} \text{ or } \begin{bmatrix} P_2 & P_1 \\ -P_1^{-1} + P_3P_1^{-1}P_2 & P_3 \end{bmatrix}$$

where P_1 is invertible and P_i are of the form

$$\begin{bmatrix} p_{i1} & p_{i2} & p_{i3} \\ 0 & p_{i1} & p_{i2} \\ 0 & 0 & p_{i1} \end{bmatrix}, \quad i = 1, 2, 3.$$

Proof. Denote

$$I = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \quad \text{and} \quad N = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

We will need the following obvious observation, which can be proved directly by comparing matrix elements:

Let Y, Y' be 6×6 matrices for which

$$Y \cdot \begin{bmatrix} 0 & I \\ -I & 0 \end{bmatrix} = \begin{bmatrix} 0 & I \\ -I & 0 \end{bmatrix} \cdot Y' \quad \text{and} \quad Y \cdot \begin{bmatrix} 0 & N \\ -N & 0 \end{bmatrix} = \begin{bmatrix} 0 & N \\ -N & 0 \end{bmatrix} \cdot Y' \quad \text{hold.}$$

Then $Y = \begin{bmatrix} Y_1 & Y_2 \\ Y_3 & Y_4 \end{bmatrix}$ and $Y' = \begin{bmatrix} Y_4^t & -Y_3^t \\ -Y_2^t & Y_1^t \end{bmatrix}$, where

$$Y_i = \begin{bmatrix} y_{i1} & y_{i2} & y_{i3} \\ 0 & y_{i1} & y_{i2} \\ 0 & 0 & y_{i1} \end{bmatrix}, \quad i = 1, 2, 3, 4.$$

We call the specific form of the above Toeplitz matrices ” Δ form”.

Now we can find all invertible

$$P = \begin{bmatrix} P_1 & P_2 \\ P_3 & P_4 \end{bmatrix}$$

that satisfy

$$P \cdot \begin{bmatrix} 0 & I \\ -I & 0 \end{bmatrix} \cdot P^t = \begin{bmatrix} 0 & I \\ -I & 0 \end{bmatrix} \quad \text{and} \quad P \cdot \begin{bmatrix} 0 & N \\ -N & 0 \end{bmatrix} \cdot P^t = \begin{bmatrix} 0 & N \\ -N & 0 \end{bmatrix}.$$

By the above observation all P_i 's are of Δ form. Moreover,

$$\begin{aligned} p_{11}p_{41} - p_{21}p_{31} &= 1, \\ p_{22}p_{31} + p_{21}p_{32} - p_{12}p_{41} - p_{11}p_{42} &= 0, \\ p_{23}p_{31} + p_{22}p_{32} + p_{21}p_{33} - p_{13}p_{41} - p_{12}p_{42} - p_{11}p_{43} &= 0. \end{aligned}$$

In other words, if P_1 is invertible then $P_4 = P_1^{-1} + P_3P_1^{-1}P_2$. The same way we see that $P_3 = -P_2^{-1} + P_1P_2^{-1}P_4$ when P_2 is invertible.

Since P is invertible and consists of Δ blocks, at least one of P_1, P_2 is also invertible. Note that

$$\begin{bmatrix} P_1 & P_2 \\ P_3 & P_4 \end{bmatrix} \cdot \begin{bmatrix} 0 & -\text{Id} \\ \text{Id} & 0 \end{bmatrix} = \begin{bmatrix} P_2 & -P_1 \\ P_4 & -P_3 \end{bmatrix}$$

exchanges P_1 and P_2 which finishes the proof. \square

The action of Lemma 3.3 enables us to reduce the number of parameters c_{ij} . We can choose such P that its action eliminates

$$(3) \quad c_{13} = c_{23} = c_{46} = c_{56} = 0, \quad c_{35} = 0 \quad \text{and} \quad c_{16} = c_{34}.$$

Indeed, if we choose $p_{11} = 1$, the above conditions determine p_{12}, p_{13} and $p_{22}, p_{23}, p_{32}, p_{33}$:

$$\begin{aligned} p_{32} &\rightarrow c_{56} - c_{35}p_{31} + c_{56}p_{31}p_{21}, \\ p_{22} &\rightarrow -c_{23} + c_{35}p_{21}, \\ p_{33} &\rightarrow \frac{1}{2}((c_{16} - c_{34} + c_{35}^2 - c_{23}c_{56})p_{31} + 2c_{46}(1 + p_{31}p_{21})), \\ p_{23} &\rightarrow \frac{1}{2}(-2c_{13} + (-c_{16} + c_{34} + c_{35}^2 - c_{23}c_{56})p_{21}), \\ p_{12} &\rightarrow -c_{35} + c_{56}p_{21}, \\ p_{13} &\rightarrow \frac{1}{2}(c_{16} - c_{34} + c_{35}^2 - c_{23}c_{56} + 2c_{46}p_{21}). \end{aligned}$$

The relations among c_{ij} then simplify to:

$$\begin{aligned} c_{14} &= 3c_{16}^2 + \lambda + 2c_{16}(1 + \lambda), \\ c_{24} &= -c_{15}, \\ (4) \quad 0 &= c_{15}^2 - 8c_{16}^3 - c_{12}c_{45} - \lambda - \lambda^2 - 8c_{16}^2(1 + \lambda) - 2c_{16}(1 + 3\lambda + \lambda^2). \end{aligned}$$

which leaves us with 4 parameters $c_{12}, c_{45}, c_{15}, c_{16}$ and equation (4) connecting them:

$$\begin{bmatrix} 0 & \mathbf{c}_{12} & 0 & c_{14} & \mathbf{c}_{15} & \mathbf{c}_{16} \\ & 0 & 0 & -c_{15} & -1-\lambda-2c_{16} & 0 \\ & & 0 & c_{16} & 0 & -1 \\ & & & 0 & \mathbf{c}_{45} & 0 \\ & & & & 0 & 0 \\ & & & & & 0 \end{bmatrix}.$$

It is easy to check that $A \mapsto P \cdot A \cdot P^t$ from Lemma 3.3 preserves all zeros and -1 in the above matrix if and only if

$$P_i = \begin{bmatrix} p_{i1} & 0 & 0 \\ 0 & p_{i1} & 0 \\ 0 & 0 & p_{i1} \end{bmatrix} \quad \text{for } i = 1, 2, 3, 4, \quad \text{together with } p_{11}p_{41} - p_{21}p_{31} = 1.$$

We can use this "diagonal" action to make $c_{45} = 0$ by choosing appropriate p_{41} like in (3). When $c_{15} \neq 0$ we can furthermore make $c_{12} = 0$ by $p_{11} = p_{41} = 1, p_{31} = 0, p_{21} = -c_{12}/2c_{15}$. The only case left to consider is $c_{15} = 0$. The action which keeps $c_{45} = 0$ maps $c_{12} \mapsto c_{12}p_{11}^2$ where $p_{11}p_{41} = 1$ and $p_{31} = 0$. Thus either $c_{12} = 0$ or we can make $c_{12} = 1$.

In order to simplify notations even further, we introduce parameters t and s by $c_{16} = \frac{1}{2}(t - 1 - \lambda)$ and $c_{15} = s$. When $c_{45} = c_{12} = 0$ the matrix becomes

$$\begin{bmatrix} 0 & 0 & 0 & \frac{3t^2 - 2t(1+\lambda) - (1-\lambda)^2}{4} & s & \frac{t-1-\lambda}{2} \\ & 0 & 0 & -s & -t & 0 \\ & & 0 & \frac{t-1-\lambda}{2} & 0 & -1 \\ & & & 0 & 0 & 0 \\ & & & & 0 & 0 \\ & & & & & 0 \end{bmatrix}$$

and relation (4) in the new parameters equals $s^2 - t(t-1)(t-\lambda) = 0$.

Additionally we get

$$\begin{bmatrix} 0 & 1 & 0 & \frac{3t^2 - 2t(1+\lambda) - (1-\lambda)^2}{4} & 0 & \frac{t-1-\lambda}{2} \\ & 0 & 0 & 0 & -t & 0 \\ & & 0 & \frac{t-1-\lambda}{2} & 0 & -1 \\ & & & 0 & 0 & 0 \\ & & & & 0 & 0 \\ & & & & & 0 \end{bmatrix}$$

where t is one of the three solutions of $0 = -t(t-1)(t-\lambda)$.

Remark 3.4. The representations in Theorem 3.2 are non-equivalent to each other, since they are not connected by the action $A \mapsto P \cdot A \cdot P^t$.

4. A REMARK ABOUT THE MODULI SPACE OF RANK 2 BUNDLES

Rank 2 bundles with trivial determinant and no sections lie in the open set

$$M_C(2, \mathcal{O}_C) \setminus \Theta_{2, \mathcal{O}_C},$$

where $M_C(2, \mathcal{O}_C)$ is the moduli space of semistable rank 2 bundles with determinant \mathcal{O}_C and $\Theta_{2, \mathcal{O}_C} = \{\mathcal{E} \in M_C(2, \mathcal{O}_C) : h^0(C, \mathcal{E}) \neq 0\}$. In [3, § 4] and [5, § 6] we can find that there are no stable bundles in $M_C(2, \mathcal{O}_C)$ and the unstable part consists of decomposable vector bundles of the form $\mathcal{L} \oplus \mathcal{L}^{-1}$ for \mathcal{L} in the Jacobian JC . Moreover, $\Theta_{2, \mathcal{O}_C} = \{\mathcal{O}_C \oplus \mathcal{O}_C\}$. This yields a 1-1 correspondence between the points in $M_C(2, \mathcal{O}_C) \setminus \Theta_{2, \mathcal{O}_C}$ and the open subset of Kummer variety

$$(JC \setminus \{\mathcal{O}_C\}) / \equiv,$$

where \equiv is the involution $\mathcal{L} \mapsto \mathcal{L}^{-1}$.

Let A and $-A^t$ be 3×3 determinantal representations with cokernels \mathcal{L} and \mathcal{L}^{-1} respectively, like in Section 2. Obviously $\mathcal{L} \oplus \mathcal{L}^{-1}$ and $\mathcal{L}^{-1} \oplus \mathcal{L}$ are isomorphic rank 2 vector bundles. This is aligned with the corresponding decomposable pffian representations. Indeed, even though A and $-A^t$ are not necessarily equivalent determinantal representations,

$$\begin{bmatrix} 0 & A \\ -A^t & 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 0 & -A^t \\ A & 0 \end{bmatrix}$$

are equivalent pfaffian representations since

$$\begin{bmatrix} 0 & \text{Id} \\ \text{Id} & 0 \end{bmatrix} \begin{bmatrix} 0 & A \\ -A^t & 0 \end{bmatrix} \begin{bmatrix} 0 & \text{Id} \\ \text{Id} & 0 \end{bmatrix} = \begin{bmatrix} 0 & -A^t \\ A & 0 \end{bmatrix}.$$

Remark 4.1. Each point in the the moduli space $M_C(2, \mathcal{O}_C)$ corresponds to a decomposable bundle which induces a decomposable pfaffian representation. However, this does not imply that every rank 2 bundle on C is decomposable. This is because the moduli space consists of S-equivalence classes rather than isomorphic bundles.

Take for example the cokernel of a symmetric 3×3 representation; one of the three cases where $s = 0$ in (2). Then \mathcal{L} is a 2-torsion point on JC , or in other words an even theta characteristic. For this line bundle $\mathcal{L} \cong \mathcal{L}^{-1}$, the direct sum $\mathcal{L} \oplus \mathcal{L}$ and the non-trivial extension of \mathcal{L} by \mathcal{L} represent the same point in the moduli space. These two bundles are clearly not isomorphic and can be realized as cokernels of matrices from Theorem 3.2,

$$x \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} + z \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} + y \begin{bmatrix} 0 & 0 & 0 & \frac{3t^2 - 2t(1+\lambda) - (1-\lambda)^2}{4} & 0 & \frac{t-1-\lambda}{2} \\ 0 & 0 & 0 & 0 & -t & 0 \\ 0 & 0 & \frac{t-1-\lambda}{2} & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

and

$$(5) \quad x \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} + z \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} + y \begin{bmatrix} 0 & 1 & 0 & \frac{3t^2 - 2t(1+\lambda) - (1-\lambda)^2}{4} & 0 & \frac{t-1-\lambda}{2} \\ 0 & 0 & 0 & 0 & -t & 0 \\ 0 & 0 & \frac{t-1-\lambda}{2} & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix},$$

for t satisfying $0 = t(t-1)(t-\lambda)$.

5. DETERMINANTAL REPRESENTATIONS OF ORDER ≥ 2

Consider a linear matrix A with $\det A = F^r$, where F defines C . Since C is smooth, we will prove that the cokernel of A is a vector bundle of rank r . Indeed, by [3, Theorem A] the cokernel is an arithmetically Cohen-Macaulay (ACM) sheaf. Over a regular scheme any Cohen-Macaulay sheaf is locally free, thus it is a vector bundle. Furthermore, localizing at the generic point of C , we can locally write A as a diagonal matrix $[F, \dots, F, 0, \dots, 0]$, where F occurs r times. The same result was obtained in [8] and [2] using purely algebraic methods for matrix factorizations of polynomials.

Atiyah [1] classified indecomposable vector bundles on C in the following theorem.

Theorem 5.1 ([1], Theorem 5, Corollary 1). *Let C be an elliptic curve.*

- (i) *For any $r > 0$ there exists a unique (up to isomorphism) indecomposable vector bundle \mathcal{F}_r with $h^0(\mathcal{F}_r) = 1$. Moreover, \mathcal{F}_r is defined inductively by the short exact sequence*

$$0 \rightarrow \mathcal{O}_C \rightarrow \mathcal{F}_{r+1} \rightarrow \mathcal{F}_r \rightarrow 0, \text{ where } \mathcal{F}_1 = \mathcal{O}_C.$$

- (ii) *For any indecomposable rank r bundle \mathcal{E} of degree 0, there exists a line bundle \mathcal{L} such that $\mathcal{E} \cong \mathcal{F}_r \otimes \mathcal{L}$ and $\mathcal{L}^{\otimes r} = \det \mathcal{E}$.*
- (iii) *\mathcal{F}_r is self-dual, $\mathcal{F}_r \cong \mathcal{F}_r^\vee$.*

By knowing all possible cokernels of determinantal representations, we can find (up to equivalence) all determinantal representations of F [3, Proposition 1.11.]. For $r = 2$ this 1–1 correspondence gives the following corollary.

Corollary 5.2. *Let A be a linear 6×6 matrix with $\det A = F^2$. Then A is either equivalent to a block matrix*

$$(6) \quad \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix},$$

or the cokernel of A equals to a nontrivial extension of a degree 0 line bundle $\mathcal{L} \neq \mathcal{O}_C$ by itself. Recall that such \mathcal{L} is the cokernel of some 3×3 determinantal representation of C .

Proof. Denote the cokernel of A by \mathcal{E} . Consider the short exact sequence

$$0 \rightarrow \mathcal{L}_1 \rightarrow \mathcal{E} \rightarrow \mathcal{L}_2 \rightarrow 0,$$

where the line bundles $\mathcal{L}_1, \mathcal{L}_2$ are cokernels of 3×3 determinantal representations of C . By the Riemann-Roch formula all $\mathcal{L}_1, \mathcal{L}_2$ and \mathcal{E} have degree 0. Then \mathcal{E} either splits and defines (6), or is a nontrivial extension if and only if

$$\mathrm{Ext}^1(\mathcal{L}_2, \mathcal{L}_1) \cong \mathrm{Ext}^1(\mathcal{L}_2 \otimes \mathcal{L}_1^{-1}, \mathcal{O}_C) \cong H^1(C, \mathcal{L}_2 \otimes \mathcal{L}_1^{-1}) \cong H^0(C, \mathcal{L}_1 \otimes \mathcal{L}_2^{-1}) \neq 0.$$

The only line bundle on C with nontrivial group of sections is \mathcal{O}_C . This means that \mathcal{E} is indecomposable if and only if $\mathcal{L}_1 \otimes \mathcal{L}_2^{-1} = \mathcal{O}_C$. This is consistent with Atiyah's Theorem 5.1 where every indecomposable rank 2 bundle of degree 0 fits into the exact sequence

$$(7) \quad 0 \rightarrow \mathcal{L} \rightarrow \mathcal{E} \rightarrow \mathcal{L} \rightarrow 0$$

for some line bundle \mathcal{L} with $\det \mathcal{E} = \mathcal{L}^{\otimes 2}$. By [3, Proposition 1.11.], given a coherent sheaf \mathcal{E} on \mathbb{P}^2 , there exists an exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^2}(-2)^6 \xrightarrow{A} \mathcal{O}_{\mathbb{P}^2}(-1)^6 \rightarrow \mathcal{E} \rightarrow 0$$

if and only if $H^0(\mathbb{P}^2, \mathcal{E}) = H^1(\mathbb{P}^2, \mathcal{E}) = 0$. This gives the 1–1 correspondence between indecomposable rank two bundles \mathcal{E} with $\det \mathcal{E} = \mathcal{L}^{\otimes 2}$ for some $\mathcal{L} \neq \mathcal{O}_C$ of degree 0, and indecomposable determinantal representations A with $\det A = F^2$ and $\mathrm{Coker} A = \mathcal{E}$. Every $\mathcal{L} \neq \mathcal{O}_C$ of degree 0 from (7) is by Proposition 2.2 a cokernel of some 3×3 determinantal representation of C . Clearly, $H^0(C, \mathcal{L}) = 0$ and by Riemann-Roch $H^1(C, \mathcal{L}) = 0$, which implies $H^0(C, \mathcal{E}) = H^1(C, \mathcal{E}) = 0$. On the other hand, minimal free resolution is unique up to isomorphism by [9]. \square

Consider again A with $\det A = F^r$ and cokernel \mathcal{E} . In [3, Theorem B] we find that when \mathcal{E} is additionally equipped with an ϵ -symmetric ($\epsilon = (-1)^{r-1}$) invertible form

$$\mathcal{E} \times \mathcal{E} \rightarrow \mathcal{O}_C(\alpha) \text{ for some } \alpha \in \mathbb{Z},$$

then A can be taken to be ϵ -symmetric. From here it follows that there are exactly three symmetric 3×3 determinantal representations of C with cokernels $\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3$, the three even theta characteristics (i.e., 2-torsion points in JC). Indeed, these are the only nontrivial line bundles of degree 0 with $\mathcal{L}^{\otimes 2} \cong \mathcal{O}_C$. In rank 2 case, Theorem 5.1 shows that the only indecomposable rank 2 bundles on C with determinant \mathcal{O}_C are the nontrivial extensions of theta characteristics \mathcal{L}_i by themselves. Take an indecomposable pfaffian representation from Theorem 3.2. Its cokernel is indeed the nontrivial extension of a 2-torsion point $\mathcal{L}_i = \mathcal{L}_i^{-1}$. The three non-block matrices in Theorem 3.2 are by the above the only indecomposable skew-symmetric matrices with determinant F^2 . For $r = 3$ Ravindra and Tripathi [17] proved the

existence of eight indecomposable 9×9 determinantal representations of C . The corresponding indecomposable cokernels are extensions of the nontrivial 3-torsion points on JC with themselves (i.e., the eight flexes on the affine Weierstrass cubic). In order to explicitly construct these determinantal representations, we would need to repeat the proof of Theorem 3.2 for 9×9 matrices. We invite the interested reader to compute them as "extensions" of block matrices with three nonzero blocks A ; here A comes from Lemma 2.1 with (s, t) being one of the eight flexes on the affine Weierstrass cubic.

As another interesting corollary of Theorem 3.2 we are able to verify the conjecture of Kippenhahn [11] for $M_6(\mathbb{C})$, the algebra of 6×6 matrices. Shapiro [19] and [20] showed that the conjecture holds for $n \leq 5$ and for $n = 6$ in the case that the minimal polynomial is cubic. On the other hand, Waterhouse [25] presented a pair of 6×6 matrices H, K that generate $M_6(\mathbb{C})$ such that $\det(xH + yK - z\text{Id})$ has repeated linear factors, thus disproving the general form of Kippenhahn conjecture for $n = 6$. Another class of counterexamples for $n = 6$ has been constructed in [14]. Laffey [12] constructed a counterexample for $n = 8$ with quartic minimal polynomial. The following corollary shows that there is no similar counterexample with $n = 6$.

Corollary 5.3. *The conjecture of Kippenhahn as stated here is true for $n = 6$.*

Let H, K be 6×6 complex Hermitian matrices and F a homogeneous polynomial defining a smooth cubic, such that

$$\det(xH + yK - z\text{Id}) = F(x, y, z)^2.$$

Then H and K are simultaneously unitarily similar to direct sums. This means that there exists an unitary matrix U and matrices $H_1, H_2, K_1, K_2 \in M_3(\mathbb{C})$ such that

$$UHU^* = \begin{bmatrix} H_1 & 0 \\ 0 & H_2 \end{bmatrix} \text{ and } UKU^* = \begin{bmatrix} K_1 & 0 \\ 0 & K_2 \end{bmatrix}.$$

Remark 5.4. Note that F in Corollary 5.3 defines a real cubic curve and that $z\text{Id} - xH - yK$ is a definite determinantal representation of F . In the terminology of linear matrix inequalities, F is a *real zero polynomial* and $(0, 0)$ lies inside the convex set of points $\{(x, y) \in \mathbb{R}^2 : \text{Id} - xH - yK \geq 0\}$ called *spectrahedron*. Spectrahedron is bounded by the compact part of the curve. For more constructions of definite determinantal representations of polynomials we refer the reader to [16] and [18] and the references therein.

Proof. Every smooth real cubic can be brought into a Weierstrass form by a real change of coordinates

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} \mapsto P \begin{bmatrix} x \\ y \\ z \end{bmatrix}, \text{ for some } P \in \text{GL}_3(\mathbb{R}).$$

In the new coordinates we get

$$\det(xA_x + yA_y + zA_z) = (-yz^2 + x^3 + \alpha xy^2 + \beta y^3)^2, \text{ where } \alpha, \beta \in \mathbb{R}.$$

Note that A_x, A_y, A_z are real linear combinations of H, K, Id and therefore Hermitian. Since $z\text{Id} - xH - yK$ is definite, $A = xA_x + yA_y + zA_z$ is also definite. We showed that the cokernel of A is a rank 2 bundle. Assume first that the cokernel is decomposable $\mathcal{L}_1 \oplus \mathcal{L}_2$. Then A is equivalent to a block matrix $\begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix}$, where \mathcal{L}_i is the cokernel of A_i . In particular, each line bundle \mathcal{L}_i satisfies the conditions

in [22, Theorem 7], therefore A_i can be brought by [23, Theorem 8] into one of the two self-adjoint forms

$$(8) \quad \pm \left(x \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} + z \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} + y \begin{bmatrix} \alpha + \frac{3}{4}t_i^2 & i s_i & \frac{t_i}{2} \\ -i s_i & -t_i & 0 \\ \frac{t_i}{2} & 0 & -1 \end{bmatrix} \right),$$

where $(s_i, t_i) \in \mathbb{R}^2$ satisfy $-s_i^2 = t_i^3 + \alpha t_i + \beta$. Moreover, A_i is unitarily equivalent to one of the above forms. It was shown in [23], that definite self-adjoint determinantal representations of C are exactly those corresponding to the points (s, t) in the compact part of $C(\mathbb{R})$. This self-adjoint canonical form can be obtained also directly, using Lancaster–Rodman canonical forms for real matrix pairs [13], like we did in Section 2. Indeed, just replace s by $i s$ in (2).

Next we assume that the cokernel \mathcal{E} of A is indecomposable. In Corollary 5.2 we showed that $\det \mathcal{E} = \mathcal{L}^{\otimes 2}$ for some line bundle $\mathcal{L} \neq \mathcal{O}_C$ of degree 0. Since A is self-adjoint, \mathcal{L} is isomorphic to the cokernel of a self-adjoint representation in (8), $\mathcal{L}_{s,t}$ where $(s, t) \in \mathbb{R}^2$. Note that $\mathcal{L}_{s,t}^{-1} \cong \mathcal{L}_{-s,t}$. Since $A^*(x, y, z) = A(\bar{x}, \bar{y}, \bar{z})$ and always holds $\mathcal{E} \cong \mathcal{E}^\vee \otimes \det \mathcal{E}$, we conclude that $\mathcal{E}^\vee = \text{Ker } A$ is a nontrivial extension of $\mathcal{L}_{-s,t}$.

Analogous computations as in the proof of Theorem 3.2 show that A is unitarily equivalent to a self-adjoint representation

$$x \begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} + z \begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} + y \begin{bmatrix} 0 & * & * & \frac{3t^2-2t(1+\lambda)-(1-\lambda)^2}{4} & i s & \frac{t-1-\lambda}{2} \\ 0 & * & & -i s & -t & 0 \\ 0 & & & \frac{t-1-\lambda}{2} & 0 & -1 \\ & & & 0 & 0 & 0 \\ & & & & 0 & 0 \\ & & & & & 0 \end{bmatrix} \text{ for } (s, t) \in \mathbb{R}^2,$$

or i times the skew-symmetric representation (5)

$$x \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & i & 0 & 0 \\ 0 & i & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} + z \begin{bmatrix} 0 & 0 & 0 & 0 & i & 0 \\ 0 & 0 & i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} + y \begin{bmatrix} 0 & i & 0 & \frac{3t^2-2t(1+\lambda)-(1-\lambda)^2}{-4i} & 0 & \frac{t-1-\lambda}{-2i} \\ 0 & 0 & & 0 & -it & 0 \\ 0 & & & \frac{t-1-\lambda}{-2i} & 0 & -i \\ & & & 0 & 0 & 0 \\ & & & & 0 & 0 \\ & & & & & 0 \end{bmatrix} \text{ for } t = 0, 1, \lambda.$$

It is easy to check that none of these are definite and can thus not provide a counterexample to Kippenhahn’s conjecture. \square

REFERENCES

[1] M. Atiyah. *Vector bundles over an elliptic curve*, Proc. London Math. Soc. (3), **7** (1957), 414–452.
 [2] J. Backelin, J. Herzog, and H. Sanders. *Matrix factorizations of homogeneous polynomials*, Algebra Some Current Trends **1352**, Lecture Notes in Mathematics, Springer (2006),1–33.
 [3] A. Beauville. *Determinantal Hypersurfaces*, Michigan Math. J. **48** (2000), 39–63.
 [4] A. Beauville. *Vector bundles on curves and generalized theta functions: recent results and open problems*, Complex Algebraic Geometry, MSRI Publications **28** (1995), 17–33.
 [5] A Buckley and T. Košir. *Plane Curves as Pfaffians*, Annali Della Scuola Normale Superiore di Pisa-Classe di Scienze (2010).
 [6] R. J. Cook and A. D. Thomas. *Line bundles and homogeneous matrices*, Quart. J. Math. Oxford Ser. (2) **30** (1979), 423–429.
 [7] L.E. Dickson. *Determination of All General Homogeneous Polynomials Expressible as Determinants with Linear Elements*, Trans. Amer. Math. Soc. **22** (1921), 167–179.

- [8] D. Eisenbud. *Homological algebra on a complete intersection, with an application to group representations*, Trans. Amer. Math. Soc. **260** (1980), 35–64.
- [9] D. Eisenbud. *The Geometry of Syzygies: A Second Course in Algebraic Geometry and Commutative Algebra*, Graduate Texts in Mathematics, Springer (2005).
- [10] H. Grassman. *Die stereometrischen Gleichungen zweiten Grades, und die dadurch erzeugten Oberflächen*, J. Reine Angew. Math. **49** (1855), 47–65.
- [11] R. Kippenhahn. *Über den Wertevorrat einer Matrix*, Mathematische Nachrichten Volume 6, **3-4** (1951), 193–228.
- [12] T. J. Laffey. *A counterexample to kippenhahn’s conjecture on hermitian pencils*, Lin. Alg. Appl. **51** (1983), 179–182.
- [13] P. Lancaster and L. Rodman. *Canonical forms for symmetric / skew-symmetric real matrix pairs under strict equivalence and congruence*, Linear Algebra Appl. **406**, (2005), 1–76.
- [14] Chi-Kwong Li and Ilya Spitkovsky. *Equality of Higher Numerical Ranges of Matrices and a Conjecture of Kippenhahn on Hermitian Pencils*, Lin. Alg. Appl. **58** (1998), 323–349.
- [15] D. Mumford. *Abelian varieties*, Tata institute of fundamental research, Bombay, 1970.
- [16] T. Netzer and A. Thom. *Polynomials with and without determinantal representations*, Linear Algebra Appl. **437** (2012), 1579–1595.
- [17] G. V. Ravindra and Amit Tripathi. *Torsion points and matrices defining elliptic curves*, International Journal of Algebra and Computation Vol 24, **6**, 2014.
- [18] R. Quarez. *Symmetric determinantal representation of polynomials*, Linear Algebra Appl. **436** (2012), 3642–3660.
- [19] H. Shapiro. *A conjecture of Kippenhahn about the characteristic polynomial of a pencil generated by two Hermitian matrices. I and II*, Lin. Alg. Appl. **43** (1982), 201–221 and **45** (1982), 97–108.
- [20] H. Shapiro. *Hermitian pencils with a cubic minimal polynomial*, Lin. Alg. Appl. **48** (1982), 81–103.
- [21] F. Schur. *Ueber die durch collineare Grundgebilde erzeugten Curven und Flächen*, Mathematische Annalen, Vol. 18 **1** (1881), 1–32.
- [22] V. Vinnikov. *Complete description of determinantal representations of smooth irreducible curves*, Lin. Alg. Appl., **125** (1989), 103–140.
- [23] V. Vinnikov. *Self-adjoint determinantal representations of real irreducible cubics*, Operator Theory: Advances and Applications, **19** (1986), 415–442.
- [24] V. Vinnikov. *LMI representations of convex semialgebraic sets and determinantal representations of algebraic hypersurfaces: Past, present, and future*, Operator Theory: Advances and Applications, **222** (2012), 325–349.
- [25] W. C. Waterhouse. *The codimension of singular matrix pairs*, Lin. Alg. Appl. **47** (1984), 227–245.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LJUBLJANA, JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA. e-mail: anita.buckley@fmf.uni-lj.si