

Oka domains in Euclidean and projective spaces

Franc Forstnerič

Univerza v Ljubljani



Complex Analysis and Geometry XXVI
Levico Terme, 20 September 2022

Oka manifolds

A complex manifold Y is called an **Oka manifold**¹ if maps $X \rightarrow Y$ from any Stein manifold (or a reduced Stein space) X satisfy the following conditions:

- every continuous map $f : X \rightarrow Y$ can be homotopically deformed to a holomorphic map $\tilde{f} : X \rightarrow Y$.
- If the map $f : X \rightarrow Y$ is holomorphic on a compact $\mathcal{O}(X)$ -convex set K and on a closed complex subvariety X' of X , then there is a homotopy from f to \tilde{f} consisting of maps which are holomorphic near K and close to f on K , and which agree with f on X' .
- The analogous approximation and (jet) interpolation properties hold for families of maps $X \rightarrow Y$ depending continuously on a parameter.

Observation: The Kobayashi–Royden pseudometric vanishes identically on an Oka manifold.

¹F. Forstnerič, Oka manifolds, C. R. Acad. Sci. Paris **347:17-18** (2009)

Oka manifolds are like bitcoins — precious but hard to discover

- **Oka 1939, Grauert 1958** Complex Lie groups and their homogeneous manifolds are Oka.
- **Gromov 1989** Every elliptic complex manifold is an Oka manifold.

A complex manifold Y is said to be elliptic if it admits a holomorphic vector bundle $\pi : E \rightarrow Y$ and a **dominating holomorphic spray** $F : E \rightarrow Y$ such that for every point $y \in Y$ we have that $F(0_y) = y$ and

$dF_{0_y} : T_{0_y}E \rightarrow T_yY$ maps the fibre E_y onto the tangent space T_yY .

- In particular, if Y admits complete holomorphic vector fields which span the tangent space at every point, then Y is elliptic and hence Oka.
- **F. 2006** A complex manifold Y is Oka iff it satisfies the following **Convex approximation property**: Every holomorphic map $K \rightarrow Y$ from a (neighbourhood of a) compact convex set K in a Euclidean space \mathbb{C}^n is a limit of entire maps $\mathbb{C}^n \rightarrow Y$.

Kusakabe's characterization of Oka manifolds

Y. Kusakabe 2021 A complex manifold Y is an Oka manifold if and only if every holomorphic map $f : K \rightarrow Y$ from (a neighbourhood of) a compact convex set $K \subset \mathbb{C}^N$ is the core map of a dominating holomorphic spray $F : K \times \mathbb{C}^n \rightarrow Y$ for some $n \geq \dim Y$:

$F(\cdot, 0) = f$ and $\left. \frac{\partial}{\partial z} \right|_{z=0} F(\zeta, z) : \mathbb{C}^n \rightarrow T_{f(\zeta)} Y$ is surjective for every $\zeta \in K$.

This is a restricted version of condition Ell_1 studied by Gromov in 1989. Kusakabe gave a short but ingenious proof that

$$\text{Ell}_1 \implies \text{CAP};$$

the rest was known before. However, Ell_1 is often easier to verify than CAP. As an application, Kusakabe proved

The localization theorem for Oka manifolds:

If a complex manifold Y is a union of Zariski-open Oka domains $Y \setminus A_i$, with A_i a closed complex subvariety of Y , then Y is Oka.

Which domains in \mathbb{C}^n and $\mathbb{C}\mathbb{P}^n$ are Oka?

- Until Kusakabe's work the only known examples of Oka domains in \mathbb{C}^n were FB-domains, complements of complex hyperplanes, and complements of closed tame (in particular, algebraic) subvarieties of dimension $\leq n - 2$.
- **Kusakabe 2021** For every compact polynomially convex set $K \subset \mathbb{C}^n$ for $n > 1$, the complement $\mathbb{C}^n \setminus K$ is Oka.
- **F. 2022** If $K \subset \mathbb{C}^n$ is as above then $\mathbb{C}\mathbb{P}^n \setminus K$ is Oka.

Furthermore, if Γ is a compact union of curves such that the complex curve $\widehat{K \cup \Gamma} \setminus K \cup \Gamma$ has at most finitely many irreducible components, then $\mathbb{C}^n \setminus K \cup \Gamma$ and $\mathbb{C}\mathbb{P}^n \setminus K \cup \Gamma$ are Oka.

- In a recent joint work with E.F. Wold, we found surprisingly small Oka domains in \mathbb{C}^n ($n > 1$) at the limit of what is possible.
F. F. and E. F. Wold 2022 Oka domains in Euclidean spaces.
<https://arxiv.org/abs/2203.12883>

Oka complements of closed convex sets

Theorem (1)

If E is a closed convex set with \mathcal{C}^1 boundary in \mathbb{C}^n for $n > 1$ such that $E \cap T_p^{\mathbb{C}} bE$ does not contain an affine real halfline for any $p \in bE$, then $\mathbb{C}^n \setminus E$ is an Oka domain. This holds in particular if E is strictly convex, i.e., bE does not contain any straight line segments.

There are many such examples of the form

$$E = \{(z', z_n) \in \mathbb{C}^n : \Im z_n \geq \phi(z', \Re z_n)\},$$

where ϕ is a convex function of class \mathcal{C}^1 . If $t > 0$, the convex domain

$$\Omega_t^+ = \{\Im z_n > t\phi(z', \Re z_n)\}$$

does not contain any affine complex line, so it is hyperbolic (Barth 1980, Harris 1979, Bracci and Saracco 2009), while the domain

$$\Omega_t^- = \{\Im z_n < t\phi(z', \Re z_n)\}$$

is Oka. For $t < 0$ the picture is reversed, while at $t = 0$ the hyperplane $\{\Im z_n = 0\}$ splits \mathbb{C}^n in a pair of halfspaces which are neither Oka nor hyperbolic.

Oka domains below convex graphs

A convex function $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$ is **irreducible** if it is not of the form $\phi = \psi \circ P + l$ where $P : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear projection with $m < n$, ψ is a convex function on \mathbb{R}^m , and l is a linear function on \mathbb{R}^n .

Corollary

If ϕ is an irreducible convex function on $\mathbb{C}^{n-1} \times \mathbb{R}$, then the domain

$$\Omega_\phi = \{(z', z_n) \in \mathbb{C}^n : \Im z_n < \phi(z', \Re z_n)\}$$

is Oka. The same holds for domains $\Omega_\phi = \{(z', z_n) \in \mathbb{C}^n : \Im z_n < \phi(z')\}$.

Proof By Azagra (2013) the condition on ϕ implies that for every $\epsilon > 0$ there is a smooth strictly convex function $\psi : \mathbb{C}^{n-1} \times \mathbb{R} \rightarrow \mathbb{R}$ such that $\phi - \epsilon < \psi < \phi$. Hence, the domain $\Omega_\psi = \{\Im z_n < \psi(z', \Re z_n)\}$ is Oka.

This gives an increasing sequence $\phi_1 < \phi_2 < \phi_3 < \dots$ of smooth strictly convex functions on $\mathbb{C}^{n-1} \times \mathbb{R}$ converging uniformly to ϕ such that the sequence of Oka domains Ω_{ϕ_j} increases to Ω_ϕ as $j \rightarrow \infty$. Hence, Ω_ϕ is Oka.

A similar argument applies in the second case.

Complements of convex domains containing no lines

Example

Every concave wedge in \mathbb{C}^n for $n > 1$ of the form

$$\Im z_n < c|\Re z_n| + \sum_{j=1}^{n-1} (a_j|\Re z_j| + b_j|\Im z_j|)$$

for $c \geq 0$ and positive numbers $a_1, \dots, a_{n-1}, b_1, \dots, b_{n-1}$ is Oka.

Corollary

If E is a closed convex set in \mathbb{C}^n for $n > 1$ which does not contain any affine real line, then $\mathbb{C}^n \setminus E$ is an Oka domain.

Proof There is a decreasing sequence $E_1 \supset E_2 \supset E_3 \supset \dots$ of smoothly bounded strongly convex sets in \mathbb{C}^n such that $E = \bigcap_{j=1}^{\infty} E_j$.

By Theorem 1, the domain $\Omega_j = \mathbb{C}^n \setminus E_j$ is Oka for every $j \in \mathbb{N}$.

Hence, $\mathbb{C}^n \setminus E = \bigcup_{j=1}^{\infty} \Omega_j$ is an increasing union of Oka domains, so it is Oka.

The main theorem

Theorem 1 follows from our main result, Theorem 2. Given a closed set $E \subset \mathbb{C}^n$, we denote by $\bar{E} \subset \mathbb{C}\mathbb{P}^n$ its projective closure.

Theorem (2)

If E is a closed subset of \mathbb{C}^n for $n > 1$ and $\Lambda \subset \mathbb{C}\mathbb{P}^n$ is a hyperplane such that $\bar{E} \cap \Lambda = \emptyset$ and \bar{E} is polynomially convex in $\mathbb{C}\mathbb{P}^n \setminus \Lambda \cong \mathbb{C}^n$, then $\mathbb{C}^n \setminus E$ and $\mathbb{C}\mathbb{P}^n \setminus \bar{E}$ are Oka domains.

*This holds in particular if $\mathbb{C}\mathbb{P}^n \setminus \bar{E}$ is a union of a connected family of projective hyperplanes. (Such a set is called **projectively convex**.)*

The second part follows from the first one by observing that if K is a compact set in $\mathbb{C}\mathbb{P}^n$ and $\Lambda_t \subset \mathbb{C}\mathbb{P}^n$ ($t \in [0, 1]$) is a path of hyperplanes not intersecting K , then $\bigcup_{t \in (0, 1]} \Lambda_t$ does not intersect the polynomial hull of K in $\mathbb{C}\mathbb{P}^n \setminus \Lambda_0$.

Fact: Every convex set $E \subset \mathbb{C}^n$ in Theorem 1 has projectively convex closure. In fact, $\mathbb{C}\mathbb{P}^n \setminus \bar{E}$ is a union of hyperplanes parallel to the complex tangent spaces $T_p^{\mathbb{C}} bE$ for $p \in bE$. **Hence, Theorem 2 implies Theorem 1.**

Proof of Theorem 2

Let $H = \mathbb{C}\mathbb{P}^n \setminus \mathbb{C}^n$ denote the hyperplane at infinity. Set $K = \overline{E}$. Choose a projective hyperplane $\Lambda \subset \mathbb{C}\mathbb{P}^n$ with $K \cap \Lambda = \emptyset$.

Let $z = (z_1, \dots, z_n)$ be affine coordinates on $\mathbb{C}\mathbb{P}^n \setminus \Lambda \cong \mathbb{C}^n$ in which $H \setminus \Lambda = \{z_n = 0\}$. By the hypothesis, K is polynomially convex in these coordinates. It now suffices to prove the following result.

Theorem (3)

If K is a compact polynomially convex set in \mathbb{C}^n then $(\mathbb{C}^{n-1} \times \mathbb{C}^) \setminus K$ is Oka.*

Assume that Theorem 3 holds. Then,

$$\mathbb{C}^n \setminus (E \cup \Lambda) = \mathbb{C}\mathbb{P}^n \setminus (H \cup E \cup \Lambda) = (\mathbb{C}\mathbb{P}^n \setminus \Lambda) \setminus (H \cup E)$$

is Oka. Choose hyperplanes $\Lambda_0 = \Lambda, \Lambda_1, \dots, \Lambda_n$ in $\mathbb{C}\mathbb{P}^n \setminus K$ close to Λ such that $\bigcap_{i=0}^n \Lambda_i = \emptyset$. Then, $\mathbb{C}^n \setminus (E \cup \Lambda_i)$ is Oka for every i and

$$\Omega = \mathbb{C}^n \setminus E = \bigcup_{i=0}^n (\mathbb{C}^n \setminus E) \setminus \Lambda_i$$

is a union of Zariski open Oka domains, so it is Oka.

Proof of Theorem 3

In view of Kusakabe's characterisation of Oka manifolds by Condition Ell_1 , it suffices to prove the following.

Theorem (4)

Assume that

- K is a compact polynomially convex set in \mathbb{C}^n for some $n > 1$,
- L is a compact (polynomially) convex set in \mathbb{C}^N for some $N \in \mathbb{N}$, and
- $f : L \rightarrow \mathbb{C}^n$ is a holomorphic map such that

$$f(\zeta) \in (\mathbb{C}^{n-1} \times \mathbb{C}^*) \setminus K \text{ holds for all } \zeta \in L.$$

Then there is a holomorphic map $F : L \times \mathbb{C}^n \rightarrow \mathbb{C}^n$ such that for every $\zeta \in L$,

$$F(\zeta, 0) = f(\zeta) \text{ and the map } F(\zeta, \cdot) : \mathbb{C}^n \rightarrow (\mathbb{C}^{n-1} \times \mathbb{C}^*) \setminus K \text{ is injective.}$$

It follows that

$$\Omega_\zeta = \{F(\zeta, z) : z \in \mathbb{C}^n\} \subset (\mathbb{C}^{n-1} \times \mathbb{C}^*) \setminus K$$

is a family of Fatou–Bieberbach domains depending holomorphically on $\zeta \in L$.

Proof of Theorem 4

Since the set $L \subset \mathbb{C}^N$ is polynomially convex, we may assume that the map f is defined on a Stein neighbourhood U of L which is a Runge in \mathbb{C}^N .

The graph

$$\Gamma = \{(\zeta, f(\zeta)) : \zeta \in U\}$$

is a closed Stein submanifold of the Stein domain $X = U \times \mathbb{C}^n$ which is a Runge in \mathbb{C}^{N+n} . The restricted graph

$$\Gamma_L = \{(\zeta, f(\zeta)) \in X : \zeta \in L\} \subset L \times ((\mathbb{C}^{n-1} \times \mathbb{C}^*) \setminus K)$$

is then polynomially convex in $\mathbb{C}^N \times \mathbb{C}^n$.

It follows that $(L \times K) \cup \Gamma_L$ is $\mathcal{O}(X)$ -convex and hence polynomially convex.

Proof of Theorem 4, part 2

Let $\pi : \mathbb{C}^N \times \mathbb{C}^n \rightarrow \mathbb{C}^N$ denote the projection. Consider the injective π -fibre preserving holomorphic map $\Phi = (\text{Id}, \phi)$ of the form

$$\Phi(\zeta, z) = (\zeta, \phi(\zeta, z)) \quad \text{for } (\zeta, z) \in \Omega$$

on a small Runge Stein neighbourhood $\Omega = \Omega' \cup \Omega''$ of $(L \times K) \cup \Gamma_L$ in $\mathbb{C}^N \times \mathbb{C}^n$ which equals the identity map on a neighbourhood Ω' of $L \times K$ and

$$\phi(\zeta, z) = f(\zeta) + \frac{1}{2}(z - f(\zeta)) = \frac{1}{2}f(\zeta) + \frac{1}{2}z$$

for (ζ, z) in a neighbourhood Ω'' of Γ_L . Thus, $\phi(\zeta, \cdot)$ is a contraction by the factor $1/2$ around the point $f(\zeta) \in \mathbb{C}^n$ for every $\zeta \in L$.

On a suitable neighbourhood Ω'' of Γ_L the map $\phi = \phi_{1/2}$ is connected to $\phi_0(\zeta, z) = z$ by the isotopy $\phi_t : \Omega'' \rightarrow \Omega''$ ($t \in [0, 1/2]$) given by

$$\phi_t(\zeta, z) = tf(\zeta) + (1-t)z.$$

On $\Omega' \supset L \times K$ we take the constant isotopy $\phi_t(\zeta, z) = \phi_0(\zeta, z) = z$ for all t . Clearly, the trace of the isotopy $\Phi_t = (\text{Id}, \phi_t)$ for $t \in [0, 1/2]$ consists of Runge domains $\Phi_t(\Omega) \subset \Omega$.

Proof of Theorem 4, part 3

Varolin 2001 The Lie algebra of holomorphic (algebraic) vector fields on \mathbb{C}^n vanishing on $\mathbb{C}^{n-1} \times \{0\}$ has the (algebraic) density property.

Hence, we can approximate Φ on $(L \times K) \cup \Gamma_L$ by a holomorphic map

$$\Psi : V \times \mathbb{C}^n \rightarrow V \times \mathbb{C}^n, \quad \Psi(\zeta, z) = (\zeta, \psi(\zeta, z)),$$

where $V \subset U$ is a neighbourhood of L , such that for every $\zeta \in V$ we have that

- $\psi(\zeta, \cdot) \in \text{Aut}(\mathbb{C}^n)$,
- $\psi(\zeta, z) = z$ for every $z = (z', 0) \in \mathbb{C}^{n-1} \times \{0\}$, and
- $\psi(\zeta, f(\zeta)) = f(\zeta)$.

Choose $a, b \in \mathbb{R}$ such that

$$0 < a < 1/2 < b < 1 \quad \text{and} \quad b^2 < a.$$

If the approximation of ϕ by ψ is close enough then the estimate

$$a|z - f(\zeta)| \leq |\psi(\zeta, z) - f(\zeta)| \leq b|z - f(\zeta)| \tag{1}$$

holds in a neighbourhood of the graph Γ_L of f , and ψ is arbitrarily close to the map $(\zeta, z) \mapsto z$ on a neighbourhood of $L \times K$.

Proof of Theorem 4, part 4

This gives a sequence of holomorphic maps ψ_k of the same kind as ψ such that the estimate (1) holds for all of them on the same neighbourhood of Γ_L , and ψ_k converges to the map $(\zeta, z) \mapsto z$ on a neighbourhood of $L \times K$ as $k \rightarrow \infty$.

Consider the sequence of random iterations

$$\theta_k(\zeta, \cdot) = \psi_k(\zeta, \cdot) \circ \psi_{k-1}(\zeta, \cdot) \circ \cdots \circ \psi_1(\zeta, \cdot) \in \text{Aut}(\mathbb{C}^n).$$

Due to the condition $b^2 < a$ in the estimate (1) the attracting basin $B_\zeta \subset \mathbb{C}^n$ of the sequence θ_k at the fixed point $f(\zeta)$ is biholomorphic to \mathbb{C}^n (Wold 2005).

If the convergence of the sequence ψ_k to the map $(\zeta, z) \mapsto z$ is fast enough on a neighbourhood of $L \times K$, then none of the basins B_ζ intersect K .

Furthermore, the condition $\psi_k(\zeta, (z', 0)) = (z', 0)$ for all $\zeta \in L$, $z' \in \mathbb{C}^{n-1}$, and $k \in \mathbb{N}$ ensures that the basin B_ζ does not intersect $\mathbb{C}^{n-1} \times \{0\}$.

This gives a holomorphic map $F : V \times \mathbb{C}^n \rightarrow \mathbb{C}^n$ such that the image B_ζ of $F(\zeta, \cdot)$ is a Fatou–Bieberbach domain in $(\mathbb{C}^{n-1} \times \mathbb{C}^*) \setminus K$ centred at $f(\zeta)$ for every $\zeta \in V$. Hence, Theorem 4 is proved.

Compact sets in $\mathbb{C}\mathbb{P}^n$ avoided by more general hypersurfaces

Theorem

Let K be a compact set in $\mathbb{C}\mathbb{P}^n$ ($n > 1$) and $\Lambda \subset \mathbb{C}\mathbb{P}^n$ be a closed complex hypersurface with $K \cap \Lambda = \emptyset$ such that K is holomorphically convex in the Stein domain $\Omega = \mathbb{C}\mathbb{P}^n \setminus \Lambda$. If Ω has the density property then $\mathbb{C}\mathbb{P}^n \setminus K$ is an Oka domain.

This follows by a similar argument as above:

- the complement of K in $\mathbb{C}\mathbb{P}^n \setminus \Lambda$ is Oka by Kusakabe's theorem,
- by moving Λ with automorphisms of $\mathbb{C}\mathbb{P}^n$ we find finitely many hypersurfaces $\Lambda_0 = \Lambda, \Lambda_1, \dots, \Lambda_m$ not intersecting K such that $\bigcap_{i=0}^m \Lambda_i = \emptyset$ and $(\mathbb{C}\mathbb{P}^n \setminus \Lambda_i) \setminus K$ is Oka for all i , and hence

$$\mathbb{C}\mathbb{P}^n \setminus K = \bigcup_{i=0}^m (\mathbb{C}\mathbb{P}^n \setminus \Lambda_i) \setminus K$$

is Oka by the localization theorem.

A corollary

Denote by

$$\mathcal{V}_k(\mathbf{CP}^n) \cong \mathbf{CP}^N \quad \text{with} \quad N = \binom{n+k}{k} - 1$$

the space of degree k complex hypersurfaces in \mathbf{CP}^n . For any $\Lambda \in \mathcal{V}_k(\mathbf{CP}^n)$ the complement $\mathbf{CP}^n \setminus \Lambda$ is a closed affine manifold, hence a Stein manifold.

Corollary

Let B be an open connected set in $\mathcal{V}_k(\mathbf{CP}^n)$ ($k \geq 1$, $n \geq 2$). If for some $\Lambda_0 \in B$ the domain $\mathbf{CP}^n \setminus \Lambda_0$ has the density property, then

$$\Omega(B) = \bigcup_{\Lambda \in B} \Lambda \subset \mathbf{CP}^n$$

is Oka. In particular, Λ_0 has a basis of open Oka neighbourhoods in \mathbf{CP}^n .

Problem

Which complex hypersurfaces in \mathbf{CP}^n have complements having the density property?

Examples and a conjecture

Example

- If $\Lambda_1, \dots, \Lambda_k \subset \mathbb{C}P^n$ ($n > 1$, $1 \leq k \leq n$) are hyperplanes in general position then $\bigcup_{i=1}^k \Lambda_k$ has a basis of Oka neighbourhoods in $\mathbb{C}P^n$.

Indeed, $\mathbb{C}P^n \setminus \bigcup_{i=1}^k \Lambda_k$ is isomorphic to $\mathbb{C}^{n-k+1} \times (\mathbb{C}^*)^{k-1}$ with $n - k + 1 \geq 1$. This domain has the density property (Varolin 2001), so the result follows from the previous corollary.

The proof fails for more than n hyperplanes. In particular, it is not known whether $(\mathbb{C}^*)^n$ for $n > 1$ has the density property.

- If Λ is a quadric hypersurface in $\mathbb{C}P^n$ ($n > 1$) then $\mathbb{C}P^n \setminus \Lambda$ has the density property.

Conjecture

For a generic hypersurface $\Lambda \subset \mathbb{C}P^n$ ($n > 1$) of degree at most n the complement $\mathbb{C}P^n \setminus \Lambda$ has the density property.

THANK YOU

FOR YOUR ATTENTION