

# Complex analytic methods in minimal surface theory

Franc Forstnerič

University of Ljubljana

<http://www.fmf.uni-lj.si/~forstneric/>

American University in Beirut

24–27 November 2014

# Lecture 3: Riemann-Hilbert problem for null holomorphic curves, the Calabi-Yau problem, and minimal hulls

The aim of this lecture is to provide **approximate solutions** to certain **Riemann-Hilbert boundary value problems** for **holomorphic null curves** in the complex Euclidean space  $\mathbb{C}^3$ , and derive geometric applications.

## References:

- Antonio Alarcón and Franc Forstnerič, *Null curves and directed immersions of open Riemann surfaces*. Invent. Math. **196** (2014), no. 3, 733–771
- Antonio Alarcón and Franc Forstnerič, *The Calabi-Yau problem, null curves, and Bryant surfaces*. <http://arxiv.org/abs/1308.0903>
- Barbara Drinovec Drnovšek and Franc Forstnerič, *Minimal hulls of compact sets in  $\mathbb{R}^3$* . <http://arxiv.org/abs/1409.6906>

# The classical Riemann-Hilbert problem

- Let  $f: \overline{\mathbb{D}} \rightarrow \mathbb{C}^n$  be a  $\mathcal{C}^0$  map that is holomorphic in  $\mathbb{D}$ . Let

$$g: b\mathbb{D} \times \overline{\mathbb{D}} \rightarrow \mathbb{C}^n$$

be a  $\mathcal{C}^0$  map such that  $g(\zeta, \cdot)$  is holomorphic in  $\mathbb{D}$  and  $g(\zeta, 0) = f(\zeta)$  for every  $\zeta \in b\mathbb{D}$ .

Let  $K \subset \mathbb{D}$  be a compact set and let  $\epsilon > 0$ .

- Problem:** Find a  $\mathcal{C}^0$  map  $\phi: \overline{\mathbb{D}} \rightarrow \mathbb{C}^n$ , holomorphic in  $\mathbb{D}$ , such that
  - $\phi$  is  $\epsilon$ -close to  $f$  over  $K$ , and
  - $\phi(\zeta)$  is  $\epsilon$ -close to the curve  $g(\zeta, b\overline{\mathbb{D}})$  for every  $\zeta \in b\mathbb{D}$ .
- The model case:**  $n = 2$ ,  $f(\zeta) = (\zeta, 0)$  and  $g(\zeta, \zeta) = (\zeta, \zeta)$ ; take  $\phi(\zeta) = (\zeta, \zeta^N)$  for large  $N \in \mathbb{N}$ .
- Solutions exist even when  $\overline{\mathbb{D}}$  is replaced by a **bordered Riemann surface** and  $\mathbb{C}^n$  by any complex manifold (**Forstnerič** and **Drinovec Drnovšek**, Duke Math. J. 139 (2007)). This is useful in a variety of problems.

# Null (holomorphic) curves in $\mathbb{C}^3$

Let  $M$  be either an open or a bordered Riemann surface.

- Recall that a holomorphic immersion

$$F = (F_1, F_2, F_3): M \rightarrow \mathbb{C}^3$$

is a **null curve** if it is directed by the conical quadric subvariety

$$\mathfrak{A} = \{z = (z_1, z_2, z_3) \in \mathbb{C}^3: z_1^2 + z_2^2 + z_3^2 = 0\},$$

in the sense that the derivative  $F' = (F'_1, F'_2, F'_3)$  with respect to any local holomorphic coordinate on  $M$  takes values in  $\mathfrak{A}_* = \mathfrak{A} \setminus \{0\}$ .

- The real and the imaginary part of a null curve  $F: M \rightarrow \mathbb{C}^3$  are conformal minimal immersions in  $\mathbb{R}^3$ .
- Every conformal minimal immersion  $\mathbb{D} \rightarrow \mathbb{R}^3$  is the real part of a null curve  $\mathbb{D} \rightarrow \mathbb{C}^3$ . (This fails on non-simply connected Riemann surfaces due to the possibly nonzero periods of the harmonic conjugate.)

# Geometry of the null quadric

$$\mathfrak{A} = \{z = (z_1, z_2, z_3) \in \mathbb{C}^3 : z_1^2 + z_2^2 + z_3^2 = 0\}.$$

- $\mathfrak{A}$  is a complex cone with vertex at 0;  $\mathfrak{A}_* = \mathfrak{A} \setminus \{0\}$  is smooth.
- $\mathfrak{A}_*$  is a holomorphic fiber bundle with fiber  $\mathbb{C}_*$  over the curve  $\Lambda = \{[z_1 : z_2 : z_3] : z_1^2 + z_2^2 + z_3^2 = 0\} \subset \mathbb{C}P^2$ . We have  $\Lambda \cong \mathbb{C}P^1$ .
- $\mathfrak{A}_*$  is an **Oka manifold**; in particular, maps  $M \rightarrow \mathfrak{A}_*$  satisfy the Runge and Mergelyan approximation properties.
- The **spinor representation**:

$$\pi: \mathbb{C}^2 \rightarrow \mathfrak{A}, \quad \pi(u, v) = (u^2 - v^2, i(u^2 + v^2), 2uv).$$

The map  $\pi: \mathbb{C}_*^2 \rightarrow \mathfrak{A}_*$  is a nonramified two-sheeted covering.

# Existence and general position theorem for null curves

Let  $\theta$  be a nowhere vanishing holomorphic 1-form on  $M$ .

There is a bijective correspondence (up to translations)

$$\{F: M \rightarrow \mathbb{C}^3 \text{ null curve}\} \longleftrightarrow \{f: M \rightarrow \mathfrak{A}_* \text{ holomorphic, } f\theta \text{ exact}\}$$

$$F(p) = F(p_0) + \int_{p_0}^p f\theta, \quad dF = f\theta.$$

## Theorem (Lecture 1, Theorem 3)

The *general position* of null curves in  $\mathbb{C}^3$  is *embedded*.

# Riemann-Hilbert problem for null discs

## Theorem

Let  $F: \overline{\mathbb{D}} \rightarrow \mathbb{C}^3$  be a null curve, let  $\vartheta \in \mathfrak{A}_*$  be a null vector, let

$$\mu: b\mathbb{D} = \{\zeta \in \mathbb{C}: |\zeta| = 1\} \rightarrow \mathbb{R}_+ := [0, +\infty)$$

be a continuous function (the *size function*), and set

$$\varkappa: b\mathbb{D} \times \overline{\mathbb{D}} \rightarrow \mathbb{C}^3, \quad \varkappa(\zeta, \xi) = F(\zeta) + \mu(\zeta) \xi \vartheta.$$

Given numbers  $\epsilon > 0$  and  $0 < r < 1$ , there exist a number  $r' \in [r, 1)$  and a null holomorphic immersion  $G: \overline{\mathbb{D}} \rightarrow \mathbb{C}^3$  satisfying

- $G$  is  $\epsilon$ -close to  $F$  in the  $\mathcal{C}^1$  topology on  $\{\zeta \in \mathbb{C}: |\zeta| \leq r'\}$ ,
- $\text{dist}(G(\zeta), \varkappa(\zeta, b\mathbb{D})) < \epsilon$  for all  $\zeta \in b\mathbb{D}$ , and
- $\text{dist}(G(\rho\zeta), \varkappa(\zeta, \overline{\mathbb{D}})) < \epsilon$  for all  $\zeta \in b\mathbb{D}$  and all  $\rho \in [r', 1)$ .

Furthermore, if  $I$  is a compact arc in  $b\mathbb{D}$ , the size function  $\mu$  vanishes everywhere on  $b\mathbb{D} \setminus I$ , and  $U$  is an open neighborhood of  $I$  in  $\overline{\mathbb{D}}$ , then

- one can choose  $G$  to be  $\epsilon$ -close to  $F$  in the  $\mathcal{C}^1$  topology on  $\overline{\mathbb{D}} \setminus U$ .

$$F' = \pi(u, v) = (u^2 - v^2, i(u^2 + v^2), 2uv) : \overline{\mathbb{D}} \rightarrow \mathfrak{A}_*$$

$$\vartheta = \pi(p, q) = (p^2 - q^2, i(p^2 + q^2), 2pq) \in \mathfrak{A}_*$$

$$\eta = \sqrt{\mu} : b\mathbb{D} \rightarrow \mathbb{R}_+$$

$$\eta(\zeta) \approx \tilde{\eta}(\zeta) = \sum_{j=1}^N A_j \zeta^{j-m} \quad (\text{rational approximation on } \mathbb{C}_*)$$

$$u_n(\xi) = u(\xi) + \sqrt{2n+1} \tilde{\eta}(\xi) \xi^n p \quad (n > m, u_n(0) = u(0))$$

$$v_n(\xi) = v(\xi) + \sqrt{2n+1} \tilde{\eta}(\xi) \xi^n q \quad (v_n(0) = v(0))$$

$$\Phi_n = \pi(u_n, v_n) = (u_n^2 - v_n^2, i(u_n^2 + v_n^2), 2u_n v_n) : \overline{\mathbb{D}} \rightarrow \mathfrak{A}_*$$

$$G_n(\zeta) = F(0) + \int_0^\zeta \Phi_n(\xi) d\xi, \quad \zeta \in \overline{\mathbb{D}}.$$

# Proof

$$F' = \pi(u, v) = (u^2 - v^2, i(u^2 + v^2), 2uv) : \overline{\mathbb{D}} \rightarrow \mathfrak{A}_*$$

$$\vartheta = \pi(p, q) = (p^2 - q^2, i(p^2 + q^2), 2pq) \in \mathfrak{A}_*$$

$$\eta = \sqrt{\mu} : b\mathbb{D} \rightarrow \mathbb{R}_+$$

$$\eta(\zeta) \approx \tilde{\eta}(\zeta) = \sum_{j=1}^N A_j \zeta^{j-m} \quad (\text{rational approximation on } \mathbb{C}_*)$$

$$u_n(\zeta) = u(\zeta) + \sqrt{2n+1} \tilde{\eta}(\zeta) \zeta^n p \quad (n > m, u_n(0) = u(0))$$

$$v_n(\zeta) = v(\zeta) + \sqrt{2n+1} \tilde{\eta}(\zeta) \zeta^n q \quad (v_n(0) = v(0))$$

$$\Phi_n = \pi(u_n, v_n) = (u_n^2 - v_n^2, i(u_n^2 + v_n^2), 2u_n v_n) : \overline{\mathbb{D}} \rightarrow \mathfrak{A}_*$$


$$G_n(\zeta) = F(0) + \int_0^\zeta \Phi_n(\xi) d\xi, \quad \zeta \in \overline{\mathbb{D}}.$$

It follows that  $G_n(\zeta) \approx F(\zeta) + \zeta^{2n+1} \mu(\zeta) \vartheta$ . Take  $G = G_n$  for large enough  $n \in \mathbb{N}$ .

# Proof

Furthermore, if  $I$  is a compact arc in  $b\mathbb{D}$ , the size function  $\mu$  vanishes everywhere on  $b\mathbb{D} \setminus I$ , and  $U$  is an open neighborhood of  $I$  in  $\overline{\mathbb{D}}$ , then

- one can choose  $G$  to be  $\epsilon$ -close to  $F$  in the  $\mathcal{C}^1$  topology on  $\overline{\mathbb{D}} \setminus U$ .



# Riemann-Hilbert problem for null curves

## Theorem

Let  $M$  be a bordered Riemann surface and let  $F: M \rightarrow \mathbb{C}^3$  be a null holomorphic immersion (the *central curve*).

Let  $I$  be a compact arc in  $bM$  that is *not a connected component of  $bM$* .

Choose a small annular neighborhood  $A \subset M$  of the component  $C$  of  $bM$  containing  $I$  and a smooth retraction  $\rho: A \rightarrow C$ .

Let  $\vartheta \in \mathfrak{A}_*$  be a null vector (the *direction vector*), let  $\mu: C \rightarrow \mathbb{R}_+$  be a continuous function supported on  $I$  (the *size function*), and consider the continuous map

$$\begin{aligned} \varkappa: bM \times \overline{\mathbb{D}} &\rightarrow \mathbb{C}^3 \\ \varkappa(x, \xi) &= \begin{cases} F(x); & x \in bM \setminus I \\ F(x) + \mu(x) \xi \vartheta; & x \in I. \end{cases} \end{aligned}$$

## Theorem (Continued)

Then for any number  $\epsilon > 0$  there exist an arbitrarily small open neighborhood  $\Omega$  of  $I$  in  $A$  and a null holomorphic immersion  $G: M \rightarrow \mathbb{C}^3$  satisfying the following properties:

- $G$  is  $\epsilon$ -close to  $F$  in the  $\mathcal{C}^1$  topology on  $M \setminus \Omega$ .
  - $\text{dist}(G(x), \kappa(x, b\mathbb{D})) < \epsilon$  for all  $x \in bM$ .
  - $\text{dist}(G(x), \kappa(\rho(x), \overline{\mathbb{D}})) < \epsilon$  for all  $x \in \Omega$ .
- 
- We do not change the conformal structure on  $M$ .
  - $I$  can be replaced by a finite family of pairwise disjoint compact subarcs; it is allowed to use different direction vectors in each subarc.

# Sketch of the proof

- Solve the problem in a small disc  $D \subset \Omega \subset M \setminus A$  containing  $I$ . (Previous Theorem.) Let  $G_0: \overline{D} \rightarrow \mathbb{C}^3$  be a solution.
- Let  $\theta$  be a nowhere vanishing holomorphic 1-form on  $M$  and write

$$dG_0 = g_0\theta, \quad g_0: \overline{D} \rightarrow \mathfrak{A}_*,$$

$$dF = f_0\theta, \quad f_0: M \rightarrow \mathfrak{A}_*.$$

- Embed  $f_0$  as the core map of a **dominating and period dominating holomorphic spray** of maps (see Lecture 2):

$$\{f_t: M \rightarrow \mathfrak{A}_*\}_{t \in B}, \quad 0 \in B \subset \mathbb{C}^N \quad (N \text{ large}).$$

- Find a **dominating spray** of holomorphic maps

$$\{g_t: \overline{D} \rightarrow \mathfrak{A}_*\}_{t \in B}.$$

which are approximate solutions to the Riemann-Hilbert problem for the maps  $f_t$  in the spray  $\{f_t\}_{t \in B}$  in  $\overline{D}$ .

# Sketch of the proof

- **Glue** the two (dominating) sprays  $\{f_t\}_{t \in B}$  and  $\{g_t\}_{t \in B}$  into a single spray of holomorphic maps

$$\{h_t: M \rightarrow \mathfrak{A}_*\}_{t \in B}$$

such that  $h_t$  is close to  $g_t$  on  $\overline{D}$  and to  $f_t$  on  $\overline{M \setminus D}$ .

- Since  $f_0\theta$  is exact and the spray  $f_t$  is period dominating, so is  $h_t$  provided the approximation is close enough. Hence the period map

$$B \ni t \mapsto \mathcal{P}(h_t) = \left( \int_{\gamma} h_t \theta \right)_{\gamma \in \mathcal{H}_1(M, \mathbb{Z})}$$

has maximal rank at  $t = 0$ . The Implicit Function Theorem gives  $t_0 \in B$  close to 0 such that  $h_{t_0}\theta$  has vanishing periods.

- Fix a point  $p_0 \in M \setminus D$ . The map  $G: M \rightarrow \mathbb{C}^3$ ,

$$G(p) = F(p_0) + \int_{p_0}^p h_{t_0} \theta$$

proves the theorem.

# The conformal Calabi-Yau problem

## Theorem

Every bordered Riemann surface carries a **complete proper null holomorphic embedding** into a **ball** of  $\mathbb{C}^3$ .

## Corollary

Every bordered Riemann surface  $M$  carries a **complete conformal minimal immersion**  $M \rightarrow \mathbb{R}^3$  with **bounded image**.

- This answers a question by [Martín](#), [Umehara](#), and [Yamada](#) 2009.
- What is new in comparison to all existing results in the literature is that

**we do not change the complex structure  
on the source Riemann surface.**

(Compare with the work of [Nadirashvili](#), [Martín](#), [Morales](#), [López](#), [Alarcón](#), [Ferrer](#), [Umehara](#), [Yamada](#), [Meeks](#), etc.)

# Idea of the construction of complete bounded holomorphic immersions - Pythagoras' theorem

- Let  $F_0: M \rightarrow \mathbb{C}^n$  be a holomorphic immersion satisfying  $|F_0| \geq r_0 > 0$  on  $bM$ . We try to increase the boundary distance on  $M$  with respect to the induced metric by a fixed number  $\delta > 0$ .
- To this end, we approximate  $F_0$  uniformly on a compact set in  $M$  by an immersion  $F_1: M \rightarrow \mathbb{C}^n$  which at a point  $x \in bM$  adds a displacement for approximately  $\delta$  in a direction  $V \in \mathbb{C}^n$ ,  $|V| = 1$ , approximately orthogonal to the point  $F_0(x) \in \mathbb{C}^n$ . The boundary distance increases by  $\approx \delta$ , while the outer radius increases to

$$|F_1(x)| \approx \sqrt{|F_0(x)|^2 + \delta^2} \approx |F_0(x)| + \frac{\delta^2}{2|F_0(x)|} \leq |F_0(x)| + \frac{\delta^2}{2r_0}.$$

- By choosing a sequence  $\delta_j > 0$  such that  $\sum_j \delta_j = +\infty$  while  $\sum_j \delta_j^2 < \infty$ , we obtain by induction a limit immersion  $F: M \rightarrow \mathbb{C}^n$  with bounded outer radius and with complete metric  $F^* ds^2$ .

# The main tools

- This idea can be realized on short arcs  $I \subset bM$ , on which  $F_0$  does not vary too much, by solving a **Riemann-Hilbert problem**.
- Globally this method alone could lead to ‘sliding curtains’, creating shortcuts in the new induced metric on  $M$ .
- To **localize the problem** and **eliminate any shortcuts**, we subdivide  $bM = \cup_j I_j$  into a finite union of short arcs such that two adjacent arcs  $I_{j-1}, I_j$  meet at a common endpoint  $x_j$ . At the point  $p_j = F(x_j) \in \mathbb{C}^n$  we attach to  $F_0(M)$  a smooth real curve  $\lambda_j$  of length  $\delta$  whose other endpoint  $q_j$  increases the outer radius by  $\delta^2$ .
- By the method of **exposing boundary points** we modify the immersion so that  $F_0(x_j) = q_j$ . Hence any curve in  $M$  terminating on  $bM$  near  $x_j$  is elongated by approximately  $\delta > 0$ .
- In the next step we use a Riemann-Hilbert problem to increase the boundary distance on the arcs  $I_j$  by approximately  $\delta$ . These local modifications are glued together by the method of sprays.

# Null curves with a bounded coordinate

## Theorem

Every bordered Riemann surface  $M$  carries a *proper* holomorphic null embedding  $F = (F_1, F_2, F_3): M \rightarrow \mathbb{C}^3$  such that  $F_3$  is *bounded* on  $M$ . (Thus  $(F_1, F_2): M \rightarrow \mathbb{C}^2$  is a proper map.)

- These are the first examples of proper null curves in  $\mathbb{C}^3$  with a bounded coordinate function.
- Every open Riemann surface properly immerses in  $\mathbb{C}^2$  as a complex curve (Remmert 1956, Bishop 1960, and Narasimhan 1961).
- The only properly immersed minimal surfaces in  $\mathbb{R}^3$  contained in a half-space are planes (Hoffman and Meeks 1990). In particular, with  $\mathbf{F}$  as in the theorem,  $\Re(e^{it}\mathbf{F}): M \rightarrow \mathbb{R}^3$  is not proper for any  $t \in \mathbb{R}$ .
- Say a word about the proof.

# Null curves in $SL_2(\mathbb{C})$ and Bryant surfaces

- A **null curve** in the special linear group

$$SL_2(\mathbb{C}) = \left\{ z = \begin{pmatrix} z_{11} & z_{12} \\ z_{21} & z_{22} \end{pmatrix} \in \mathbb{C}^4 : \det z = z_{11}z_{22} - z_{12}z_{21} = 1 \right\}$$

is a holomorphic immersion  $F: M \rightarrow SL_2(\mathbb{C})$  of an open Riemann surface  $M$  into  $SL_2(\mathbb{C})$  which is directed by the quadric variety

$$\left\{ z = \begin{pmatrix} z_{11} & z_{12} \\ z_{21} & z_{22} \end{pmatrix} : \det z = z_{11}z_{22} - z_{12}z_{21} = 0 \right\} \subset \mathbb{C}^4.$$

- The projection of a null curve in  $SL_2(\mathbb{C})$  to the hyperbolic 3-space  $\mathbb{H}^3 = SL_2(\mathbb{C})/SU(2)$  is a conformal **Bryant surface** (a conformal immersion with constant mean curvature one in  $\mathbb{H}^3$ ).  
Bryant surfaces are closely related to minimal surfaces in  $\mathbb{R}^3$ .

# Null curves in $SL_2(\mathbb{C})$ and CMC-1 surfaces in $\mathbb{H}^3$

- The biholomorphic map  $\mathbf{T}: \mathbb{C}^3 \setminus \{z_3 = 0\} \rightarrow SL_2(\mathbb{C}) \setminus \{z_{11} = 0\}$ ,

$$\mathbf{T}(z_1, z_2, z_3) = \frac{1}{z_3} \begin{pmatrix} 1 & z_1 + iz_2 \\ z_1 - iz_2 & z_1^2 + z_2^2 + z_3^2 \end{pmatrix},$$

carries null curves to null curves (Martín, Umehara & Yamada 2009).

- If  $F = (F_1, F_2, F_3): M \rightarrow \mathbb{C}^3$  is a **proper** null curve such that  $c_1 < |F_3| < c_2$  on  $M$  for a pair of constants  $0 < c_1 < c_2$ , then  $\mathbf{T} \circ F: M \rightarrow SL_2(\mathbb{C})$  is a **proper** null curve in  $SL_2(\mathbb{C})$ .
- Since  $SU(2)$  is compact, **proper** null curves in  $SL_2(\mathbb{C})$  project into **proper** Bryant surfaces in  $\mathbb{H}^3$ .

## Corollary

*Every bordered Riemann surface carries a **proper holomorphic null embedding** into  $SL_2(\mathbb{C})$  and a **proper conformal Bryant surface** in  $\mathbb{H}^3$ .*

- These are the first examples of **proper** null curves in  $SL_2(\mathbb{C})$ , and Bryant surfaces in  $\mathbb{H}^3$ , with finite topology and **hyperbolic** conformal structure.
- Morales 2003 Analogous problem for minimal surfaces in  $\mathbb{R}^3$ .

# Final remarks and open problems

- Our method enables one to construct examples with **control on the conformal structure**. If we do not take care of the conformal structure, our method gives null curves in  $\mathbb{C}^3$  with **arbitrary topology**.
- Is **every open Riemann surface** conformally equivalent to a properly immersed Bryant surface in  $\mathbb{H}^3$ ?  
Does every open Riemann surface admit a proper holomorphic null embedding (or immersion) into  $SL_2(\mathbb{C})$ ?
- **Work in progress**: Every bordered Riemann surface admits a conformal complete proper minimal immersion into a ball of  $\mathbb{R}^3$ , and also a bounded complete immersion in  $\mathbb{R}^3$  which extends continuously to the boundary and maps the boundary onto a union of Jordan curves. There exists embeddings with these properties into  $\mathbb{R}^5$ .

# Minimal hulls of compact sets in $\mathbb{R}^3$

(With Barbara Drinovec Drnovšek.)

When discussing hulls in various geometries, one typically deals with dual sets of objects.

Given a set  $\mathcal{P}$  of real functions on a manifold  $X$ , the  $\mathcal{P}$ -hull of a compact subset  $K \subset X$  is

$$\widehat{K}_{\mathcal{P}} = \{x \in X : f(x) \leq \sup_K f \quad \forall f \in \mathcal{P}\}.$$

Suppose that  $\mathcal{G}$  is a class of geometric objects in  $X$  such that the restriction  $f|_C$  satisfies the maximum principle for every  $f \in \mathcal{P}$  and  $C \in \mathcal{G}$ . Then  $C \subset \widehat{K}_{\mathcal{P}}$  for every  $C \in \mathcal{G}$  with boundary  $\partial C \subset K$ .

The main question is how closely is the hull described by such objects.

A basic example is the convex hull  $\text{Co}(K)$  of a compact set in an affine space  $X \cong \mathbb{R}^n$ ; here  $\mathcal{P}$  is the class of all affine linear functions on  $X$  and  $\mathcal{G}$  is the collection of straight line segments in  $X$ .

# Polynomial hulls

Another classical case is the **polynomial hull**,  $\widehat{K}$ , of a compact set  $K \subset \mathbb{C}^n$ . This is the hull  $\widehat{K}_{\mathcal{P}}$  for  $\mathcal{P} = \{|f| : f \in \mathcal{O}(\mathbb{C}^n)\}$  or  $\mathcal{P} = \text{Psh}(\mathbb{C}^n)$  (plurisubharmonic functions). A natural dual class  $\mathcal{G}$  consists of complex curves.

Wermer (1958), Stolzenberg (1966), Alexander (1971): If  $K$  is a compact sets of finite linear measure then  $A = \widehat{K} \setminus K$  is a (possibly empty) one dimensional closed complex subvariety of  $\mathbb{C}^n \setminus K$ .

Stolzenberg' (1963): in general one cannot describe  $\widehat{K}$  by bounded complex curves with boundaries **exactly** in  $K$ .

## Theorem (Poletsky 1991, 1993)

*Let  $K$  be a compact set in  $\mathbb{C}^n$ , and let  $B \subset \mathbb{C}^n$  be a ball containing  $K$ . A point  $p \in B$  belongs to  $\widehat{K}$  iff there exists a sequence of holomorphic discs  $f_j: \overline{\mathbb{D}} \rightarrow B$  satisfying the following for every  $j = 1, 2, \dots$ :*

$$f_j(0) = p \text{ and } |\{t \in [0, 2\pi] : \text{dist}(f(e^{it}), K) < 1/j\}| \geq 2\pi - 1/j.$$

# Minimal plurisubharmonic functions

Is there a similar notion of the **minimal hull**,  $\widehat{K}_{\mathfrak{M}}$ , of a compact set  $K \subset \mathbb{R}^n$ ? Which class of function should one use?

## Definition (Harvey & Lawson; Drinovec Drnovšek & F.)

An upper semicontinuous function  $u$  on a domain  $\omega \subset \mathbb{R}^n$  ( $n \geq 3$ ) is **minimal plurisubharmonic ( $\mathfrak{M}$ -psh)** if the restriction of  $u$  to any affine 2-dimensional plane  $L \subset \mathbb{R}^n$  is subharmonic on  $L \cap \omega$  (in isothermal coordinates). The set of all such functions is denoted by  $\mathfrak{MPsh}(\omega)$ .

## Observations:

- The restriction of an  $\mathfrak{M}$ -psh function to an immersed minimal surface is subharmonic (in isothermal coordinates on the surface).
- A  $\mathcal{C}^2$  function  $u$  is  $\mathfrak{M}$ -psh if and only if the sum of the two smallest eigenvalues of  $\text{Hess } u$  is nonnegative at every point.
- A function  $u: \omega \rightarrow \mathbb{R}$  on  $\omega \subset \mathbb{R}^n$  is  $\mathfrak{M}$ -psh iff the function  $U: \omega \times \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $U(x + iy) = u(x)$ , is **null plurisubharmonic**, i.e., subharmonic on any null holomorphic curve.

# Minimal hulls

## Definition

The **minimal hull** of a compact set  $K \subset \mathbb{R}^n$  ( $n \geq 3$ ) is the set

$$\widehat{K}_{\text{MH}} = \{x \in \mathbb{R}^n : u(x) \leq \sup_K u \quad \forall u \in \mathfrak{MPsh}(\mathbb{R}^n)\}.$$

By the max principle, every minimal surface  $M$  with  $\partial M \subset K$  lies in  $\widehat{K}_{\text{MH}}$ .

## Theorem

Let  $K$  be a compact set in  $\mathbb{R}^3$ , and let  $\omega \Subset \mathbb{R}^3$  be a bounded open convex set containing  $K$ . A point  $p \in \omega$  belongs to the minimal hull  $\widehat{K}_{\text{MH}}$  if and only if there exists a sequence of conformal minimal discs  $f_j: \overline{\mathbb{D}} \rightarrow \omega$  such that for all  $j = 1, 2, \dots$  we have  $f_j(0) = p$  and

$$|\{t \in [0, 2\pi] : \text{dist}(f_j(e^{it}), K) < 1/j\}| \geq 2\pi - 1/j.$$

[Barbara Drinovec Drnovšek & Franc Forstnerič, Minimal hulls of compact sets in  $\mathbb{R}^3$ . <http://arxiv.org/abs/1409.6906>

# Largest $\mathfrak{M}$ -psh minorant

This is a consequence of the following result, analogous to the Poletsky and Bu-Schachermayer theorem (1991, 1993) for psh minorants.

**Theorem (B. Drinovec Drnovšek & Forstnerič, 2014)**

*Assume that  $\omega$  is a domain in  $\mathbb{R}^3$  and  $\phi: \omega \rightarrow \mathbb{R} \cup \{-\infty\}$  is an upper semicontinuous function. Then the function  $u: \omega \rightarrow \mathbb{R} \cup \{-\infty\}$ , given for any  $x \in \omega$  by*

$$u(x) = \inf \left\{ \int_0^{2\pi} \phi(f(e^{it})) \frac{dt}{2\pi} : f \in \text{CMI}(\overline{\mathbb{D}}, \omega), f(0) = x \right\}$$

*is minimal plurisubharmonic or identically  $-\infty$ ; moreover,  $u$  is the supremum of functions in  $\mathfrak{MPsh}(\omega)$  which are bounded above by  $\phi$ .*

The proof uses the approximate solution to the [Riemann-Hilbert problem for null holomorphic discs in  \$\mathbb{C}^3\$](#) , together with the relationship between  $\mathfrak{M}$ -psh functions on domains  $\omega \subset \mathbb{R}^3$  and null-psh functions on tubes  $\omega \times i\mathbb{R}^3 \subset \mathbb{C}^3$ .