

MINIMAL SURFACES IN MINIMALLY CONVEX DOMAINS

ANTONIO ALARCÓN, BARBARA DRINOVEC DRNOVŠEK, FRANC FORSTNERIČ,
AND FRANCISCO J. LÓPEZ

Dedicated to Josip Globevnik for his seventieth birthday

ABSTRACT. In this paper, we prove that every conformal minimal immersion of a compact bordered Riemann surface M into a minimally convex domain $D \subset \mathbb{R}^3$ can be approximated uniformly on compacts in $\dot{M} = M \setminus bM$ by proper complete conformal minimal immersions $\dot{M} \rightarrow D$. We also obtain a rigidity theorem for complete immersed minimal surfaces of finite total curvature contained in a minimally convex domain in \mathbb{R}^3 , and we characterize the minimal surface hull of a compact set K in \mathbb{R}^n for any $n \geq 3$ by sequences of conformal minimal discs whose boundaries converge to K in the measure theoretic sense.

1. INTRODUCTION

A major problem in minimal surface theory is to understand which domains in \mathbb{R}^3 admit complete properly immersed minimal surfaces and how the geometry of the domain influences the conformal properties of such surfaces. (For background on this topic, see e.g. [43, Section 3].) In the present paper, we obtain general existence and approximation results for complete proper conformal minimal immersions from an arbitrary bordered Riemann surface into any minimally convex domain in \mathbb{R}^3 ; see Theorems 1.1, 1.7, and 1.9. We also show that one cannot expect similar results in a wider class of domains in \mathbb{R}^3 .

Let $n \geq 3$. A domain $D \subset \mathbb{R}^n$ is said to be *minimally convex* if it admits a smooth exhaustion function $\rho: D \rightarrow \mathbb{R}$ that is *strongly 2-plurisubharmonic* (also called *minimal strongly plurisubharmonic*), meaning that for every point $\mathbf{x} \in D$, the sum of the smallest two eigenvalues of the Hessian $\text{Hess}_\rho(\mathbf{x})$ is positive. (See Definitions 2.1 and 2.3.) A domain D with \mathcal{C}^2 boundary is minimally convex if and only if $\kappa_1(\mathbf{x}) + \kappa_2(\mathbf{x}) \geq 0$ for each point $\mathbf{x} \in bD$, where $\kappa_1(\mathbf{x}) \leq \kappa_2(\mathbf{x}) \leq \dots \leq \kappa_{n-1}(\mathbf{x})$ are the normal curvatures of bD at the point \mathbf{x} with respect to the inner normal (see Theorem 1.2). In particular, a domain in \mathbb{R}^3 bounded by a properly embedded minimal surface is minimally convex (see Corollary 1.3). Clearly, every convex

Received by the editors August 5, 2016, and, in revised form, June 27, 2017.

2010 *Mathematics Subject Classification*. Primary 53A10; Secondary 32B15, 32E30, 32H02.

Key words and phrases. Riemann surface, minimal surface, minimally convex domain.

The first author was supported by the Ramón y Cajal program of the Spanish Ministry of Economy and Competitiveness.

The first and fourth authors were partially supported by the MINECO/FEDER grants MTM2014-52368 and MTM2017-89677-P, Spain.

The second and third authors were partially supported by the research program P1-0291 and grants J1-5432 and J1-7256 from ARRS, Republic of Slovenia.

domain is also minimally convex, but there exist minimally convex domains without any convex boundary points (see Example 1.4).

Our first main result is the following.

Theorem 1.1. *Assume that D is a minimally convex domain in \mathbb{R}^3 , and let M be a compact bordered Riemann surface with nonempty boundary bM . Then, every conformal minimal immersion $F_0: M \rightarrow D$ can be approximated uniformly on compacts in $\mathring{M} = M \setminus bM$ by proper complete conformal minimal immersions $F: \mathring{M} \rightarrow D$ with $\text{Flux}(F) = \text{Flux}(F_0)$.*

Recall that a *compact bordered Riemann surface* is a compact connected oriented surface, M , endowed with a complex structure whose boundary $bM \neq \emptyset$ consists of finitely many smooth Jordan curves. The interior, $\mathring{M} = M \setminus bM$, of such M is an (open) *bordered Riemann surface*. A *conformal minimal immersion* $F: M \rightarrow \mathbb{R}^n$ is an immersion which is angle preserving and harmonic; such a map parametrizes a minimal surface in \mathbb{R}^n . The *flux* of F is the group homomorphism $\text{Flux}(F): H_1(M, \mathbb{Z}) \rightarrow \mathbb{R}^n$ whose value on any closed oriented curve $\gamma \subset M$ is $\text{Flux}(F)(\gamma) = \oint_\gamma \Im(\partial F)$; here, ∂F is the $(1, 0)$ -differential of F and \Im denotes the imaginary part. An immersion $F: \mathring{M} \rightarrow \mathbb{R}^n$ is said to be *complete* if the pull-back F^*ds^2 of the Euclidean metric on \mathbb{R}^n is a complete Riemannian metric on \mathring{M} .

Note that Theorem 1.1 pertains to a fixed conformal structure on the surface M . The analogous result for *convex domains* in \mathbb{R}^n for any $n \geq 3$ is [2, Theorem 1.4]; see also [7, Theorem 1]. Theorem 1.1 seems to be the first general existence and approximation result for (complete) proper minimal surfaces in a class of domains in \mathbb{R}^3 which contains all convex domains, but also many nonconvex ones; convexity has been impossible to avoid with the existing construction methods. Comparing with the results in the literature, it is known that there are properly immersed minimal surfaces in \mathbb{R}^3 with arbitrary conformal structure (see [6, 8, 9]) and that every domain $D \subset \mathbb{R}^3$ which is convex, or has a smooth strictly convex boundary point, admits complete properly immersed minimal surfaces that are conformally equivalent to any given bordered Riemann surface (see [2]). These were the most general known results in this direction up to now.

As shown by Remark 1.11 and Examples 1.13 and 1.14, the hypothesis of minimal convexity is essentially optimal in Theorem 1.1. In Example 1.13 we exhibit a bounded, simply connected domain $D \subset \mathbb{R}^3$ such that a certain conformal minimal disc $F_0: \mathbb{D} \rightarrow D$ cannot be approximated by *proper* conformal minimal discs $\mathbb{D} \rightarrow D$. (Here, $\mathbb{D} = \{\zeta \in \mathbb{C} : |\zeta| < 1\}$.) In another direction, Martín, Meeks, and Nadirashvili constructed bounded (nonsimply connected) domains in \mathbb{R}^3 which do not admit any complete properly immersed minimal surfaces with an annular end (see [42]). We point out in Example 1.14 that there is a domain from [42] which does not admit any proper minimal discs. Clearly, Theorem 1.1 fails in both these examples even without the completeness condition.

In Remark 3.8, we indicate a generalization of Theorem 1.1, and of the related subsequent results in this paper, to a certain class of not necessarily convex domains in \mathbb{R}^n for any $n > 3$. However, we have optimal results only in dimension $n = 3$.

Theorem 1.1 is proved in Section 3; here is a brief outline. Let $\rho: D \rightarrow \mathbb{R}$ be a Morse minimal strongly plurisubharmonic exhaustion function with the (discrete)

critical locus P . For every point $\mathbf{x} \in D \setminus P$ we find a small embedded conformal minimal disc $\mathbf{x} \in \mathcal{M}_{\mathbf{x}} \subset D$ such that the restriction of ρ to $\mathcal{M}_{\mathbf{x}}$ has a strict minimum at \mathbf{x} and increases quadratically. Furthermore, for points in a simply connected compact set in $D \setminus P$ we can choose a smooth family of such discs satisfying uniform estimates for the rate of growth of ρ (see Lemma 3.1). By using these discs and an approximate solution of a Riemann-Hilbert type boundary value problem (see Theorem 3.2), we can lift the boundary of a given conformal minimal immersion $M \rightarrow D$ to a higher level of the function ρ , paying attention not to decrease the level of ρ much anywhere on M and to approximate the given immersion on a chosen compact subset of \dot{M} (see Proposition 3.3). This procedure can be carried out so that the image of the boundary bM avoids the critical locus of ρ . A recursive application of this lifting method leads to the construction of a proper conformal minimal immersion $\dot{M} \rightarrow D$. (Analogous results for proper holomorphic maps can be found in [18, 19].) This construction method is geometrically simpler than the one developed by the authors in [2], the main advantage being the higher flexibility of the Riemann-Hilbert method that is available in dimension $n = 3$ (compare Theorem 3.2 with [2, Theorem 3.5]).

Completeness of the immersion is achieved by combining the boundary lifting procedure with a technique, developed recently in [2], that enables one to increase the intrinsic boundary distance in M by an arbitrarily big amount while staying \mathcal{C}^0 close to a given conformal minimal immersion $M \rightarrow \mathbb{R}^n$ (see [2, Lemmas 4.1 and 4.2]). A recursive application of these two techniques yields Theorem 1.1. (See Section 3 for the details.)

Before proceeding, we place the class of minimal plurisubharmonic functions and minimally convex domains into a wider framework, and we provide some examples.

Minimal plurisubharmonic functions are a special case, with $p = 2$, of the class of p -*plurisubharmonic functions* which have been studied by Harvey and Lawson in [34]; see also [21, 31–33]. A real-valued \mathcal{C}^2 function u on a domain $D \subset \mathbb{R}^n$ is said to be (strongly) p -plurisubharmonic for some integer $p \in \{1, 2, \dots, n\}$ if the restriction of u to any p -dimensional affine subspace of \mathbb{R}^n is (strongly) subharmonic (see Definition 2.1); equivalently, if the sum of the p smallest eigenvalues of the Hessian of u is nonnegative (positive) at every point (see Proposition 2.2). The restriction of a p -plurisubharmonic function to a p -dimensional minimal submanifold is a subharmonic function on the submanifold (see Proposition 2.2). Note that 1-plurisubharmonic functions are convex functions, while n -plurisubharmonic functions are subharmonic functions. The set $\text{Psh}_p(D)$ of all p -plurisubharmonic functions on D is closed under addition and multiplication by nonnegative numbers.

A domain $D \subset \mathbb{R}^n$ is said to be p -convex if it admits a strongly p -plurisubharmonic exhaustion function $\rho: D \rightarrow \mathbb{R}$ (see Definition 2.3 and Proposition 2.6). Thus, 1-convex domains are linearly convex, while 2-convex domains are minimally convex. Every domain in \mathbb{R}^n is n -convex; this is a special case of a theorem of Greene and Wu [30] (see also Demailly [16]) that every connected noncompact Riemannian manifold admits a smooth strongly subharmonic exhaustion function. Harvey and Lawson proved that for smoothly bounded domains in \mathbb{R}^n , p -convexity is a local property of the boundary, akin to Levi pseudoconvexity in complex analysis. For future reference, we state the following summary of their main results from [34]. Harvey and Lawson considered bounded domains in \mathbb{R}^n , but we show in Section 2.3 that Theorem 1.2 also holds for unbounded domains.

Theorem 1.2 (Section 3 in [34]). *Let $1 \leq p < n$ be integers, and let $D \subset \mathbb{R}^n$ be a domain with \mathcal{C}^2 boundary, not necessarily bounded. The following conditions are equivalent.*

- (a) *D is p -convex.*
- (b) *There exist a neighborhood $U \subset \mathbb{R}^n$ of bD and a \mathcal{C}^2 function $\rho: U \rightarrow \mathbb{R}$ such that $D \cap U = \{\rho < 0\}$, $d\rho \neq 0$ on $bD \cap U = \{\rho = 0\}$, and*

$$(1.1) \quad \operatorname{tr}_L \operatorname{Hess}_\rho(\mathbf{x}) \geq 0 \quad \text{for every tangent } p\text{-plane } L \subset T_{\mathbf{x}}bD, \mathbf{x} \in bD.$$
(Here $\operatorname{Hess}_\rho(\mathbf{x})$ is the Hessian (2.1) of ρ at \mathbf{x} , and tr_L denotes the trace of the restriction to L .) Property (1.1) is independent of the choice of ρ .
- (c) *If $\mathbf{x} \in bD$ and $\kappa_1 \leq \kappa_2 \leq \dots \leq \kappa_{n-1}$ are the principal curvatures of bD from the inner side at \mathbf{x} , then $\kappa_1 + \kappa_2 + \dots + \kappa_p \geq 0$.*
- (d) *There exists a neighborhood U of bD such that the function $-\log \operatorname{dist}(\cdot, bD)$ is p -plurisubharmonic on $D \cap U$.*

Theorem 1.2 shows in particular that a domain $D \subset \mathbb{R}^3$ with \mathcal{C}^2 boundary is minimally convex if and only if the principal curvatures of the boundary bD satisfy $\kappa_1(\mathbf{x}) + \kappa_2(\mathbf{x}) \geq 0$ at every point $\mathbf{x} \in bD$. Theorem 1.1 applies to any such domain.

The following is a corollary to Theorem 1.2 in the case $D = \mathbb{R}^3$ (note that we have $\kappa_1 + \kappa_2 = 0$ on a minimal surface $S \subset \mathbb{R}^3$); the general case is proved in Section 2.2.

Corollary 1.3. *If S is a properly embedded minimal surface in \mathbb{R}^3 , then every connected component of $\mathbb{R}^3 \setminus S$ is a minimally convex domain. More generally, if $D \subset \mathbb{R}^3$ is a minimally convex domain and S is a closed embedded minimal surface in a neighborhood of \overline{D} , then every connected component of $D \setminus S$ is minimally convex.*

Example 1.4. Let D be the domain

$$D = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 > \cosh^2 z\}.$$

Since the boundary of D is a minimal surface (a catenoid), D is minimally convex by Corollary 1.3. Clearly, D does not have any convex boundary point, and its fundamental group $\pi_1(D)$ equals \mathbb{Z} .

Remark 1.5. Note that the Hessian of a minimal strongly plurisubharmonic function on a domain in \mathbb{R}^3 has at most one negative eigenvalue at every point. Hence, Morse theory implies that a minimally convex domain D has the homotopy type of a 1-dimensional CW complex; in particular, the higher homotopy groups $\pi_k(D)$ for $k > 1$ all vanish. Similarly, a p -convex domain has the homotopy type of a CW complex of dimension at most $p - 1$.

Remark 1.6. In the literature on minimal surfaces, a smoothly bounded domain D in \mathbb{R}^n is said to be (strongly) *mean-convex* if the sum of the principal curvatures of bD from the interior side is nonnegative (resp. positive) at each point. This is precisely condition (c) in Theorem 1.2 with $p = n - 1$; hence, a smoothly bounded domain in \mathbb{R}^n is mean-convex if and only if it is $(n - 1)$ -convex. In particular, mean-convex domains in \mathbb{R}^3 coincide with smoothly bounded minimally convex domains. Mean-convex domains have been studied as natural barriers for minimal hypersurfaces in view of the maximum principle; see Section 2.4 and Remark 5.7. Nontrivial proper minimal hypersurfaces in mean-convex domains often arise as solutions to Plateau problems. For instance, Meeks and Yau [45] proved that every

null-homotopic Jordan curve in the boundary of a mean-convex domain $D \subset \mathbb{R}^3$ bounds an area minimizing minimal disc in D . This method does not seem to provide examples of complete minimal surfaces or those normalized by a given bordered Riemann surface other than the disc. For a discussion of this subject, see e.g. [15, Section 6.5].

Our proof of Theorem 1.1 also shows that boundaries of conformal minimal surfaces can be pushed to a *minimally convex end* of a domain $D \subset \mathbb{R}^3$ as in the following theorem. An analogous result in the holomorphic category is [19, Theorem 1.1].

Theorem 1.7. *Assume that $\Omega \subset D$ are open sets in \mathbb{R}^3 and that $\rho: \Omega \rightarrow (0, +\infty)$ is a smooth minimal strongly plurisubharmonic function such that for any pair of numbers $0 < c_1 < c_2$ the set $\Omega_{c_1, c_2} = \{\mathbf{x} \in \Omega: c_1 \leq \rho(\mathbf{x}) \leq c_2\}$ is compact. Let M be a compact bordered Riemann surface with nonempty boundary bM . Every conformal minimal immersion $F_0: M \rightarrow D$ satisfying $F_0(bM) \subset \Omega$ can be approximated uniformly on compacts in $\dot{M} = M \setminus bM$ by complete conformal minimal immersions $F: \dot{M} \rightarrow D$ such that $F(z) \in \Omega$ for every $z \in \dot{M}$ sufficiently close to bM and*

$$(1.2) \quad \lim_{z \rightarrow bM} \rho(F(z)) = +\infty.$$

In a typical application of Theorem 1.7, the set Ω is a collar around a minimally convex boundary component $S \subset bD$. (By Theorem 1.2, a smooth boundary component $S \subset bD$ is minimally convex if and only if $-\log \text{dist}(\cdot, S)$ is minimal plurisubharmonic near S .) Theorem 1.7 furnishes a proper complete conformal minimal immersion $F: \dot{M} \rightarrow D$ whose boundary cluster set is contained in S as shown by condition (1.2).

Next, we consider the class of strongly minimally convex domains.

Definition 1.8. A domain $D \subset \mathbb{R}^n$ with \mathcal{C}^2 boundary is *strongly p -convex* for some $p \in \{1, \dots, n - 1\}$ if it admits a \mathcal{C}^2 defining function ρ on a neighborhood U of bD (i.e., $D \cap U = \{\rho < 0\}$ and $d\rho \neq 0$ on $bD = \{\mathbf{x} \in U: \rho(\mathbf{x}) = 0\}$) whose Hessian satisfies the strict inequality in (1.1):

$$\text{tr}_L \text{Hess}_\rho(\mathbf{x}) > 0 \quad \text{for every tangent } p\text{-plane } L \subset T_{\mathbf{x}}bD, \mathbf{x} \in bD.$$

A strongly 2-convex domain is said to be *strongly minimally convex*.

The analogue of Theorem 1.2 holds in this setting. In particular, a bounded domain $D \Subset \mathbb{R}^n$ with \mathcal{C}^2 boundary is strongly p -convex for some $p \in \{1, \dots, n - 1\}$ if and only if the principal curvatures $\kappa_1 \leq \kappa_2 \leq \dots \leq \kappa_{n-1}$ of bD at any point $\mathbf{x} \in bD$ satisfy $\kappa_1 + \kappa_2 + \dots + \kappa_p > 0$. Note that D is strongly $(n - 1)$ -convex if and only if it is strongly mean-convex (see Remark 1.6).

Our next result improves Theorem 1.1 for bounded strongly minimally convex domains.

Theorem 1.9. *Let D be a bounded strongly minimally convex domain in \mathbb{R}^3 (Definition 1.8). Given a compact bordered Riemann surface M with nonempty boundary bM and a conformal minimal immersion $F_0: M \rightarrow D$, we can approximate F_0 uniformly on compacts in \dot{M} by continuous maps $F: M \rightarrow \overline{D}$ such that*

$F(bM) \subset bD$, $F: \overset{\circ}{M} \rightarrow D$ is a proper complete conformal minimal immersion, $\text{Flux}(F_0) = \text{Flux}(F)$, and

$$(1.3) \quad \sup_{\zeta \in \overset{\circ}{M}} \|F(\zeta) - F_0(\zeta)\| \leq C \sqrt{\max_{\zeta \in bM} \text{dist}(F_0(\zeta), bD)}$$

for some constant $C > 0$ depending only on D .

The improvement over Theorem 1.1 is that the approximating map F can now be chosen continuous up to the boundary of M , so $F(bM) \subset bD$ is a union of finitely many closed curves, and we have the estimate (1.3). Since F is complete, the minimal surface $F(\overset{\circ}{M}) \subset D$ has infinite area, and hence its boundary $F(bM)$ is necessarily nonrectifiable in view of the isoperimetric inequality. The corresponding result for smoothly bounded strongly convex domains in \mathbb{R}^n for any $n \geq 3$ is [2, Theorem 1.2]; see also [1] for a previous partial result in this line. As in the latter result, we are unable to achieve that F is a topological embedding on bM , so $F(bM)$ needs not consist of Jordan curves.

Theorem 1.9 implies the following corollary.

Corollary 1.10. *Every domain $D \subset \mathbb{R}^3$ having a \mathcal{C}^2 strongly minimally convex boundary point contains complete properly immersed minimal surfaces extending continuously up to the boundary and normalized by any given bordered Riemann surface.*

Proof. Assume that $\mathbf{x}_0 \in bD$ is a strongly minimally convex boundary point, i.e., such that $\kappa_1(\mathbf{x}_0) + \kappa_2(\mathbf{x}_0) > 0$. Then there are a neighborhood U of \mathbf{x}_0 and a strongly minimally convex domain $D' \subset D$ such that $D \cap U = D' \cap U$. (It suffices to intersect D by a small ball around \mathbf{x}_0 and smooth the corners.) Given a conformal minimal immersion $F_0: M \rightarrow D'$ whose image $F_0(M)$ lies close enough to the point \mathbf{x}_0 (such exists since M is compact), the map $F: M \rightarrow \overline{D}'$, furnished by Theorem 1.9, satisfies $F(bM) \subset bD \cap U$ in view of the estimate (1.3), and hence the map $F: \overset{\circ}{M} \rightarrow D$ is proper. \square

Following Meeks and Pérez [43, Section 3], a domain $W \subset \mathbb{R}^3$ is said to be *universal for minimal surfaces* if every complete, connected, properly immersed minimal surface in W is either recurrent (when the surface is open) or a parabolic surface with boundary. Since every open bordered Riemann surface $\overset{\circ}{M} = M \setminus bM$ is transient, Theorem 1.7 and Corollary 1.10 show that every domain $D \subset \mathbb{R}^3$ which has a minimally convex end or a strongly minimally convex boundary point fails to be universal for minimal surfaces. In particular, there are domains in \mathbb{R}^3 which are not universal for minimal surfaces and have no convex boundary points; for example, the catenoidal domain in Example 1.4.

Remark 1.11. The conclusion of Theorem 1.7 fails along a compact smooth boundary component $S \subset bD$ which is *strongly minimally concave*; i.e., $\kappa_1(\mathbf{x}) + \kappa_2(\mathbf{x}) < 0$ for every point $\mathbf{x} \in S$. Indeed, Theorem 1.2 furnishes an open neighborhood U of S in \mathbb{R}^3 and a minimal strongly plurisubharmonic function $\phi: U \rightarrow \mathbb{R}$ that vanishes on S and is positive on $D \cap U$. The maximum principle applied with ϕ shows that there is no minimal surface in $\overline{D} \cap U$ with boundary in S . The same argument holds locally near a smooth strongly minimally concave boundary point $\mathbf{x}_0 \in bD$. In this case there is a neighborhood $U \subset \mathbb{R}^3$ of \mathbf{x}_0 such that there are no proper minimal surfaces in $D \cap U$ with boundary in $bD \cap U$.

For a *complete* proper minimal surface there is another restriction on the location of its boundary points. Assume that $D \subset \mathbb{R}^3$ is a domain with \mathcal{C}^2 boundary and $F: \mathbb{D} \rightarrow D$ is a complete conformal proper minimal immersion from the disc $\mathbb{D} = \{\zeta \in \mathbb{C} : |\zeta| < 1\}$ extending continuously to $\overline{\mathbb{D}}$. Then the boundary curve $F(b\mathbb{D}) \subset bD$ does not contain any strongly concave boundary points of D (see [10]). This is especially relevant in connection to Theorem 1.9. However, we do not know whether $F(b\mathbb{D})$ could contain a strongly minimally concave boundary point; the following remains an open problem.

Problem 1.12. Let D be a smoothly bounded domain in \mathbb{R}^3 and let $F: \mathbb{D} \rightarrow D$ be a complete conformal proper minimal immersion extending continuously to $\overline{\mathbb{D}}$ (hence $F(b\mathbb{D}) \subset bD$). Do we have $\kappa_1(\mathbf{x}) + \kappa_2(\mathbf{x}) \geq 0$ for every point $\mathbf{x} \in F(b\mathbb{D})$?

We now illustrate by a couple of examples that Theorem 1.1 fails in general for domains in \mathbb{R}^3 which are not minimally convex. Since every domain in \mathbb{R}^3 is 3-convex (i.e., it admits a strongly subharmonic exhaustion function; see [16, 30]), we see in particular that the hypothesis of 2-convexity cannot be replaced by 3-convexity in Theorem 1.1.

Example 1.13. We exhibit a simply connected domain $D \subset \mathbb{R}^3$ of the form $D = 2\mathbb{B} \setminus K$, where \mathbb{B} is the unit ball of \mathbb{R}^3 and K is a compact set contained in a thin shell around the unit sphere $\mathbb{S} = b\mathbb{B}$, such that the image of every proper conformal minimal disc $F: \mathbb{D} \rightarrow D$ avoids the ball $\frac{1}{2}\mathbb{B} \subset D$. Clearly, Theorem 1.1 fails in this example even without the completeness condition. Note however that D admits complete properly immersed minimal surfaces normalized by any given bordered Riemann surface in view of Theorem 1.7 applied to the strongly convex boundary component $2\mathbb{S} \subset bD$.

The example is essentially the one given in [27, Section 5] in the context of holomorphic discs in domains in \mathbb{C}^2 . We cover the unit sphere $\mathbb{S} \subset \mathbb{R}^3$ by small open spherical caps C_1, \dots, C_m (i.e., every C_j is the intersection of \mathbb{S} by a half-space defined by an affine plane $H_j \subset \mathbb{R}^3$) such that $\bigcup_{j=1}^m \text{Co}(\overline{C}_j) \cap \frac{1}{2}\overline{\mathbb{B}} = \emptyset$. (Here, Co denotes the convex hull.) Pick a number $r > 1$ so close to 1 that $\mathbb{S} \subset \bigcup_{j=1}^m \text{Co}(rC_j)$. Choose pairwise distinct numbers ρ_1, \dots, ρ_m very close to r such that the pairwise disjoint spherical caps $\Gamma_j = \rho_j C_j$ satisfy $\mathbb{S} \subset \bigcup_{j=1}^m \text{Co}(\Gamma_j)$ and $\bigcup_{j=1}^m \text{Co}(\overline{\Gamma}_j) \cap \frac{1}{2}\overline{\mathbb{B}} = \emptyset$. Let $D = 2\mathbb{B} \setminus \bigcup_{j=1}^m \overline{\Gamma}_j$. For any proper conformal minimal disc $F: \mathbb{D} \rightarrow D$, its boundary cluster set $\Lambda(F)$ (i.e., the set of all limit points $\lim_{j \rightarrow \infty} F(\zeta_j) \in bD$ along sequences $\zeta_j \in \mathbb{D}$ with $\lim_{j \rightarrow \infty} |\zeta_j| = 1$) is a connected compact set in bD ; hence it is contained in the sphere $2\mathbb{S}$ or in one of the caps Γ_j . Assume now that $F(\zeta_0) \in \frac{1}{2}\overline{\mathbb{B}}$ for some $\zeta_0 \in \mathbb{D}$. If $\Lambda(F) \subset \Gamma_j$ for some $j \in \{1, \dots, m\}$, then $F(\mathbb{D}) \subset \text{Co}(\Gamma_j)$ by the maximum principle, a contradiction since $\text{Co}(\Gamma_j)$ does not intersect the ball $\frac{1}{2}\overline{\mathbb{B}}$. If on the other hand $\Lambda(F) \subset 2\mathbb{S}$, there is a point $\zeta_1 \in \mathbb{D}$ with $F(\zeta_1) \in \mathbb{S}$. Pick $j \in \{1, \dots, m\}$ such that $F(\zeta_1) \in \text{Co}(\Gamma_j)$. Since F has no cluster points on Γ_j , the set $U = \{\zeta \in \mathbb{D} : F(\zeta) \in \text{Co}(\Gamma_j)\}$ is a nonempty relatively compact domain in \mathbb{D} , and $F(bU)$ lies in the affine plane H_j which determines the spherical cap Γ_j . By the maximum principle it follows that F maps all of U , and hence the whole disc \mathbb{D} , into H_j , a contradiction.

Martín, Meeks, and Nadirashvili constructed bounded domains in \mathbb{R}^3 which do not admit any proper complete minimal surfaces of finite topology (see [42]). In the next example we show that the collection in [42] includes a domain $D \subset \mathbb{R}^3$ which

carries no proper minimal discs, irrespective of completeness. A similar result in the holomorphic category is due to Dor [17], who constructed a bounded domain D in \mathbb{C}^m for any $m \geq 2$ which does not admit any proper holomorphic discs.

Example 1.14. Let S be the cylindrical shell

$$S = \left\{ (x, y, z) \in \mathbb{R}^3 : 1 < \|(x, y)\| = \sqrt{x^2 + y^2} < 2, 0 < z < 1 \right\}.$$

For $0 < t < 1$, let $C_t := S \cap \{(x, y, z) \in \mathbb{R}^3 : z = t\}$ denote the planar round open annulus obtained by intersecting the cylinder S with the plane $z = t$. For $j \in \mathbb{N}$, denote by $C_{t,j}$ the planar round compact annulus $C_{t,j} = \{(x, y, z) \in C_t : 1 + \frac{1}{2j} \leq \|(x, y)\| \leq 2 - \frac{1}{2j}\}$. Obviously, $bC_{t,j} = \{(x, y, z) \in \mathbb{R}^3 : \|(x, y)\| \in \{1 + \frac{1}{2j}, 2 - \frac{1}{2j}\}, z = t\}$. Let t_1, t_2, t_3, \dots denote the sequence

$$\frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{4}, \frac{2}{4}, \frac{3}{4}, \dots, \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n}, \dots.$$

Set $\Gamma = \bigcup_{j \in \mathbb{N}} bC_{t_j,j} \subset S$ and $D = S \setminus \Gamma$. By [42, Proof of Theorem 1], D is a domain in \mathbb{R}^3 , and the boundary cluster set $\Lambda(E) \subset \overline{D} \setminus D$ of any proper minimal annular end $E \subset D$ lies in a horizontal plane of \mathbb{R}^3 . By the maximum principle, this implies that every proper minimal disc $\mathbb{D} \rightarrow D$ is contained in a horizontal plane, but clearly D does not admit any such discs. More generally, D does not carry any proper minimal surfaces of finite genus and with a single end.

All minimal surfaces in Theorems 1.1, 1.7, and 1.9 are images of bordered Riemann surfaces, hence finitely connected. If one does not insist on the approximation and the control of the conformal structure on these surfaces, then our methods also give complete proper minimal surfaces of arbitrary topological type.

Corollary 1.15. *If D is a domain in \mathbb{R}^3 which has a minimally convex end in the sense of Theorem 1.7 or a strongly minimally convex boundary point, then every open orientable smooth surface S carries a complete proper minimal immersion $S \rightarrow D$ with arbitrary flux.*

Corollary 1.15 is proved at the end of Section 3. For domains $D \subset \mathbb{R}^n$ ($n \geq 3$) that are convex or have a \mathcal{C}^2 smooth strictly convex boundary point, this has already been established in [2]; for $n = 3$ see also Ferrer, Martín, and Meeks [25].

Theorems 1.1, 1.7, and 1.9 show that every minimally convex domain in \mathbb{R}^3 admits many complete properly immersed minimal surfaces of *hyperbolic* conformal type. In contrast, the following rigidity type result shows that only very few minimally convex domains contain a complete proper minimal surface $S \subset \mathbb{R}^3$ of *finite total curvature*. (Note that these are the simplest complete minimal surfaces of *parabolic* conformal type.)

Theorem 1.16. *Let $S \subset \mathbb{R}^3$ be a complete connected properly immersed minimal surface with finite total curvature in \mathbb{R}^3 . If $D \subset \mathbb{R}^3$ is a smoothly bounded minimally convex domain containing S , then $D = \mathbb{R}^3$ or S is a plane; in the latter case, the connected component of D containing S is a slab, a half-space, or \mathbb{R}^3 .*

By a *slab* in \mathbb{R}^3 , we mean a domain bounded by two parallel planes.

In particular, if D is a connected component of $\mathbb{R}^3 \setminus S$ where S is a nonflat properly embedded minimal surface of finite total curvature in \mathbb{R}^3 , then Theorem 1.16 shows that D is a *maximal minimally convex domain*, in the sense that the only smoothly bounded minimally convex domain containing \overline{D} is \mathbb{R}^3 itself.

Theorem 1.16 is proved in Section 4 as an application of a general maximum principle at infinity for complete, finite total curvature, noncompact minimal surfaces with compact boundary in minimally convex domains of \mathbb{R}^3 ; see Theorem 4.1. Maximum principles at infinity have been the key in many celebrated classification results in the theory of minimal surfaces; see for instance [44] and the references therein. In the proof of Theorem 4.1, we exploit the geometry of complete minimal surfaces of finite total curvature along with the *Kontinuitätssatz* for conformal minimal surfaces; see Proposition 2.9 for the latter.

In Section 5, we indicate how the Riemann-Hilbert technique, developed in [2], allows us to extend all main results of the paper [21] to null hulls of compact sets in \mathbb{C}^n (see Definition 5.4) and minimal hulls of compact sets in \mathbb{R}^n (see Definition 2.5) for any $n \geq 3$.

After the completion of this paper, Alarcón, Forstnerič, and López obtained analogues of Theorems 1.1 and 1.9 in the nonorientable framework (see [4]).

2. p -PLURISUBHARMONIC FUNCTIONS AND p -CONVEX DOMAINS

We begin this preparatory section by summarizing basic results concerning p -plurisubharmonic functions and p -convex domains in \mathbb{R}^n which are used in the paper, referring to the papers of Harvey and Lawson [32–34] and the references therein for a more complete account. We add the proof of Theorem 1.2 for unbounded domains (see Subsection 2.3) and formulate the *Kontinuitätssatz* for minimal submanifolds (see Proposition 2.9). In Subsection 2.5, we recall the notion of a *null plurisubharmonic function* and develop one of the main tools that will be used in the proof of Theorems 1.1, 1.7, and 1.9.

We denote by $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$ the standard Euclidean inner product and the Euclidean norm on \mathbb{R}^n , respectively. We shall use the same notation for the Euclidean norm on \mathbb{C}^n .

2.1. p -plurisubharmonic functions. Let $\mathbf{x} = (x_1, \dots, x_n)$ be coordinates on \mathbb{R}^n . Given a domain $D \subset \mathbb{R}^n$ and a \mathcal{C}^2 function $u: D \rightarrow \mathbb{R}$, the *Hessian* of u at a point $\mathbf{x} \in D$ is the quadratic form $\text{Hess}_u(\mathbf{x}) = \text{Hess}_u(\mathbf{x}; \cdot)$ on the tangent space $T_{\mathbf{x}}\mathbb{R}^n \cong \mathbb{R}^n$, given by

$$(2.1) \quad \text{Hess}_u(\mathbf{x}; \xi) = \sum_{j,k=1}^n \frac{\partial^2 u}{\partial x_j \partial x_k}(\mathbf{x}) \xi_j \xi_k, \quad \xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n.$$

The trace of the Hessian is the Laplace operator on \mathbb{R}^n : $\text{tr}(\text{Hess}_u) = \Delta u = \sum_{j=1}^n \frac{\partial^2 u}{\partial x_j^2}$.

The Euclidean metric $ds^2 = \sum_{j=1}^n dx_j \otimes dx_j$ on \mathbb{R}^n induces a Riemannian metric $g = g_M$ on any smoothly immersed submanifold $M \rightarrow \mathbb{R}^n$. A function $u \in \mathcal{C}^2(D)$ is subharmonic on a submanifold $M \subset D$ if $\Delta_M(u|_M) \geq 0$, where Δ_M is the Laplace operator on M associated to the metric g_M induced by the immersion. In particular, if L is an affine p -dimensional subspace of \mathbb{R}^n given by

$$L = \left\{ \mathbf{x}(\xi) = \mathbf{a} + \sum_{j=1}^p \xi_j \mathbf{v}_j \in \mathbb{R}^n : \xi_1, \dots, \xi_p \in \mathbb{R} \right\},$$

where $\mathbf{a} \in \mathbb{R}^n$ and $\mathbf{v}_1, \dots, \mathbf{v}_p \in \mathbb{R}^n$ is an orthonormal set, then u is subharmonic on $L \cap D$ if and only if the function $\xi \mapsto u(\mathbf{x}(\xi))$ is subharmonic on $\{\xi \in \mathbb{R}^p : \mathbf{x}(\xi) \in D\}$.

Definition 2.1. An upper semicontinuous function $u: D \rightarrow \mathbb{R} \cup \{-\infty\}$ on a domain $D \subset \mathbb{R}^n$ is p -plurisubharmonic for some integer $p \in \{1, \dots, n\}$ if the restriction $u|_{L \cap D}$ to any affine p -dimensional plane $L \subset \mathbb{R}^n$ is subharmonic on $L \cap D$. A 2-plurisubharmonic function is also called *minimal plurisubharmonic*.

The set of all p -plurisubharmonic functions on D is denoted by $\text{Psh}_p(D)$. Following the notation introduced in [21], we shall write

$$\text{Psh}_2(D) = \mathfrak{MPsh}(D).$$

It is obvious that $\text{Psh}_1(D) \subset \text{Psh}_2(D) \subset \dots \subset \text{Psh}_n(D)$. An n -plurisubharmonic function on a domain $D \subset \mathbb{R}^n$ is a subharmonic function in the usual sense, and a 1-plurisubharmonic function is a convex function. Clearly, $\text{Psh}_p(D)$ is closed under addition and multiplication by nonnegative real numbers. Most of the familiar properties of plurisubharmonic functions on domains in \mathbb{C}^n extend to p -plurisubharmonic functions on domains in \mathbb{R}^n (see e.g. [32, Section 6]). In particular, every p -plurisubharmonic function can be approximated by smooth p -plurisubharmonic functions.

Proposition 2.2 (Proposition 2.3 and Theorem 2.13 in [34]). *Let $1 \leq p \leq n$ be integers and let D be a domain in \mathbb{R}^n . The following conditions are equivalent for a function $u \in \mathcal{C}^2(D)$:*

- (a) u is p -plurisubharmonic on D .
 - (b) $\text{tr}_L \text{Hess}_u(\mathbf{x}) \geq 0$ for every point $\mathbf{x} \in D$ and every p -dimensional linear subspace $L \subset \mathbb{R}^n$ (here, tr_L denotes the trace of the restriction to L).
 - (c) If $\lambda_1(\mathbf{x}) \leq \lambda_2(\mathbf{x}) \leq \dots \leq \lambda_n(\mathbf{x})$ are the eigenvalues of $\text{Hess}_u(\mathbf{x})$, then
- (2.2)
$$\lambda_1(\mathbf{x}) + \dots + \lambda_p(\mathbf{x}) \geq 0 \quad \text{for every } \mathbf{x} \in D.$$
- (d) $u|_M$ is subharmonic on every minimal p -dimensional submanifold $M \subset D$.

Sketch of proof. The equivalences (a) \Leftrightarrow (b) \Leftrightarrow (c) are easily seen, and (d) \Rightarrow (a) is obvious. The nontrivial implication (b) \Rightarrow (d) follows from the following formula which holds for every smooth submanifold $M \subset \mathbb{R}^n$ (cf. [31, Proposition 2.10]):

(2.3)
$$\Delta_M(u|_M) = \text{tr}_M \text{Hess}_u - H_M u.$$

Here, $\text{tr}_M \text{Hess}_u$ is the trace of the restriction of the Hessian of u to the tangent bundle of M , and H_M is the mean curvature vector field of M . If M is a minimal submanifold, then $H_M = 0$, and we get that $\Delta_M(u|_M) = \text{tr}_M \text{Hess}_u \geq 0$. \square

Definition 2.3. A function $u \in \mathcal{C}^2(D)$ on a domain $D \subset \mathbb{R}^n$ is *strongly p -plurisubharmonic* if $\text{tr}_L \text{Hess}_u(\mathbf{x}) > 0$ for every p -dimensional affine linear subspace $L \subset \mathbb{R}^n$ and every point $\mathbf{x} \in D \cap L$. Equivalently, if $\lambda_1(\mathbf{x}) \leq \lambda_2(\mathbf{x}) \leq \dots \leq \lambda_n(\mathbf{x})$ are the eigenvalues of $\text{Hess}_u(\mathbf{x})$, then $\lambda_1(\mathbf{x}) + \dots + \lambda_p(\mathbf{x}) > 0$ for all $\mathbf{x} \in D$.

The analogue of Proposition 2.2 holds for strongly p -plurisubharmonic functions; in particular, we have the following result.

Proposition 2.4. *A function $u \in \mathcal{C}^2(D)$ on a domain $D \subset \mathbb{R}^n$ is strongly p -plurisubharmonic if and only if $u|_M$ is strongly subharmonic on every minimal p -dimensional submanifold $M \subset D$.*

Observe that for any $u \in \text{Psh}_p(D) \cap \mathcal{C}^2(D)$, the function $u(\mathbf{x}) + \epsilon \|\mathbf{x}\|^2$ is strongly p -plurisubharmonic for every $\epsilon > 0$. It follows that every p -plurisubharmonic function can be approximated by smooth strongly p -plurisubharmonic functions.

If h is a smooth real function on \mathbb{R} and u is a \mathcal{C}^2 function on a domain $D \subset \mathbb{R}^n$, then for each point $\mathbf{x} \in D$ and vector $\xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n$ we have

$$(2.4) \quad \text{Hess}_{h \circ u}(\mathbf{x}, \xi) = h'(u(\mathbf{x})) \text{Hess}_u(\mathbf{x}, \xi) + h''(u(\mathbf{x})) \|\nabla u(\mathbf{x}) \cdot \xi\|^2.$$

It follows that if u is (strongly) p -plurisubharmonic and h is (strongly) increasing and convex on the range of u , then $h \circ u$ is also (strongly) p -plurisubharmonic.

2.2. p -convex hulls and p -convex domains.

Definition 2.5 (Definitions 3.1 and 3.3 in [34]). Let K be a compact set in a domain $D \subset \mathbb{R}^n$ and let $p \in \{1, 2, \dots, n\}$. The p -convex hull (or the p -hull) of K in D is the set

$$\widehat{K}_{p,D} = \{\mathbf{x} \in D : u(\mathbf{x}) \leq \sup_K u \text{ for all } u \in \text{Psh}_p(D)\}.$$

We shall write $\widehat{K}_p = \widehat{K}_{p,\mathbb{R}^n}$. The 2-hull is also called the *minimal hull* and denoted

$$\widehat{K}_{\mathfrak{M},D} = \widehat{K}_{2,D}; \quad \widehat{K}_{\mathfrak{M}} = \widehat{K}_{\mathfrak{M},\mathbb{R}^n}.$$

A domain $D \subset \mathbb{R}^n$ is p -convex if $\widehat{K}_{p,D}$ is compact for every compact set $K \subset D$. A 2-convex domain is also called *minimally convex*.

Since $\text{Psh}_p(D) \subset \text{Psh}_{p+1}(D)$ for $p = 1, \dots, n-1$, we have $K \subset \widehat{K}_n \subset \dots \subset \widehat{K}_2 \subset \widehat{K}_1 = \text{Co}(K)$. Simple examples show that these inclusions are strict in general.

The following result is [34, Theorem 3.4]; the proof is similar to the classical one concerning holomorphically convex domains in \mathbb{C}^n .

Proposition 2.6. *A domain $D \subset \mathbb{R}^n$ is p -convex for some $p \in \{1, 2, \dots, n\}$ (see Definition 2.5) if and only if it admits a smooth strongly p -plurisubharmonic exhaustion function.*

The proof of the next result follows the familiar case of plurisubharmonic functions; see e.g. Hörmander [35, Theorem 5.1.5, p. 117].

Proposition 2.7. *Let D be a p -convex domain in \mathbb{R}^n , and let $K \subset D$ be a compact p -convex set, i.e., $K = \widehat{K}_{p,D}$. Then the following conditions hold.*

- (a) *There exists a smooth p -plurisubharmonic exhaustion function $\rho : D \rightarrow \mathbb{R}_+$ such that $\rho^{-1}(0) = K$ and ρ is strongly p -plurisubharmonic on $D \setminus K$.*
- (b) *For every p -plurisubharmonic function u on a neighborhood U of K there exists a p -plurisubharmonic exhaustion function $v : D \rightarrow \mathbb{R}$ which agrees with u on K and is smooth strongly p -plurisubharmonic on $D \setminus K$.*

Proof of (a). For any point $\mathbf{x} \in D \setminus K$ there exists a smooth strongly p -plurisubharmonic function u on D such that $u < 0$ on K and $u(\mathbf{x}) > 0$. Pick a smooth function $h : \mathbb{R} \rightarrow \mathbb{R}_+$ which equals zero on $(-\infty, 0]$ and is strongly increasing and strongly convex on $(0, \infty)$. Then $h \circ u \geq 0$ vanishes on a neighborhood of K and is strongly p -plurisubharmonic on a neighborhood V of \mathbf{x} in view of (2.4). Hence we can pick a countable collection $\{(V_j, u_j)\}_{j \in \mathbb{N}}$, where V_j is an open set in $D \setminus K$, $u_j \geq 0$ is a smooth p -plurisubharmonic function on D that vanishes near K and is strongly p -plurisubharmonic on V_j , and $\bigcup_{j=1}^\infty V_j = D \setminus K$. If the numbers $\epsilon_j > 0$ are chosen small enough, then the series $v = \sum_{j=1}^\infty \epsilon_j u_j \geq 0$ converges in $\mathcal{C}^\infty(D)$. By the construction, v vanishes precisely on K and is strongly p -plurisubharmonic on $D \setminus K$. Finally, take $\rho = v + h \circ \tau$, where τ is a smooth p -plurisubharmonic exhaustion function on D that is negative on K .

Proof of (b). We may assume that \bar{U} is compact. Choose a smooth function χ on \mathbb{R}^n such that $\chi = 1$ on a neighborhood of K and $\text{supp } \chi \subset U$. Let ρ be as in part (a). The function $v = \chi u + C\rho$ then satisfies condition (b) if the constant $C > 0$ is chosen big enough. Indeed, the (very) positive Hessian of $C\rho$ compensates the bounded negative part of the Hessian of χu on the compact support of $d\chi$ which is contained in $U \setminus K$. \square

2.3. Domains with smooth p -convex boundaries.

Proof of Theorem 1.2. As pointed out in the Introduction, these results were proved by Harvey and Lawson [34] for bounded domains; here we extend their arguments to unbounded domains.

Thus, let $D \subset \mathbb{R}^n$ be a domain with boundary bD of class \mathcal{C}^2 . Assume first that condition (a) holds; i.e., D is p -convex. It is immediate that such D is also locally p -convex, in the sense that every point $\mathbf{x} \in bD$ has a neighborhood $U \subset \mathbb{R}^n$ such that $D \cap U$ is p -convex (cf. [34, (3.1) and Theorem 3.7]; the cited results also give the converse implication for bounded domains). Furthermore, local p -convexity admits the following differential theoretic characterization (cf. [34, Remark 3.11]):

A smoothly bounded domain $D \subset \mathbb{R}^n$ is locally p -convex at $\mathbf{x} \in bD$ if and only if there are a neighborhood $U \subset \mathbb{R}^n$ of \mathbf{x} and a local smooth defining function ρ for D (i.e., $D \cap U = \{\rho < 0\}$ and $d\rho \neq 0$ on $bD \cap U = \{\rho = 0\}$) such that

$$\text{tr}_L \text{Hess}_\rho(\mathbf{y}) \geq 0 \quad \text{for every tangent } p\text{-plane } L \subset T_{\mathbf{y}} bD, \mathbf{y} \in bD \cap U.$$

This property is independent of the choice of ρ and is equivalent to property (c) in Theorem 1.2 (that the sum of p smallest principal curvatures of bD is nonnegative). Furthermore, setting $\delta = \text{dist}(\cdot, bD)$, D is locally p -convex if and only if the function $-\log \delta$ is p -plurisubharmonic on a collar around bD in D (cf. [34, Summary 3.16]).

This justifies the implications (a) \Rightarrow (b) \Leftrightarrow (c) \Rightarrow (d) in Theorem 1.2.

It remains to prove that (d) \Rightarrow (a). Assume that (d) holds; i.e., the \mathcal{C}^2 function $-\log \delta$ is p -plurisubharmonic on an interior collar $U \subset D$ around bD . Choose a smooth cut-off function $\chi: \mathbb{R}^n \rightarrow [0, 1]$ which equals 0 on an open set $V \subset D$ containing $D \setminus U$ and equals 1 on an open set $W \subset \mathbb{R}^n$ containing $\mathbb{R}^n \setminus D$. Its differential $d\chi$ has support in the set $U \setminus W$ whose closure is contained in D . The product $-\chi \log \delta$ is then a function of class $\mathcal{C}^2(D)$ which is p -plurisubharmonic near bD and tends to $+\infty$ along bD . Let $h: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a smooth, increasing, strongly convex function. If h is chosen such that its derivative $h'(t) > 0$ grows sufficiently fast as $t \rightarrow +\infty$, then we see from (2.4) that the function

$$\rho(\mathbf{x}) = -\chi(\mathbf{x}) \log \delta(\mathbf{x}) + h(\|\mathbf{x}\|^2), \quad \mathbf{x} \in D,$$

is a strongly p -plurisubharmonic exhaustion function on D , so condition (a) holds. \square

Corollary 2.8. *A domain $D \subset \mathbb{R}^n$ (not necessarily bounded) whose boundary bD is a smooth embedded minimal hypersurface is $(n-1)$ -convex (also called mean-convex; see Remark 1.6). In particular, a domain in \mathbb{R}^3 bounded by a closed embedded minimal surface is minimally convex.*

2.4. The maximum principle and the Kontinuitätssatz. Since the restriction of a p -plurisubharmonic function u on a domain $D \subset \mathbb{R}^n$ to a minimal p -dimensional submanifold $M \subset D$ is subharmonic on M (cf. Propositions 2.2 and 2.4), it follows from the maximum principle for subharmonic functions that, for any compact

minimal p -dimensional submanifold $M \subset D$ with boundary bM , we have the implication

$$bM \subset K \implies M \subset \widehat{K}_{p,D}.$$

The same conclusion holds for immersed minimal submanifolds and for minimal p -dimensional currents. Furthermore, we have the following result, which is analogous to the classical Kontinuitätssatz (also called the *continuity principle*) in complex analysis. (Compare with Harvey and Lawson [34], proof of Theorem 3.9 on p. 159.)

Proposition 2.9 (Kontinuitätssatz for minimal submanifolds). *Assume that D is a p -convex domain in \mathbb{R}^n for some $p \in \{1, \dots, n\}$ and $\{M_t\}_{t \in [0,1]}$ is a continuous family of immersed compact minimal p -dimensional submanifolds of \mathbb{R}^n with boundaries bM_t . If $M_0 \subset D$ and $\bigcup_{t \in [0,1]} bM_t$ is contained in a compact subset of D , then $\bigcup_{t \in [0,1]} M_t$ is also contained in a compact subset of D .*

Proof. Let K denote the closure of the set $M_0 \cup \bigcup_{t \in [0,1]} bM_t$ in D . By the hypothesis, K is compact. Since D is p -convex, the p -hull $L = \widehat{K}_{p,D} \subset D$ of K is also compact. Consider the set $J = \{t \in [0,1] : M_t \subset L\}$. We have $0 \in J$ by the hypothesis. We claim that $J = [0,1)$. Since the family M_t is continuous in t and L is compact, J is closed. It remains to see that J is also open. Assume that $t_0 \in J$; then $M_{t_0} \subset L \subset D$. By continuity, it follows that $M_t \subset D$ for all $t \in [0,1)$ sufficiently close to t_0 , and the maximum principle implies that $M_t \subset L$ for all such t . \square

Problem 2.10. Assume that $1 < p < n$ and D is a domain in \mathbb{R}^n which satisfies the conclusion of Proposition 2.9 for minimal p -dimensional submanifolds. Does it follow that D is p -convex? Is the function $-\log \text{dist}(\cdot, bD)$ p -plurisubharmonic on D ?

If bD is smooth, then the validity of the Kontinuitätssatz for D implies (by Harvey and Lawson; cf. Theorem 1.2 above) that $-\log \text{dist}(\cdot, bD)$ is p -plurisubharmonic near bD ; even in this case, it is not clear whether it is p -plurisubharmonic on all of D . The analogous result in complex analysis is Oka’s theorem, saying that the function $-\log \text{dist}(\cdot, bD)$ is plurisubharmonic on any Hartogs pseudoconvex domain $D \subset \mathbb{C}^n$ (see e.g. [51, Theorem 5.6, p. 96]). Its proof breaks down in the present situation since the sum of two minimal discs in \mathbb{R}^n is not a minimal disc in general.

The following result will be used in the proof of Theorem 1.16 in Section 4.

Proposition 2.11 (The maximum principle for minimal submanifolds). *Let D be a proper p -convex domain in \mathbb{R}^n and let $M \subset D$ be a compact, connected, immersed minimal p -dimensional submanifold with boundary bM . Then the following hold:*

- (a) $\text{dist}(bM, bD) = \text{dist}(M, bD)$.
- (b) *If D has smooth boundary and $\text{dist}(\mathbf{x}_0, bD) = \text{dist}(bM, bD)$ for some point $\mathbf{x}_0 \in \overset{\circ}{M} = M \setminus bM$, then bD contains a translate of M .*
- (c) *If the assumption in part (b) holds for $p = 2, n = 3$ (i.e., M is a compact minimal surface with boundary in a minimally convex domain $D \subset \mathbb{R}^3$ and $\text{dist}(\mathbf{x}_0, bD) = \text{dist}(bM, bD)$ for some $\mathbf{x}_0 \in \overset{\circ}{M}$), then M is a piece of a plane. Moreover, if $\mathbf{y}_0 \in bD$ is such that $\|\mathbf{x}_0 - \mathbf{y}_0\| = \text{dist}(bM, bD)$, then $\bigcup_{t \in [0,1]} t(\mathbf{y}_0 - \mathbf{x}_0) + M \subset D$ and $(\mathbf{y}_0 - \mathbf{x}_0) + M \subset bD$.*

Proof of (a). Assume that $\text{dist}(\mathbf{x}_0, bD) < \text{dist}(bM, bD)$ for some $\mathbf{x}_0 \in \overset{\circ}{M}$. Pick a point $\mathbf{y}_0 \in bD$ such that $\text{dist}(\mathbf{x}_0, bD) = \|\mathbf{x}_0 - \mathbf{y}_0\|$ and a number t_0 with

$\|\mathbf{x}_0 - \mathbf{y}_0\| < t_0 < \text{dist}(bM, bD)$. The family of translates $M_t = M + t(\mathbf{y}_0 - \mathbf{x}_0)/\|\mathbf{y}_0 - \mathbf{x}_0\|$ for $t \in [0, t_0]$ then violates Proposition 2.9. This contradiction proves part (a).

Proof of (b). By Theorem 1.2, there are a neighborhood $U \subset \mathbb{R}^n$ of bD and a p -plurisubharmonic function ρ on U such that $U \cap D = \{\mathbf{x} \in U : \rho(\mathbf{x}) < 0\}$. Let $\mathbf{x}_0 \in \overset{\circ}{M}$ be such that $c = \text{dist}(\mathbf{x}_0, bD) = \text{dist}(M, bD)$. Pick a point $\mathbf{y}_0 \in bD$ with $\|\mathbf{x}_0 - \mathbf{y}_0\| = c$. There is a compact connected neighborhood $V \subset M$ of \mathbf{x}_0 in M such that the translate $W = V + \mathbf{y}_0 - \mathbf{x}_0$ is contained in U , and hence in $U \cap \overline{D}$ by part (a). Clearly $\mathbf{y}_0 \in W$. Since the function $\rho|_W \leq 0$ is subharmonic and $\rho(\mathbf{y}_0) = 0$, it is constantly equal to zero by the maximum principle, and hence $W \subset bD$. This means that for every $\mathbf{x} \in V$, we have

$$(2.5) \quad \mathbf{x} + \mathbf{y}_0 - \mathbf{x}_0 \in bD \quad \text{and} \quad \text{dist}(\mathbf{x}, bD) = \text{dist}(M, bD).$$

This argument shows that the set of points $\mathbf{x} \in M$ satisfying (2.5) is open, and clearly it is also closed, so it equals M . Thus $M + \mathbf{y}_0 - \mathbf{x}_0 \subset bD$.

Proof of (c). Let \mathbf{x}_0 and \mathbf{y}_0 be as in the statement of (c). Then M does not intersect the open ball centered at \mathbf{y}_0 of radius $\|\mathbf{x}_0 - \mathbf{y}_0\| = \text{dist}(bM, bD)$. This implies that $\mathbf{y}_0 = \mathbf{x}_0 + cN(\mathbf{x}_0)$, where $N: V \rightarrow \mathbb{S}^2$ is a Gauss map of the orientable surface $V \subset M$ introduced in part (b). Since (2.5) holds for all $\mathbf{x} \in V$, we see that $N(\mathbf{x}) = N(\mathbf{x}_0)$ for all $\mathbf{x} \in V$. This shows that V , and hence also M , is a piece of a plane and (c) follows. \square

2.5. Null plurisubharmonic functions. Let $\mathbf{z} = (z_1, \dots, z_n) = \mathbf{x} + i\mathbf{y}$ be complex coordinates on \mathbb{C}^n , with $z_j = x_j + iy_j$ for $j = 1, \dots, n$. We shall write $\mathbf{0} = (0, \dots, 0)$ for the origin in \mathbb{R}^n or in \mathbb{C}^n . Given a \mathcal{C}^2 function $\rho: \Omega \rightarrow \mathbb{R}$ on a domain $\Omega \subset \mathbb{C}^n$, we denote by $\mathcal{L}_\rho(\mathbf{z}; \cdot)$ its *Levi form* at a point $\mathbf{z} \in \Omega$:

$$(2.6) \quad \mathcal{L}_\rho(\mathbf{z}; \mathbf{w}) = \sum_{j,k=1}^n \frac{\partial^2 \rho}{\partial z_j \partial \bar{z}_k}(\mathbf{z}) w_j \bar{w}_k, \quad \mathbf{w} = (w_1, \dots, w_n) \in \mathbb{C}^n.$$

We shall use the following lemma, whose proof amounts to a simple calculation.

Lemma 2.12. *Let $B = (b_{j,k})$ be a real symmetric $n \times n$ matrix, and let $\mathbf{w} = \mathbf{u} + i\mathbf{v} \in \mathbb{C}^n$. Then*

$$2 \sum_{j,k=1}^n b_{j,k} u_j u_k = \Re \left(\sum_{j,k=1}^n b_{j,k} w_j w_k \right) + \sum_{j,k=1}^n b_{j,k} w_j \bar{w}_k.$$

A function $\rho: D \rightarrow \mathbb{R}$ on a domain $D \subset \mathbb{R}^n$ will also be considered as a function on the tube $\mathcal{T}_D = D \times i\mathbb{R}^n \subset \mathbb{C}^n$ which is independent of the imaginary variable:

$$(2.7) \quad \rho(\mathbf{x} + i\mathbf{y}) = \rho(\mathbf{x}) \quad \text{for all } \mathbf{x} \in D \text{ and } \mathbf{y} \in \mathbb{R}^n.$$

Fix a point $\mathbf{x} \in D$ and a vector $\mathbf{u} \in \mathbb{R}^n$. The Hessian $\text{Hess}_\rho(\mathbf{x}; \cdot)$ (2.1) has coefficients

$$b_{j,k} := \frac{\partial^2 \rho}{\partial x_j \partial x_k}(\mathbf{x}) = 4 \frac{\partial^2 \rho}{\partial z_j \partial \bar{z}_k}(\mathbf{x}) \in \mathbb{R}.$$

Lemma 2.12 shows that for every $\mathbf{w} = \mathbf{u} + i\mathbf{v} \in \mathbb{C}^n$, we have

$$(2.8) \quad \frac{1}{2} \text{Hess}_\rho(\mathbf{x}; \mathbf{u}) = \frac{1}{4} \Re \left(\sum_{j,k=1}^n b_{j,k} w_j w_k \right) + \mathcal{L}_\rho(\mathbf{x}; \mathbf{w}).$$

Replacing \mathbf{w} by $-\mathbf{i}\mathbf{w} = \mathbf{v} - \mathbf{u}$ and noting that $\mathcal{L}_\rho(\mathbf{x}; \pm\mathbf{i}\mathbf{w}) = \mathcal{L}_\rho(\mathbf{x}; \mathbf{w})$ while the first term on the right hand side of (2.8) changes sign, we obtain

$$\text{Hess}_\rho(\mathbf{x}; \mathbf{u}) + \text{Hess}_\rho(\mathbf{x}; \mathbf{v}) = 4\mathcal{L}_\rho(\mathbf{x}; \mathbf{u} + \mathbf{i}\mathbf{v}).$$

In particular, if (\mathbf{u}, \mathbf{v}) is an orthonormal pair of vectors in \mathbb{R}^n and we set

$$L = \mathbf{x} + \text{span}_{\mathbb{R}}\{\mathbf{u}, \mathbf{v}\} \subset \mathbb{R}^n, \quad \Lambda = \mathbf{x} + \text{span}_{\mathbb{C}}\{\mathbf{u} + \mathbf{i}\mathbf{v}\} \subset \mathbb{C}^n,$$

then it follows that

$$(2.9) \quad \Delta(\rho|_L)(\mathbf{x}) = 4\mathcal{L}_\rho(\mathbf{x}; \mathbf{u} + \mathbf{i}\mathbf{v}) = \Delta(\rho|_\Lambda)(\mathbf{x}).$$

Set $a_j = \frac{\partial \rho}{\partial x_j}(\mathbf{x}) \in \mathbb{R}$ for $j = 1, \dots, n$. The identity (2.8) implies that for every point $\mathbf{z} = \mathbf{x} + \mathbf{i}\mathbf{y} \in \mathcal{T}_D$ and vector $\mathbf{w} = \mathbf{u} + \mathbf{i}\mathbf{v} \in \mathbb{C}^n$ near $\mathbf{0} \in \mathbb{C}^n$, we have the Taylor expansion

$$\begin{aligned} \rho(\mathbf{z} + \mathbf{w}) &= \rho(\mathbf{x}) + \sum_{j=1}^n a_j u_j + \frac{1}{2} \text{Hess}_\rho(\mathbf{x}; \mathbf{u}) + o(\|\mathbf{u}\|^2) \\ &= \rho(\mathbf{x}) + \Re \left(\sum_{j=1}^n a_j w_j + \frac{1}{4} \sum_{j,k=1}^n b_{j,k} w_j w_k \right) + \mathcal{L}_\rho(\mathbf{x}; \mathbf{w}) + o(\|\mathbf{w}\|^2). \end{aligned}$$

Denote by $\Sigma_{\mathbf{x}} \subset \mathbb{C}^n$ the local complex hypersurface near the origin $\mathbf{0} \in \mathbb{C}^n$ given by

$$(2.10) \quad \Sigma_{\mathbf{x}} = \left\{ \mathbf{w} : \sum_{j=1}^n a_j w_j + \frac{1}{4} \sum_{j,k=1}^n b_{j,k} w_j w_k = 0 \right\}.$$

It follows that

$$(2.11) \quad \rho(\mathbf{z} + \mathbf{w}) = \rho(\mathbf{z}) + \mathcal{L}_\rho(\mathbf{x}; \mathbf{w}) + o(\|\mathbf{w}\|^2), \quad \mathbf{z} = \mathbf{x} + \mathbf{i}\mathbf{y} \in \mathcal{T}_D, \mathbf{w} \in \Sigma_{\mathbf{x}}.$$

We need to recall the connection between minimal plurisubharmonic functions on a domain $D \subset \mathbb{R}^n$ and null plurisubharmonic functions on the tube $\mathcal{T}_D = D \times \mathbf{i}\mathbb{R}^n \subset \mathbb{C}^n$; the latter class of functions was introduced in [21].

Let $\mathfrak{A} \subset \mathbb{C}^n$ denote the *null quadric*:

$$(2.12) \quad \mathfrak{A} = \{ \mathbf{z} = (z_1, \dots, z_n) \in \mathbb{C}^n : z_1^2 + z_2^2 + \dots + z_n^2 = 0 \}, \quad \mathfrak{A}_* = \mathfrak{A} \setminus \{ \mathbf{0} \}.$$

Definition 2.13 (Definitions 2.1 and 2.4 in [21]). Let Ω be a domain in \mathbb{C}^n for some $n \geq 3$.

- (a) An upper semicontinuous function $u : \Omega \rightarrow \mathbb{R} \cup \{-\infty\}$ is *null plurisubharmonic* ($u \in \mathfrak{NPsh}(\Omega)$) if, for any affine complex line $L \subset \mathbb{C}^n$ directed by a null vector $\theta \in \mathfrak{A}_*$, the restriction of u to $L \cap \Omega$ is subharmonic. (If $u \in \mathcal{C}^2(\Omega)$, this is equivalent to the condition that $\mathcal{L}_u(\mathbf{z}; \mathbf{w}) \geq 0$ for every $\mathbf{z} \in \Omega$ and $\mathbf{w} \in \mathfrak{A}_*$.)
- (b) A function $u \in \mathcal{C}^2(\Omega)$ is *null strongly plurisubharmonic* if $\mathcal{L}_u(\mathbf{z}; \mathbf{w}) > 0$ for every $\mathbf{z} \in \Omega$ and $\mathbf{w} \in \mathfrak{A}_*$.

Note that a vector $0 \neq \mathbf{w} = \mathbf{u} + \mathbf{i}\mathbf{v} \in \mathbb{C}^n$ belongs to the null quadric \mathfrak{A} (2.12) if and only if the vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ are orthogonal and have equal length:

$$(2.13) \quad \mathbf{u} + \mathbf{i}\mathbf{v} \in \mathfrak{A}_* \iff \mathbf{u} \cdot \mathbf{v} = 0 \text{ and } \|\mathbf{u}\| = \|\mathbf{v}\|.$$

Assume that (\mathbf{u}, \mathbf{v}) is an orthonormal pair in \mathbb{R}^n . In view of (2.9), we have the following result for functions $u \in \mathcal{C}^2(D)$; the general case for upper semicontinuous functions is seen similarly.

Lemma 2.14 (Lemma 4.3 in [21]). *Let D be a domain in \mathbb{R}^n and let $\mathcal{T}_D = D \times i\mathbb{R}^n \subset \mathbb{C}^n$.*

- *If u is (strongly) minimal plurisubharmonic on D , then the function $U(\mathbf{x} + i\mathbf{y}) = u(\mathbf{x})$ is (strongly) null plurisubharmonic on \mathcal{T}_D .*
- *Conversely, assume that a function $U: \mathcal{T}_D \rightarrow \mathbb{R}$ is independent of the variable $\mathbf{y} = \Im \mathbf{z}$, and let $u(\mathbf{x}) = U(\mathbf{x} + i\mathbf{0})$ for $\mathbf{x} \in D$. If U is (strongly) null plurisubharmonic on \mathcal{T}_D , then u is (strongly) minimal plurisubharmonic on D .*

Recall that a *null holomorphic disc* in \mathbb{C}^n ($n \geq 3$) is a holomorphic map $F = (F_1, \dots, F_n): \mathbb{D} \rightarrow \mathbb{C}^n$ satisfying the nullity condition $F'(\zeta) \in \mathfrak{A}$; equivalently,

$$(2.14) \quad F'_1(\zeta)^2 + F'_2(\zeta)^2 + \dots + F'_n(\zeta)^2 = 0, \quad \zeta \in \mathbb{D}.$$

More generally, a holomorphic immersion $F: M \rightarrow \mathbb{C}^n$ from an open Riemann surface M is a *holomorphic null curve* if the derivative of F in any local holomorphic coordinate on M satisfies the condition (2.14). It follows from (2.13) and the Cauchy-Riemann equations that the real and the imaginary parts of a holomorphic null disc $F: \mathbb{D} \rightarrow \mathbb{C}^n$ are conformal minimal discs in \mathbb{R}^n ; conversely, every conformal minimal disc is the real part of a holomorphic null disc. We have the following observation.

Proposition 2.15 (Proposition 2.7 in [21]). *An upper semicontinuous function u on a domain $\Omega \subset \mathbb{C}^n$ ($n \geq 3$) is null plurisubharmonic if and only if the function $u \circ F$ is subharmonic on \mathbb{D} for every null holomorphic disc $F: \mathbb{D} \rightarrow \Omega$.*

3. PROOF OF THEOREMS 1.1, 1.7, AND 1.9

We begin with technical preparations.

Let $\rho: D \rightarrow \mathbb{R}$ be a smooth minimal strongly plurisubharmonic exhaustion function on a domain $D \subset \mathbb{R}^3$. We extend ρ to a function on the tube $\mathcal{T}_D = D \times i\mathbb{R}^3 \subset \mathbb{C}^3$ which is independent of the imaginary variable; see (2.7). By Lemma 2.14, the extended function ρ is null strongly plurisubharmonic on \mathcal{T}_D . For every point $\mathbf{x} \in D$ we denote by $\Sigma_{\mathbf{x}} \subset \mathbb{C}^3$ the local complex hypersurface at $\mathbf{0} \in \mathbb{C}^3$ given by (2.10):

$$(3.1) \quad \Sigma_{\mathbf{x}} = \left\{ \mathbf{w} = (w_1, w_2, w_3) : \sum_{j=1}^3 \frac{\partial \rho}{\partial x_j}(\mathbf{x}) w_j + \sum_{j,k=1}^3 \frac{\partial^2 \rho}{\partial z_j \partial \bar{z}_k}(\mathbf{x}) w_j w_k = 0 \right\}.$$

Let P denote the critical locus of ρ . We assume in the sequel that $\mathbf{x} \in D \setminus P$; then $\Sigma_{\mathbf{x}}$ is nonsingular at $\mathbf{0} \in \Sigma_{\mathbf{x}}$ and its tangent space is

$$(3.2) \quad T_{\mathbf{0}} \Sigma_{\mathbf{x}} = \left\{ \mathbf{w} \in \mathbb{C}^3 : \sum_{j=1}^3 \frac{\partial \rho}{\partial x_j}(\mathbf{x}) w_j = 0 \right\}.$$

Note that the coefficients $a_j = \frac{\partial \rho}{\partial x_j}(\mathbf{x})$ of the equation in (3.2) are real. By shrinking $\Sigma_{\mathbf{x}}$ around $\mathbf{0}$ if necessary, we may assume that the hypersurface $\Sigma_{\mathbf{x}}$ is nonsingular.

The intersection of the null quadric \mathfrak{A} (2.12) with any complex 2-plane $\Lambda \subset \mathbb{C}^3$ consists of two complex lines which may coincide for certain Λ . However, for a 2-plane $\Lambda = \{ \mathbf{w} = (w_1, w_2, w_3) \in \mathbb{C}^3 : \sum_{j=1}^3 a_j w_j = 0 \}$ with real coefficients $a_1, a_2, a_3 \in \mathbb{R}$ not all equal to 0, the intersection $\mathfrak{A} \cap \Lambda$ consists of two distinct complex lines as is seen by a simple calculation. Identifying the tangent space $T_{\mathbf{z}} \mathbb{C}^3$ with \mathbb{C}^3 , we may consider the null quadric \mathfrak{A} as a subset of $T_{\mathbf{z}} \mathbb{C}^3$ for any

point $\mathbf{z} \in \mathbb{C}^3$. By what has been said above, for any point $\mathbf{z} \in \Sigma_{\mathbf{x}}$ sufficiently close to $\mathbf{0}$ the intersection $\mathfrak{A} \cap T_{\mathbf{z}}\Sigma_{\mathbf{x}}$ is a union of two distinct complex lines. This defines on $\Sigma_{\mathbf{x}}$ a couple of holomorphic direction fields, and hence (by integration) a couple of 1-dimensional complex analytic foliations by holomorphic null curves. In particular, for any point $\mathbf{x} \in D \setminus P$ we have two distinct embedded holomorphic null discs $\mathcal{N}_{\mathbf{x}}^1, \mathcal{N}_{\mathbf{x}}^2 \subset \Sigma_{\mathbf{x}}$ passing through $\mathbf{0}$. Although there is no well defined global ordering of these two null discs when \mathbf{x} runs over $D \setminus P$, such an ordering clearly exists on every simply connected subset. By the definition of $\Sigma_{\mathbf{x}}$ and (2.11), we have that

$$\rho(\mathbf{z} + \mathbf{w}) = \rho(\mathbf{z}) + \mathcal{L}_{\rho}(\mathbf{x}; \mathbf{w}) + o(\|\mathbf{w}\|^2), \quad \mathbf{w} \in \Sigma_{\mathbf{x}}.$$

This holds in particular for all $\mathbf{w} \in \mathcal{N}_{\mathbf{x}}^1 \cup \mathcal{N}_{\mathbf{x}}^2 \subset \Sigma_{\mathbf{x}}$. Since ρ is null strongly plurisubharmonic on \mathcal{T}_D , the Levi form $\mathcal{L}_{\rho}(\mathbf{x}; \cdot)$ is positive on the null lines $T_{\mathbf{0}}\mathcal{N}_{\mathbf{x}}^j$ for $j = 1, 2$. It follows that for every point $\mathbf{z} = \mathbf{x} + i\mathbf{y} \in \mathcal{T}_D$ with $\mathbf{x} \in D \setminus P$ there exist constants $C_{\mathbf{x}} > 0$ and $\delta_{\mathbf{x}} > 0$ such that

$$(3.3) \quad \rho(\mathbf{z} + \mathbf{w}) \geq \rho(\mathbf{z}) + C_{\mathbf{x}}\|\mathbf{w}\|^2, \quad \mathbf{w} \in \mathcal{N}_{\mathbf{x}}^1 \cup \mathcal{N}_{\mathbf{x}}^2, \quad \|\mathbf{w}\| \leq \delta_{\mathbf{x}}.$$

Moreover, the constants $C_{\mathbf{x}}$ and $\delta_{\mathbf{x}}$ can clearly be chosen uniform for all points \mathbf{x} in any given compact subset of $D \setminus P$. By projecting the discs $\mathcal{N}_{\mathbf{x}}^1, \mathcal{N}_{\mathbf{x}}^2$ to \mathbb{R}^3 we get a corresponding family of conformal minimal discs with the analogous properties.

We summarize the above discussion in the following lemma.

Lemma 3.1. *Let D be a domain in \mathbb{R}^3 , and let $\rho: D \rightarrow \mathbb{R}$ be a \mathcal{C}^2 minimal strongly plurisubharmonic function with the critical locus P . For every compact set $L \subset D \setminus P$ there exist a constant $c = c_L > 0$ and families of embedded null holomorphic discs $\sigma_{\mathbf{x}}^j = \alpha_{\mathbf{x}}^j + i\beta_{\mathbf{x}}^j: \overline{\mathbb{D}} \rightarrow \mathbb{C}^3$ ($\mathbf{x} \in L, j = 1, 2$), depending locally \mathcal{C}^1 smoothly on the point $\mathbf{x} \in L$ and satisfying the following conditions:*

- (a) $\sigma_{\mathbf{x}}^j(0) = 0$;
- (b) $\{\mathbf{x} + \alpha_{\mathbf{x}}^j(\zeta) : \zeta \in \overline{\mathbb{D}}\} \subset D$;
- (c) *the function $\overline{\mathbb{D}} \ni \zeta \mapsto \rho(\mathbf{x} + \alpha_{\mathbf{x}}^j(\zeta))$ is strongly convex and satisfies*

$$(3.4) \quad \rho(\mathbf{x} + \alpha_{\mathbf{x}}^j(\zeta)) \geq \rho(\mathbf{x}) + c\|\zeta\|^2, \quad \zeta \in \overline{\mathbb{D}}.$$

The conformal minimal discs $\alpha_{\mathbf{x}}^j: \overline{\mathbb{D}} \rightarrow \mathbb{R}^3$, furnished by Lemma 3.1, will be used to push the boundary $F(bM)$ of a given conformal minimal immersion $F: M \rightarrow D$ to a higher level set of ρ , except near the critical points of ρ , which shall be avoided by a different method explained in what follows. The relevant tool for this lifting is the following. (Related results on the Riemann-Hilbert problem for null curves are given by [3, Theorem 4] in dimension $n = 3$ and by [2, Theorem 3.5] in arbitrary dimension $n \geq 3$.)

Theorem 3.2 (Riemann-Hilbert problem for conformal minimal surfaces in \mathbb{R}^3). *Let M be a compact bordered Riemann surface with nonempty boundary $bM \neq \emptyset$, let I_1, \dots, I_k be pairwise disjoint compact subarcs of bM which are not connected components of bM , and set $I = \bigcup_{j=1}^k I_j$. Choose a thin annular neighborhood $A \subset M$ of bM and a smooth retraction $\rho: A \rightarrow bM$. Assume that*

- $F: M \rightarrow \mathbb{R}^3$ is a conformal minimal immersion of class $\mathcal{C}^1(M)$,
- $r: bM \rightarrow [0, 1]$ is a continuous function supported on I , and
- $\alpha: I \times \overline{\mathbb{D}} \rightarrow \mathbb{R}^3$ is a map of class \mathcal{C}^1 such that for every $\zeta \in I$ the map $\overline{\mathbb{D}} \ni \xi \mapsto \alpha(\zeta, \xi) \in \mathbb{R}^3$ is a conformal minimal immersion with $\alpha(\zeta, 0) = 0$.

Let the map $\varkappa: bM \times \overline{\mathbb{D}} \rightarrow \mathbb{R}^3$ be given by

$$(3.5) \quad \varkappa(\zeta, \xi) = F(\zeta) + \alpha(\zeta, r(\zeta)\xi),$$

where we take $\alpha(\zeta, r(\zeta)\xi) = 0$ for $\zeta \in bM \setminus I$. Given a number $\eta > 0$ and an open neighborhood $\Omega \subset M$ of I , there exists a conformal minimal immersion $G: M \rightarrow \mathbb{R}^3$ of class $\mathcal{C}^1(M)$ satisfying the following conditions:

- (i) $\text{dist}(G(\zeta), \varkappa(\zeta, \mathbb{T})) < \eta$ for all $\zeta \in bM$;
- (ii) $\text{dist}(G(\zeta), \varkappa(\rho(\zeta), \overline{\mathbb{D}})) < \eta$ for all $\zeta \in \Omega$;
- (iii) $\|G - F\|_{1, M \setminus \Omega} < \eta$;
- (iv) $\text{Flux}(G) = \text{Flux}(F)$.

Proof. If M is the disc $\overline{\mathbb{D}}$, the conclusion follows from [2, Lemma 3.1], which gives an analogous result for null holomorphic immersions in \mathbb{C}^3 . Since every conformal minimal disc $\overline{\mathbb{D}} \rightarrow \mathbb{R}^3$ is the real part of a holomorphic null disc $\overline{\mathbb{D}} \rightarrow \mathbb{C}^3$, the cited lemma can be used for the corresponding families of null discs; the real part G of the resulting null disc then satisfies the conclusion of Theorem 3.2. (The loss of smoothness in harmonic conjugates is not important since we can restrict our maps to a slightly smaller disc.)

In the general case, for an arbitrary bordered Riemann surface M , one follows the proof of [2, Theorems 3.5 and 3.6], but replacing [2, Lemma 3.3] by [2, Lemma 3.1]. The former one holds in any dimension $n \geq 3$ but only applies to flat conformal minimal discs $\alpha(\zeta, \cdot): \overline{\mathbb{D}} \rightarrow \mathbb{R}^n$ lying in parallel 2-planes, while the latter one holds without any such restriction on α but only in dimension $n = 3$. □

The next result is the main technical ingredient in the proofs of Theorems 1.1, 1.7, and 1.9. Similar techniques have been used for lifting boundaries of complex curves and Stein varieties in q -convex manifolds; see e.g. [18, 19] and the references therein.

Proposition 3.3 (Lifting boundaries of conformal minimal surfaces). *Let D be a domain in \mathbb{R}^3 and let $\rho: D \rightarrow \mathbb{R}$ be a \mathcal{C}^2 minimal strongly plurisubharmonic function with the critical locus P . Given a compact set $L \subset D \setminus P$, there exist constants $\epsilon_0 > 0$ and $C_0 > 0$ such that the following holds.*

Let M be a compact bordered Riemann surface, and let $F: M \rightarrow D$ be a conformal minimal immersion of class $\mathcal{C}^1(M)$. Given a continuous function $\epsilon: bM \rightarrow [0, \epsilon_0]$ supported on the set $J = \{\zeta \in bM : F(\zeta) \in L\}$, an open set $U \subset M$ containing $\text{supp}(\epsilon)$ in its relative interior, and a constant $\delta > 0$, there exists a conformal minimal immersion $G: M \rightarrow D$ satisfying the following conditions:

- (1) $|\rho(G(\zeta)) - \rho(F(\zeta)) - \epsilon(\zeta)| < \delta$ for every $\zeta \in bM$;
- (2) $\rho(G(\zeta)) \geq \rho(F(\zeta)) - \delta$ for every $\zeta \in M$;
- (3) $\|G - F\|_{1, M \setminus U} < \delta$;
- (4) $\|G - F\|_{0, M} \leq C_0\sqrt{\epsilon_0}$;
- (5) $\text{Flux}(G) = \text{Flux}(F)$.

Proof. By approximation, we may assume that F is of class $\mathcal{C}^\infty(M)$ (see [5, 6]).

Pick a compact set $L_0 \subset D \setminus P$ which contains L in its interior. Let c_{L_0} be the constant furnished by Lemma 3.1 for the set L_0 , and choose a number ϵ_0 such that $0 < \epsilon_0 < c_{L_0}$. Set $J_0 = \{\zeta \in bM : F(\zeta) \in L_0\}$. By approximation, we may assume that the function $\epsilon: bM \rightarrow [0, \epsilon_0]$ in Proposition 3.3 is smooth and supported in the relative interior of $J_0 \cap U$.

Assume first that the support of ϵ does not contain any boundary curves of M ; the general case will be obtained by two consecutive applications of this special case. Choose finitely many closed pairwise disjoint segments $I_1, I_2, \dots, I_m \subset J_0 \cap U$ whose union $I = \bigcup_{j=1}^m I_j$ contains $\text{supp}(\epsilon)$ in its relative interior. Note that $F(I) \subset L_0$. Since I is simply connected, Lemma 3.1 (see in particular (3.4)) furnishes a family of conformal minimal discs $\alpha_{F(\zeta)}: \mathbb{D} \rightarrow \mathbb{R}^3$, depending smoothly on $\zeta \in I$, such that

$$(3.6) \quad \rho(F(\zeta) + \alpha_{F(\zeta)}(\xi)) \geq \rho(F(\zeta)) + c_{L_0} > \rho(F(\zeta)) + \epsilon_0, \quad \zeta \in I, |\xi| = 1.$$

Without loss of generality we may assume that $\delta < 3\epsilon_0$. Let $\tilde{\epsilon}: I \rightarrow [\delta/3, \epsilon_0]$ be obtained by smoothing the function $\max\{\epsilon, \delta/3\}$; in particular, we assume that $\tilde{\epsilon} = \epsilon$ on the set where $\epsilon \geq \delta/2$ and $\delta/3 \leq \tilde{\epsilon} < \delta/2$ on the complementary set. The properties of the discs $\alpha_{\mathbf{x}}$, furnished by Lemma 3.1, imply that for every fixed $\zeta \in I$ the function $\mathbb{D} \ni \xi \mapsto \rho(F(\zeta) + \alpha_{F(\zeta)}(\xi))$ is strongly convex, with a minimum at $\xi = 0$ and no other critical points. In view of (3.6) the set

$$(3.7) \quad \mathcal{D}_\zeta := \{\xi \in \mathbb{D} : \rho(F(\zeta) + \alpha_{F(\zeta)}(\xi)) < \rho(F(\zeta)) + \tilde{\epsilon}(\zeta)\}$$

contains the origin, is simply connected (a disc), and is compactly contained in \mathbb{D} ; furthermore, the discs \mathcal{D}_ζ depend smoothly on the point $\zeta \in I$. Choose a smooth family of diffeomorphisms $\phi_\zeta: \mathbb{D} \rightarrow \mathcal{D}_\zeta$ ($\zeta \in I$) which are holomorphic in \mathbb{D} and satisfy $\phi_\zeta(0) = 0$. Let $\alpha: I \times \mathbb{D} \rightarrow \mathbb{R}^3$ be defined by

$$(3.8) \quad \alpha(\zeta, \xi) = \alpha_{F(\zeta)}(\phi_\zeta(\xi)), \quad \zeta \in I, \xi \in \mathbb{D}.$$

Pick a smooth function $r: I \rightarrow [0, 1]$ such that $r(\zeta) = 1$ when $\epsilon(\zeta) \geq \delta/2$ and the support of r is contained in the relative interior of $J_0 \cap U$.

We now apply Theorem 3.2 to the conformal minimal immersion $F: M \rightarrow D$, the map α given by (3.8), and the function r . It is straightforward to verify that the resulting conformal minimal immersion $G: M \rightarrow D$ satisfies the conclusion of Proposition 3.3 provided that the number $\eta > 0$ in Theorem 3.2 is chosen small enough. The existence of a constant $C_0 > 0$ satisfying the estimate (4) in Proposition 3.3 is immediate from the geometry of the discs $\alpha_{\mathbf{x}}^j(\cdot)$ furnished by Lemma 3.1. Indeed, we clearly have a uniform estimate $\|\alpha_{\mathbf{x}}^j(\xi)\| \leq b|\xi|$ ($\xi \in \mathbb{D}, \mathbf{x} \in L, j = 1, 2$) for some constant $b > 0$. From (3.8), we get $\|\alpha(\zeta, \xi)\| \leq b|\phi_\zeta(\xi)|$ for $\zeta \in I$ and $\xi \in \mathbb{D}$. Together with (3.4), (3.7), and (3.8) one obtains

$$\epsilon_0 \geq \tilde{\epsilon}(\zeta) \geq \rho(F(\zeta) + \alpha(\zeta, \xi)) - \rho(F(\zeta)) \geq c|\phi_\zeta(\xi)|^2 \geq c/b^2 \|\alpha(\zeta, \xi)\|^2,$$

which gives $\|\alpha(\zeta, \xi)\| \leq C_0\sqrt{\epsilon_0}$ with $C_0 = b/\sqrt{c}$. By increasing C_0 slightly, this gives (4) provided that the approximation in Theorem 3.2 (see (3.5) and (i)) is close enough.

If the support of the function ϵ contains a boundary curve of M , then we write $\epsilon = \epsilon_1 + \epsilon_2$ where each of the two nonnegative functions $\epsilon_1, \epsilon_2: bM \rightarrow [0, \epsilon_0]$ satisfies the conditions of the special case considered above. By first deforming F to G_1 using the function ϵ_1 and subsequently deforming G_1 to $G = G_2$ using the function ϵ_2 , the resulting conformal minimal immersion G satisfies the conclusion of Proposition 3.3, provided that the approximations are sufficiently close at each step. \square

We now explain how to avoid critical points of a Morse exhaustion function $\rho: D \rightarrow \mathbb{R}$ when applying Proposition 3.3. To this end, we adapt the method from [26, Section 3.11].

Definition 3.4. A critical point \mathbf{x}_0 of a \mathcal{C}^2 function ρ is *nice* if, in some neighborhood of \mathbf{x}_0 , ρ agrees with its second order Taylor polynomial at \mathbf{x}_0 .

Lemma 3.5. *Every Morse function ρ can be approximated arbitrarily closely in the fine \mathcal{C}^2 topology by a Morse function $\tilde{\rho}$ with the same critical locus and with nice critical points. Furthermore, $\tilde{\rho}$ can be chosen to agree with ρ outside an arbitrarily small neighborhood of the critical locus.*

Proof. Assume that \mathbf{x}_0 is an (isolated) critical point of ρ and that

$$\rho(\mathbf{x}) = Q(\mathbf{x}) + \eta(\mathbf{x}), \quad \lim_{\mathbf{x} \rightarrow \mathbf{x}_0} \frac{\eta(\mathbf{x})}{\|\mathbf{x} - \mathbf{x}_0\|^2} = 0.$$

Choose a smooth increasing function $\chi: \mathbb{R} \rightarrow [0, 1]$ such that $\chi(t) = 0$ for $t \leq 1$ and $\chi(t) = 1$ for $t \geq 2$. Given $\epsilon > 0$, we consider the function

$$\rho_\epsilon(\mathbf{x}) = Q(\mathbf{x}) + \chi(\epsilon^{-1}\|\mathbf{x} - \mathbf{x}_0\|) \eta(\mathbf{x}).$$

Then $\rho_\epsilon = Q$ on the ball $\|\mathbf{x} - \mathbf{x}_0\| \leq \epsilon$ and $\rho_\epsilon = \rho$ on $\|\mathbf{x} - \mathbf{x}_0\| \geq 2\epsilon$. As $\epsilon \rightarrow 0$, the \mathcal{C}^2 norm of $\rho(\mathbf{x}) - \rho_\epsilon(\mathbf{x}) = (1 - \chi(\epsilon^{-1}\|\mathbf{x} - \mathbf{x}_0\|)) \eta(\mathbf{x})$ tends to zero. If $\epsilon > 0$ is chosen small enough, then ρ_ϵ satisfies the conclusion of the lemma at the critical point \mathbf{x}_0 . The same modification can be performed simultaneously at all critical points of ρ . □

A minimal strongly plurisubharmonic function has no critical points of index greater than 1 (see Remark 1.5). Critical points of index zero are local minima and are not approached by the boundary $F(bM)$ when applying Proposition 3.3.

Assume now that \mathbf{x}_0 is a nice Morse critical point of ρ with Morse index 1. The subsequent analysis is local near \mathbf{x}_0 , so we may assume, after a rigid motion of \mathbb{R}^3 , that $\mathbf{x}_0 = \mathbf{0} \in \mathbb{R}^3$, $\rho(\mathbf{x}_0) = 0$, and

$$(3.9) \quad \rho(\mathbf{x}) = \rho(x_1, x_2, x_3) = -a_1x_1^2 + a_2x_2^2 + a_3x_3^2 + \eta(\mathbf{x}),$$

where $-a_1 < 0 < a_2 \leq a_3$ and the function η vanishes in a neighborhood of the origin. Note that $a_1 < a_2$ since ρ is minimal strongly plurisubharmonic. Choose a number $c_0 > 0$ small enough such that η vanishes on the set

$$(3.10) \quad P_{c_0} := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : a_1x_1^2 \leq c_0, a_2x_2^2 + a_3x_3^2 \leq 4c_0\}.$$

The straight line arc $E \subset \mathbb{R}^3$, defined by

$$(3.11) \quad E = \{(x_1, 0, 0) \in \mathbb{R}^3 : a_1x_1^2 \leq c_0\},$$

is a local stable manifold of the critical point $\mathbf{0}$ of ρ . Set $\lambda = a_2/a_1 > 1$. Choose a number $\mu \in \mathbb{R}$ with $1 < \mu < \lambda$ and set

$$(3.12) \quad t_0 = c_0(1 - 1/\mu)^2;$$

hence $0 < t_0 < c_0(1 - 1/\lambda)^2 < c_0$.

The following is [26, Lemma 3.11.1, p. 98], adapted to the situation at hand.

Lemma 3.6 (Assumptions as above). *Assume that 0 is the only critical value of the function ρ (3.9) in the set $\{-c_0 < \rho < 3c_0\}$. Then there exists a minimal strongly plurisubharmonic function $\tau: D \cap \{\rho < 3c_0\} \rightarrow \mathbb{R}$ satisfying the following conditions:*

- (a) $\{\rho \leq -c_0\} \cup E \subset \{\tau \leq 0\} \subset \{\rho \leq -t_0\} \cup E$ (here, E is given by (3.11));
- (b) $\{\rho \leq c_0\} \subset \{\tau \leq 2c_0\} \subset \{\rho < 3c_0\}$;

- (c) *there is a constant $t_1 \in (t_0, c_0)$ such that $\tau = \rho + t_1$ outside the set P_{c_0} (3.10);*
- (d) *τ has no critical values in the interval $(0, 2c_0]$.*

The sublevel sets $\{\tau < c\}$ for $c > 0$ in a neighborhood of the origin are shown in [26, Figure 3.5, p. 100] (in a similar setting of strongly plurisubharmonic functions).

Proof. The choice of the number t_0 (3.12) implies that there is a smooth convex increasing function $h: \mathbb{R} \rightarrow [0, +\infty)$ satisfying the following conditions:

- (i) $h(t) = 0$ for $t \leq t_0$;
- (ii) for $t \geq c_0$ we have $h(t) = t - t_1$ with $t_1 = c_0 - h(c_0) \in (t_0, c_0)$;
- (iii) for $t_0 \leq t \leq c_0$ we have $t - t_1 \leq h(t) \leq t - t_0$;
- (iv) for all $t \in \mathbb{R}$ we have that $0 \leq \dot{h}(t) \leq 1$ and $2t\ddot{h}(t) + \dot{h}(t) < \lambda$.

The construction of such a function is entirely elementary (cf. [26, pp. 98-99]; its graph is shown in [26, Figure 3.4, p. 99]). Let $\tau: \mathbb{R}^3 \rightarrow \mathbb{R}$ be given by

$$(3.13) \quad \tau(\mathbf{x}) = -h(a_1x_1^2) + a_2x_2^2 + a_3x_3^2 + \eta(\mathbf{x}).$$

Setting $t = a_1x_1^2$, a calculation shows that on the set $P_{c_0} \subset \{\eta = 0\}$ (3.10) we have

$$-\frac{\partial^2\tau(\mathbf{x})}{\partial x_1^2} = 2a_1 \left(2t\ddot{h}(t) + \dot{h}(t) \right) < 2a_2 = \frac{\partial^2\tau(\mathbf{x})}{\partial x_2^2},$$

where the inequality holds by property (iv) of h (recall that $\lambda = a_2/a_1$). This shows that τ is minimal strongly plurisubharmonic on P_{c_0} . The other properties of τ follow immediately from the properties of h . (Compare with the proof of [26, Lemma 3.11.1, p. 98].) Condition (c) shows that τ is minimal strongly plurisubharmonic also on the complement of P_{c_0} . Condition (d) obviously holds on P_{c_0} , while on the complement of P_{c_0} it follows from (c) and the assumptions on ρ . □

Combining Proposition 3.3 and Lemma 3.6, we now prove the following lemma, which provides the induction step in the proof of Theorem 1.1.

Lemma 3.7. *Let ρ be a minimal strongly plurisubharmonic function on a domain $D \subset \mathbb{R}^3$, and let $a < b$ be real numbers such that the set*

$$(3.14) \quad D_{a,b} = \{\mathbf{x} \in D : a < \rho(\mathbf{x}) < b\}$$

is relatively compact in D . Given numbers $0 < \eta < b - a$, $\epsilon > 0$, $\delta > 0$, a conformal minimal immersion $F: M \rightarrow D$ such that $F(bM) \subset D_{a,b}$, a point $p_0 \in \overset{\circ}{M}$, a number $d > 0$, and a compact set $K \subset \overset{\circ}{M}$, there exists a conformal minimal immersion $G: M \rightarrow D$ satisfying the following conditions:

- (a) $G(bM) \subset D_{b-\eta,b}$ (equivalently, $b - \eta < \rho(G(\zeta)) < b$ for every $\zeta \in bM$);
- (b) $\rho(G(\zeta)) \geq \rho(F(\zeta)) - \delta$ for every $\zeta \in M$;
- (c) $\|G - F\|_{1,K} < \epsilon$;
- (d) $\text{dist}_G(p_0, bM) > d$;
- (e) $\text{Flux}(G) = \text{Flux}(F)$.

Proof. If the domain $D_{a,b}$ (3.14) does not contain any critical points of ρ , then a finite number of applications of Proposition 3.3 furnishes a conformal minimal immersion $G: M \rightarrow D$ satisfying all conditions except (d); this last condition can be achieved by an arbitrarily \mathcal{C}^0 small deformation of G using [2, Lemma 4.1]. (The cited lemma allows one to increase the interior boundary distance of a conformal

minimal immersion by an arbitrarily big amount while staying arbitrarily \mathcal{C}^0 -close to the given map.)

Assume now that $\mathbf{x}_1, \dots, \mathbf{x}_m$ are the (nice) critical points of ρ in $D_{a,b}$ (3.14). We may assume that the numbers $c_j = \rho(\mathbf{x}_j)$ are distinct, and we enumerate the points so that $a < c_1 < c_2 < \dots < c_m < b$. We may also assume that $\min_{bM} \rho \circ F \leq c_1$, since otherwise a may be replaced by a constant satisfying $c_1 < a < \min_{bM} \rho \circ F$.

Pick $c_0 > 0$ such that the conclusion of Lemma 3.6 applies to the critical point \mathbf{x}_1 of ρ and the constant c_0 . Applying Proposition 3.3 finitely many times, we can replace F by a conformal minimal immersion $F_1: M \rightarrow D$ such that $F_1(bM) \subset D_{c_1-c_0,b}$ and F_1 satisfies conditions (b), (c), and (e) in Lemma 3.7 (with F_1 in place of G , and for some new constants ϵ_1 and δ_1 in place of ϵ and δ). By general position, we can assume that $F_1(bM)$ avoids the local stable manifold E (see (3.11)) of the point \mathbf{x}_1 . Let τ be the function furnished by Lemma 3.6 (for the point \mathbf{x}_1 and the constant c_0). Applying Proposition 3.3 with the function τ finitely many times, we can lift the boundary $F_1(bM)$ above the level $c_1 = \rho(\mathbf{x}_1)$ and thus obtain a new conformal minimal immersion $G_1: M \rightarrow D$ satisfying $G_1(bM) \subset D_{c_1,b}$. As before, G_1 is chosen to satisfy conditions (b), (c), and (e) in Lemma 3.7, with G_1 in place of G and F_1 in place of F (and for some new constants $\epsilon_2 > 0, \delta_2 > 0$).

Now, we repeat the same procedure, first using Proposition 3.3 to push $G_1(bM)$ close to the level $\rho = c_2$, and subsequently lifting the boundary across $\rho = c_2$ by using Lemma 3.6. This furnishes a conformal minimal immersion $G_2: M \rightarrow D$ with $G_2(bM) \subset D_{c_2,b}$.

In finitely many steps of this kind we find a conformal minimal immersion $G: M \rightarrow D$ satisfying $G(bM) \subset D_{b-\eta,b}$ (condition (a)) and condition (e). Since the number of steps depends only on the geometry of ρ , we can fulfill conditions (b) and (c) by choosing the corresponding numbers $\epsilon_j > 0$ and $\delta_j > 0$ sufficiently small at every step. Finally, condition (d) is achieved as in the special case by appealing to [2, Lemma 4.1]. \square

Proof of Theorem 1.1. Let $F_0: M \rightarrow D$ be a conformal minimal immersion and let K be a compact set in \mathring{M} . Given $\epsilon > 0$, we shall find a complete proper conformal minimal immersion $F: \mathring{M} \rightarrow D$ satisfying $\|F - F_0\|_{0,K} = \sup_{\zeta \in K} \|F(\zeta) - F_0(\zeta)\| < \epsilon$. Such F will be found as the limit $F = \lim_{j \rightarrow \infty} F_j$ of a sequence of conformal minimal immersions $F_j: M \rightarrow D$ that will be constructed by an inductive application of Lemma 3.7.

Choose a minimal strongly plurisubharmonic Morse exhaustion function $\rho: D \rightarrow \mathbb{R}$. Let $P = \{\mathbf{x}_1, \mathbf{x}_2, \dots\} \subset D$ be the (discrete) critical locus of ρ , where the points \mathbf{x}_j are enumerated so that $\rho(\mathbf{x}_1) < \rho(\mathbf{x}_2) < \dots$. By Lemma 3.5, we may assume that every \mathbf{x}_j is a nice critical point of ρ . Pick increasing sequences $a_1 < a_2 < a_3 \dots$ and $d_1 < d_2 < d_3 \dots$ such that $\sup_M \rho \circ F_0 < a_1$, $\lim_{j \rightarrow \infty} a_j = +\infty$, and $\lim_{j \rightarrow \infty} d_j = +\infty$. Also, choose a decreasing sequence $\delta_j > 0$ with $\delta = \sum_{j=1}^\infty \delta_j < \infty$. Fix a point $p_0 \in \mathring{K}$. We shall construct a sequence of smooth conformal minimal immersions $F_j: M \rightarrow D$, an increasing sequence of compacts $K = K_0 \subset K_1 \subset \dots \subset \bigcup_{j=1}^\infty K_j = \mathring{M}$, and a decreasing sequence of positive numbers $\epsilon_j > 0$ such that the following conditions hold for every $j = 1, 2, \dots$:

- (i)_j $a_j < \rho \circ F_j < a_{j+1}$ on $M \setminus K_j$;
- (ii)_j $\rho \circ F_j > \rho \circ F_{j-1} - \delta_j$ on M ;
- (iii)_j $\|F_j - F_{j-1}\|_{1,K_{j-1}} < \epsilon_j$;

- (iv_j) $\text{dist}_{F_j}(p_0, M \setminus K_j) > d_j$;
- (v_j) $\text{Flux}(F_j) = \text{Flux}(F_{j-1})$;
- (vi_j) $\epsilon_j < 2^{-1} \min\{\epsilon_{j-1}, \text{dist}(F_{j-1}(M), bD), \inf_{\zeta \in K_{j-1}} \|dF_{j-1}(\zeta)\|\}$.

To begin the induction, set $\epsilon_0 = \epsilon/2$ and $K = K_0$. Assume inductively that, for some $j \in \mathbb{N}$, we have found maps F_0, \dots, F_{j-1} , numbers $\epsilon_0, \dots, \epsilon_{j-1}$, and compact sets K_0, \dots, K_{j-1} such that the above properties hold. Pick a number $\epsilon_j > 0$ satisfying condition (vi_j). Applying Lemma 3.7 with the data $(F_{j-1}, K_{j-1}, \epsilon_j, d_j)$ furnishes a conformal minimal immersion $F_j: M \rightarrow D$ satisfying condition (i_j) on the boundary bM , conditions (ii_j), (iii_j), (v_j), and such that $\text{dist}_{F_j}(p_0, bM) > d_j$. Next, pick a compact set $K_j \subset \overset{\circ}{M}$ such that $K_{j-1} \subset \overset{\circ}{K}_j$ and conditions (i_j) and (iv_j) hold. (It suffices to take K_j big enough.) This completes the induction step.

Condition (vi_j) implies that $\sum_{k=j+1}^{\infty} \epsilon_k < \epsilon_j$ for every $j = 0, 1, \dots$; in particular, $\sum_{k=0}^{\infty} \epsilon_k < 2\epsilon_0 = \epsilon$. Condition (iii_j) ensures that the sequence F_j converges uniformly on compacts in $\bigcup_{j=1}^{\infty} K_j = M$ to a harmonic map $F = \lim_{j \rightarrow \infty} F_j: M \rightarrow \overline{D}$. Conditions (iii_j) and (vi_j) show that for every $j = 0, 1, \dots$ we have that

$$(3.15) \quad \|F - F_j\|_{1, K_j} \leq \sum_{k=j}^{\infty} \|F_{k+1} - F_k\|_{1, K_j} < \sum_{k=j}^{\infty} \epsilon_{k+1} < 2\epsilon_{j+1} < \epsilon_j.$$

In particular, $\|F - F_0\|_{0, K} < \epsilon$. The estimate (3.15), together with (vi_{j+1}), also shows that $F(K_j) \subset D$; since this holds for all j , we have $F(\overset{\circ}{M}) \subset D$. Since $2\epsilon_{j+1} < \inf_{\zeta \in K_j} \|dF_j(\zeta)\|$ by (vi_{j+1}), it follows from (3.15) that F is a conformal immersion on K_j . As this holds for all j , $F: \overset{\circ}{M} \rightarrow D$ is a conformal harmonic (hence minimal) immersion. In view of (v), we have $\text{Flux}(F) = \text{Flux}(F_0)$. Finally, conditions (i_j)–(iii_j) ensure that F is proper into D , while conditions (iii_j) and (iv_j) show that F is complete. □

Proof of Theorem 1.7. The proof is the same as that of Theorem 1.1 modulo the obvious modifications, replacing conditions pertaining to the distance from bD (see condition (vi_j) above) by the corresponding conditions pertaining to the distance from the end of the domain Ω on which the function ρ tends to $+\infty$. □

Proof of Theorem 1.9. Choose a minimal strongly plurisubharmonic function ρ on an open set $D' \supset \overline{D}$ such that $D = \{\mathbf{x} \in D' : \rho(\mathbf{x}) < 0\}$ and $d\rho \neq 0$ on $bD = \{\rho = 0\}$. Pick $\eta > 0$ such that the set $\{\rho < \eta\}$ is relatively compact in D' and $d\rho \neq 0$ on the compact set

$$(3.16) \quad L = \{\mathbf{x} \in D' : -\eta \leq \rho(\mathbf{x}) \leq \eta\}.$$

Let $C_0 > 0$ be a constant satisfying the conclusion of Proposition 3.3 for the data (D', ρ, L) . In view of Theorem 1.1, we may assume that the given conformal minimal immersion $F_0: M \rightarrow D$ satisfies

$$(3.17) \quad a_0 = a_0(F_0) := \inf_{\zeta \in bM} \rho(F_0(\zeta)) > -\eta.$$

(Equivalently, $F_0(bM) \subset D \cap \overset{\circ}{L}$.) For every $j = 0, 1, 2, \dots$ we set

$$a_j = 2^{-j} a_0, \quad \eta_j = a_{j+1} - a_j = 2^{-j-1} |a_0|.$$

Pick an increasing sequence $0 < d_1 < d_2 < \dots$ with $\lim_{j \rightarrow \infty} d_j = +\infty$ and a decreasing sequence $\delta_j > 0$ with $\delta = \sum_{j=1}^{\infty} \delta_j < \infty$. By following the proof of Theorem 1.1, using also the estimate (4) in Proposition 3.3 with the constant C_0

introduced above, we find a sequence of conformal minimal immersions $F_j: M \rightarrow D$ ($j = 1, 2, \dots$), an increasing sequence of compacts $K = K_0 \subset K_1 \subset \dots \subset \bigcup_{j=1}^\infty K_j = \overset{\circ}{M}$, and a decreasing sequence of numbers $\epsilon_j > 0$ such that the following conditions hold for all $j = 1, 2, \dots$:

- (i_j) $\rho \circ F_j > a_j$ on $M \setminus K_j$;
- (ii_j) $\rho \circ F_j > \rho \circ F_{j-1} - \delta_j$ on M ;
- (iii_j) $\|F_j - F_{j-1}\|_{1, K_{j-1}} < \epsilon_j$;
- (iv_j) $\text{dist}_{F_j}(p_0, M \setminus K_j) > d_j$;
- (v_j) $\text{Flux}(F_j) = \text{Flux}(F_{j-1})$;
- (vi_j) $\epsilon_j < 2^{-1} \min\{\epsilon_{j-1}, \text{dist}(F_{j-1}(M), bD), \inf_{\zeta \in K_{j-1}} \|dF_{j-1}(\zeta)\|\}$;
- (vii_j) $\|F_j - F_{j-1}\|_{0, M} \leq C_0 \sqrt{2\eta_j} = C_0 \sqrt{2^{-j}} \sqrt{|a_0|}$.

These properties correspond to those in the proof of Theorem 1.1, except that condition (i_j) is adjusted to the present setting, and the additional condition (vii_j) follows from the estimate (4) in Proposition 3.3. (By [2, Lemma 4.1], condition (iv_j) can be achieved by a deformation which is arbitrarily small in the $\mathcal{C}^0(M)$ norm, and the error made by this deformation is absorbed by the constant C_0 in (vii_j).

Set $C_1 = C_0 \sum_{j=1}^\infty \sqrt{2^{-j}}$. Condition (vii_j) ensures that the sequence F_j converges uniformly on M to a continuous map $F: M \rightarrow \overline{D}$ satisfying $\|F - F_0\|_{0, M} \leq C_1 \sqrt{|a_0|}$. On the set L (3.16) the function $|\rho|$ is proportional to the distance from bD , so the number $|a_0|$, defined by (3.17), is proportional to $\max_{\zeta \in bM} \text{dist}(F_0(\zeta), bD)$. This gives the estimate (1.3) in Theorem 1.9 for a suitable choice of the constant $C > 0$ which depends only on the geometry of ρ in L . We can see as in the proof of Theorem 1.1 that $F|_{\overset{\circ}{M}}: \overset{\circ}{M} \rightarrow D$ is a proper complete conformal minimal immersion. \square

Proof of Corollary 1.15. This follows from Lemma 3.7 by a similar inductive procedure as those in the proofs of Theorems 1.1 and 1.7. In this case we use in addition the Mergelyan approximation theorem for conformal minimal immersions (cf. [5, 6]) in order to add either a handle or an end to the surface at each step in the recursive construction. In this way, we may prescribe the topology of the limit surface. For the details of this construction, we refer to the proof of Theorem 1.4(b) and Corollary 1.5(b) in [2]. \square

Remark 3.8. The methods developed in [2] and in this paper allow us to generalize Theorems 1.1 and 1.9 to $(n - 2)$ -convex domains $D \subset \mathbb{R}^n$ for any $n > 3$. We shall not state these generalizations but will give a brief sketch of the proof. By definition, such a domain admits a smooth strongly $(n - 2)$ -plurisubharmonic exhaustion function $\rho: D \rightarrow \mathbb{R}$ (see Definition 2.5 and Proposition 2.6). Furthermore, by convexifying in the normal direction, ρ can be chosen such that the level sets $S_c = \{\rho = c\}$ for noncritical values of ρ are strongly $(n - 2)$ -convex hypersurfaces, which means in particular that at every point $p_0 \in S_c$ there is a 2-dimensional plane $L \subset T_{p_0} S_c$ on which Hess_ρ is strongly positive. (See Definition 1.8.) By choosing suitably shaped small flat discs $\Delta_p \subset \mathbb{R}^n$ for points p near p_0 , lying in affine 2-planes parallel to L , and solving the associated Riemann-Hilbert boundary value problem (see [2, Theorem 3.6]), one can lift a small part of the boundary of any conformal minimal disc $F: \mathbb{D} \rightarrow D$ in a neighborhood of p_0 to a higher level set of ρ (see Proposition 3.3). The rest of the proof goes through as before.

However, if the level sets of ρ are merely $(n - 1)$ -convex (which is the same as mean-convex), this approach would require the existence of approximate solutions of the Riemann-Hilbert boundary value problem for nonflat conformal minimal discs (i.e., the analogue of [2, Lemma 3.1] for $n > 3$). We are unable to prove optimal results for $n > 3$ at this time.

The corresponding optimal results in complex analysis, pertaining to the existence of proper holomorphic maps from strongly pseudoconvex Stein domains to q -convex manifolds, were obtained in the papers [18, 19].

4. MAXIMAL MINIMALLY CONVEX DOMAINS AND A MAXIMUM PRINCIPLE AT INFINITY

This section is devoted to the proof of Theorem 1.16. The main ingredient is the maximum principle for minimal surfaces with finite total curvature in minimally convex domains in \mathbb{R}^3 , given by the following theorem.

Theorem 4.1. *Assume that $S \subset \mathbb{R}^3$ is a complete, connected, immersed minimal surface with compact boundary $bS \neq \emptyset$ and finite total curvature. If $D \subset \mathbb{R}^3$ is a smoothly bounded minimally convex domain containing S , then $\text{dist}(S, \mathbb{R}^3 \setminus D) = \text{dist}(bS, \mathbb{R}^3 \setminus D)$.*

The particular case of Theorem 4.1 when S is compact (with boundary) is already ensured by Proposition 2.11. The main difficulty in the general case (when the surface S is not compact) is that one must deal with the contact at infinity, a rather delicate task.

Before proving Theorem 4.1, we show how it implies Theorem 1.16 by a Kontinuitätssatz type argument (see Proposition 2.9).

Proof of Theorem 1.16, assuming Theorem 4.1. Assume that $D^c = \mathbb{R}^3 \setminus D \neq \emptyset$ and let us prove that S is a plane. Choose a relatively compact disc $\Omega \subset S$ and set $S' = S \setminus \Omega$. By Theorem 4.1 we have that $\text{dist}(S', D^c) = \text{dist}(bS', D^c)$, and hence $\text{dist}(S, D^c) = \text{dist}(\bar{\Omega}, D^c)$. Thus, there exist points $\mathbf{x}_0 \in S$ and $\mathbf{y}_0 \in bD$ such that $\|\mathbf{x}_0 - \mathbf{y}_0\| = \text{dist}(S, D^c)$, and we infer from Proposition 2.11 that S is a plane. Without loss of generality we may assume that $S = \{(x, y, z) \in \mathbb{R}^3 : z = 0\}$. Set $W_+ = \{(x, y, z) \in \mathbb{R}^3 : z > 0\}$. We claim that if the set $W_+ \setminus D$ is nonempty, then it is a half-space. Indeed, assume that $W_+ \setminus D \neq \emptyset$ and let us prove first that

$$(4.1) \quad d_+ := \text{dist}(S, bD \cap W_+) = \text{dist}(S, W_+ \setminus D) > 0.$$

Consider the family of vertical negative half-catenoids

$$\Sigma_a = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 = a^2 \cosh^2(z/a), z \leq 0\}, \quad 0 < a \leq 1.$$

Let A_+ denote the cylinder

$$A_+ := \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 \leq 2, 0 \leq z \leq \tau_+\},$$

where $\tau_+ > 0$ is chosen small enough such that $((0, 0, \tau_+) + \Sigma_1) \cap \{z \geq 0\} \subset A_+ \subset D$. The Kontinuitätssatz for minimal surfaces (cf. Proposition 2.9) implies that

$$\Sigma_a^+ := ((0, 0, \tau_+) + \Sigma_a) \cap \{z \geq 0\} \subset D \quad \text{for all } 0 < a \leq 1.$$

Indeed, Σ_a^+ are minimal surfaces with boundaries in $A_+ \cup S \subset D$ and $\Sigma_1^+ \subset A_+ \subset D$. It is easily seen that $\bigcup_{0 < a \leq 1} \Sigma_a^+$ contains the set $\{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 \geq 2, 0 \leq z < \tau_+\}$. Since $A_+ \subset D$, we infer that D contains the slab $\{0 \leq z < \tau_+\}$. This implies that $d_+ \geq \tau_+ > 0$, thereby proving (4.1).

If there is a point $(x_0, y_0, d_+) \in bD$, then, arguing as in the proof of Proposition 2.11 and using that D is connected, we easily infer that the plane $\Pi_+ := \{z = d_+\}$ lies in $bD \cap \{z > 0\} = b(W_+ \setminus D)$, and hence $W_+ \setminus D$ is a half-space. Otherwise, $\Pi_+ \subset D$, and we may reason as above (replacing S by Π_+) to see that $\text{dist}(\Pi_+, W_+ \setminus D) > 0$ in contradiction to (4.1).

A symmetric argument guarantees that $\{(x, y, z) \in \mathbb{R}^3 : z < 0\} \setminus D$ is either empty or a half-space. This concludes the proof. \square

The proof of Theorem 4.1 also follows from a Kontinuitätssatz argument; however, the construction of a suitable family of minimal surfaces is much more delicate. The surfaces will be multigraphs, obtained as solutions of suitable Dirichlet problems for the minimal surface equation over finite coverings of annuli in \mathbb{R}^2 ; see Lemma 4.3.

Before going into the construction, we introduce some notation.

Definition 4.2. For each pair of numbers $0 \leq R_0 < R \leq +\infty$ we set

$$A_{R_0,R} := \{(x, y) \in \mathbb{R}^2 : R_0 < \|(x, y)\| < R\}, \quad A_{R_0} = A_{R_0,+\infty}.$$

Endow $A_0 = \mathbb{R}^2 \setminus \{(0, 0)\}$ with the Euclidean metric and denote by

$$\pi_n : A_0^n \rightarrow A_0$$

the n -sheeted isometric covering, $n \in \mathbb{N}$. We also set:

- $A_{R_0,R}^n := \pi_n^{-1}(A_{R_0,R})$ for all $0 \leq R_0 < R < +\infty$ and $A_{R_0}^n = \pi_n^{-1}(A_{R_0})$;
- $c_R := \{(x, y) \in \mathbb{R}^2 : \|(x, y)\| = R\}$ and $c_R^n := \pi_n^{-1}(c_R)$, $R > 0$.

Obviously, $bA_{R_0}^n = c_{R_0}^n$ and $bA_{R_0,R}^n = c_{R_0}^n \cup c_R^n$, $0 < R_0 < R < +\infty$, $n \in \mathbb{N}$.

A function $u \in \mathcal{C}^2(A_{R_0,R}^n)$ is said to *satisfy the minimal surface equation in $A_{R_0,R}^n$* if

$$\text{div}\left(\frac{\nabla u}{\sqrt{1 + \|\nabla u\|^2}}\right) = 0 \quad \text{in } A_{R_0,R}^n,$$

equivalently, if $\{(p, u(p)) : p \in A_{R_0,R}^n\}$ is a minimal surface (a minimal multigraph).

Given $\phi \in \mathcal{C}^0(bA_{R_0,R}^n)$, a function $u \in \mathcal{C}^2(A_{R_0,R}^n) \cap \mathcal{C}^0(\overline{A_{R_0,R}^n})$ is said to be a *solution of the Dirichlet problem for the minimal surface equation in $A_{R_0,R}^n$ with boundary data ϕ* if u satisfies the minimal surface equation in $A_{R_0,R}^n$ and the boundary condition $u|_{bA_{R_0,R}^n} = \phi$.

Lemma 4.3. Let $0 < R_0 < R_1$, $K \in (0, 1)$, and $n \in \mathbb{N}$. There exists a number $\epsilon > 0$, depending only on R_0 , R_1 , and K , such that the following holds. If $R \geq R_1$, $\delta \in [0, \epsilon]$,

- (a) $v : \overline{A_{R_0}^n} \rightarrow \mathbb{R}$ is a real analytic solution of the minimal surface equation in $A_{R_0}^n$,
- (b) $\|\nabla v\| < K/2$ in $\overline{A_{R_0}^n}$, and
- (c) we set $\phi_{R,\delta} : bA_{R_0,R}^n \rightarrow \mathbb{R}$, $\phi_{R,\delta} = v$ in $c_{R_0}^n$, and $\phi_{R,\delta} = v + \delta$ in c_R^n ,

then the Dirichlet problem for the minimal surface equation in $A_{R_0,R}^n$ with boundary data $\phi_{R,\delta}$ has a unique solution $u_{R,\delta}$. Furthermore, $u_{R,\delta}$ enjoys the following conditions:

- (i) $v \leq u_{R,\delta} \leq v + \delta$ on $\overline{A_{R_0,R}^n}$;
- (ii) $u_{R,\delta}$ depends continuously on $(R, \delta) \in [R_1, +\infty) \times [0, \epsilon]$;
- (iii) $\{u_{R,\delta}\}_{R > R_1} \rightarrow v$ as $R \rightarrow +\infty$ on compact subsets of $\overline{A_{R_0}^n}$ for all $\delta \in [0, \epsilon]$.

The number $\epsilon > 0$ in the lemma will only depend on the existence of suitable barrier functions ν_{p_0} at boundary points $p_0 \in bA_{R_0,R}^n$ adapted to our problem. The construction of these barrier functions in turn only depends on the constants R_0, R_1 , and K .

Proof. For the existence part of the lemma, we use Perron’s method for the minimal surface equation; see for instance [29, 47].

Take arbitrary numbers $R > R_0 > 0$ and $\delta \geq 0$. If $w \in \mathcal{C}^0(\overline{A}_{R_0,R}^n)$, D is a convex disc in $\overline{A}_{R_0,R}^n$, and w_D is the solution of the minimal surface equation in D which equals w on bD (such exists by classical Rado’s theorem), we denote by \widehat{w}_D the continuous function in $\overline{A}_{R_0,R}^n$ which coincides with w in $\overline{A}_{R_0,R}^n \setminus D$ and with w_D in D .

By definition, a function $w \in \mathcal{C}^0(\overline{A}_{R_0,R}^n)$ is said to be a sub-solution of the Dirichlet problem for the minimal surface equation in $\overline{A}_{R_0,R}^n$, defined by $\phi_{R,\delta}$ given in (c), if $w \leq \phi_{R,\delta}$ in $bA_{R_0,R}^n$ and $w \leq \widehat{w}_D$ in D (hence in $\overline{A}_{R_0,R}^n$) for all discs D as above. We denote by $\mathcal{F}_{R,\delta}^-$ the family of all such sub-solutions. Likewise, w is said to be a super-solution for this problem if $w \geq \phi_{R,\delta}$ in $bA_{R_0,R}^n$ and $w \geq \widehat{w}_D$ in D (hence in $\overline{A}_{R_0,R}^n$) for all discs D in $\overline{A}_{R_0,R}^n$. The corresponding space of super-solutions will be denoted by $\mathcal{F}_{R,\delta}^+$. Note that $v|_{\overline{A}_{R_0,R}^n} \in \mathcal{F}_{R,\delta}^-$ and $(v + \delta)|_{\overline{A}_{R_0,R}^n} \in \mathcal{F}_{R,\delta}^+$ for all $R > R_0$ and $\delta > 0$, where v is the function given in item (a) in the statement of the lemma; hence these are nonempty families. If w_1 is a sub-solution and w_2 is a super-solution, then the maximum principle ensures that $w_1 \leq w_2$. On the other hand, if w_1 and w_2 are sub-solutions (respectively, super-solutions), then $\max\{w_1, w_2\}$ (respectively, $\min\{w_1, w_2\}$) also is.

We define

$$(4.2) \quad u_{R,\delta}: A_{R_0,R}^n \rightarrow \mathbb{R}, \quad u_{R,\delta}(p) = \sup_{w \in \mathcal{F}_{R,\delta}^-} w(p).$$

It is well known that $u_{R,\delta}$ is a solution of the minimal surface equation in $A_{R_0,R}^n$. Further,

$$(4.3) \quad w_1 \leq u_{R,\delta} \leq w_2 \quad \text{for any } w_1 \in \mathcal{F}_{R,\delta}^- \text{ and } w_2 \in \mathcal{F}_{R,\delta}^+.$$

In particular,

$$(4.4) \quad v \leq u_{R,\delta} \leq v + \delta \quad \text{in } A_{R_0,R}^n \text{ for all } R > R_0 \text{ and all } \delta \geq 0.$$

Claim 4.4. Given numbers $R_1 > R_0 > 0$ and $K \in (0, 1)$, there exists $\epsilon > 0$, depending only on R_0, R_1 , and K , such that the following holds. If $R \geq R_1$ and $0 \leq \delta \leq \epsilon$, then the function $u_{R,\delta}$ given by (4.2) is a solution to the Dirichlet problem for the minimal surface equation with boundary data $\phi_{R,\delta}$ in $A_{R_0,R}^n$; that is to say,

$$\lim_{p \rightarrow p_0} u_{R,\delta}(p) = \phi_{R,\delta}(p_0) \quad \text{for all } p_0 \in bA_{R_0,R}^n.$$

Proof. Choose $R \geq R_1$ and a point $p_0 \in bA_{R_0,R}^n$, and let us distinguish cases.

Case 1 ($p_0 \in c_{R_0}^n$). Let us prove the existence of a number $\epsilon_1 > 0$, depending only on R_0, R_1 , and K , for which the following statement holds. Given $\delta \in [0, \epsilon_1]$ there exists $\nu_{p_0} \in \mathcal{C}^0(\overline{A}_{R_0}^n)$ such that $\nu_{p_0}(p_0) = v(p_0) = \phi_{R,\delta}(p_0)$ and $\nu_{p_0}|_{\overline{A}_{R_0,R}^n} \in \mathcal{F}_{R,\delta}^+$.

Indeed, write $\pi_n(p_0) = (x_0, y_0) \in c_{R_0}$. Set $C_{R_0} := \{(x, y, z) \in \mathbb{R}^3 : \|(x, y)\| < R_0\}$ and $J_K := \{(x, y, z) \in \mathbb{R}^3 : 0 \leq z - v(p_0) = K\|(x - x_0, y - y_0)\|\}$. Pick $\mu > 0$ small enough in terms of R_0 , R_1 , and K , such that the set

$$\gamma := ((bC_{R_0} \cap J_K) \cap \{0 \leq z - v(p_0) \leq \mu\}) \cup ((J_K \setminus C_{R_0}) \cap \{z - v(p_0) = \mu\})$$

is a Jordan curve. It follows that γ has one-to-one orthogonal projection γ_0 to the plane $\{z = 0\} \equiv \mathbb{R}^2$. Ensure also that γ_0 bounds a topological disc $U \subset \mathbb{R}^2$ with $\overline{U} \subset A_{R_0, R_1} \cup c_{R_0}$. Thus, $\pi_n^{-1}(U)$ consists of n disjoint isometric copies of U ; write $\widehat{U} \subset A_0^n$ for the connected component of $\pi_n^{-1}(U)$ containing p_0 . Denote by $\phi: bU \rightarrow \mathbb{R}$ the (unique) continuous function such that $\{(p, \phi(p)) : p \in bU = \gamma_0\} = \gamma$.

Further, the domain U satisfies an exterior sphere condition with radius R_0 (cf. [56, Definition 1.4(i)]), and thus, if $\mu > 0$ is sufficiently small in terms of R_0 , R_1 , and K , the Dirichlet problem for the minimal surface equation in U with boundary data ϕ has a unique solution $f: U \rightarrow \mathbb{R}$ satisfying $(f \circ \pi_n)(p_0) = \phi(p_0) = v(p_0)$ and $f \circ \pi_n > v$ in $\widehat{U} \setminus \{p_0\}$; see [56] and take into account condition (b) in the statement of the lemma and that $\gamma \subset J_K$. It follows that

$$\inf\{(f \circ \pi_n)(p) - v(p) : p \in b\widehat{U} \setminus c_{R_0}^n\} = \inf\{v(p_0) + \mu - v(p) : p \in b\widehat{U} \setminus c_{R_0}^n\} > 0.$$

Finally, take $\epsilon_1 > 0$ smaller than this infimum. Note that this number does not depend on v ; take into account (b). Further, since μ depends on R_0 , R_1 , and K but not on $p_0 \in c_{R_0}^n$, the same holds for ϵ_1 . It suffices to set $\nu_{p_0}: \overline{A}_{R_0}^n \rightarrow \mathbb{R}$, $\nu_{p_0} = \min\{f \circ \pi_n, v + \epsilon_1\}$ on \widehat{U} , and $\nu_{p_0} = v + \epsilon_1$ on $\overline{A}_{R_0}^n \setminus \widehat{U}$.

Since $\nu_{p_0}|_{\overline{A}_{R_0, R}^n} \in \mathcal{F}_{R, \delta}^+$, $\delta \in [0, \epsilon_1]$, and $\nu_{p_0}(p_0) = v(p_0) = \phi_{R, \delta}(p_0)$, the bounds (4.3) and (4.4) trivially ensure that $\lim_{p \rightarrow p_0} u_{R, \delta}(p) = \phi_{R, \delta}(p_0)$.

Case 2 ($p_0 \in c_R^n$). Let us now prove the existence of a number $\epsilon_2 > 0$, depending only on R_0 , R_1 , and K , for which the following statement holds. Given $\delta \in [0, \epsilon_2]$, there exists $\nu_{p_0} \in \mathcal{C}^0(\overline{A}_{R_0, R}^n) \cap \mathcal{F}_{R, \delta}^-$ such that $\nu_{p_0}(p_0) = v(p_0) + \delta = \phi_{R, \delta}(p_0)$.

Indeed, consider a small disc $V \subset \mathbb{R}^2$ centered at the origin and with radius less than $R_1 - R_0$, set $U_R := (\pi_n(p_0) + V) \cap \overline{A}_{R_0, R}^n$, and let $\widehat{U}_R \subset \overline{A}_{R_0, R}^n$ denote the connected component of $\pi_n^{-1}(U_R)$ containing p_0 ; obviously, $\pi_n|_{\widehat{U}_R}: \widehat{U}_R \rightarrow U_R$ is an isometry. Choose a linear function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ satisfying $(f \circ \pi_n)(p_0) = v(p_0)$ and $f \circ \pi_n < v$ on $\widehat{U}_R \setminus \{p_0\}$, and take $0 < \epsilon_2 < \inf\{v(p) - (f \circ \pi_n)(p) : p \in b\widehat{U} \setminus c_R^n\}$. The existence of such an ϵ_2 follows from item (b), and it can be chosen depending on neither v nor R (it only depends on R_0 , R_1 , and K). Given $\delta \in [0, \epsilon_2]$, it suffices to set $\nu_{p_0}: \overline{A}_{R_0, R}^n \rightarrow \mathbb{R}$, $\nu_{p_0} = \max\{f \circ \pi_n + \delta, v + \delta - \epsilon_2\}$ on \widehat{U}_R , and $\nu_{p_0} = v + \delta - \epsilon_2$ on $\overline{A}_{R_0, R}^n \setminus \widehat{U}_R$.

As above, since $\nu_{p_0} \in \mathcal{F}_{R, \delta}^-$ and $\nu_{p_0}(p_0) = v(p_0) + \delta = \phi_{R, \delta}(p_0)$, (4.3) and (4.4) guarantee that $\lim_{p \rightarrow p_0} u_{R, \delta}(p) = \phi_{R, \delta}(p_0)$.

To finish the proof of the claim, it suffices to choose $\epsilon := \min\{\epsilon_1, \epsilon_2\}$. \square

We continue with the proof of Lemma 4.3. In view of Claim 4.4 it remains to check that, given numbers $\delta \in [0, \epsilon]$ and $R \geq R_1$, the solution $u_{R, \delta}$ given by (4.2) is unique and satisfies conditions (i), (ii), and (iii). Uniqueness follows directly from the maximum principle. Property (i) is ensured by (4.4). Since the boundary data $\phi_{R, \delta}$ depend continuously on $(R, \delta) \in [R_1, +\infty) \times [0, \epsilon]$, the same holds for the solutions $u_{R, \delta}$, proving (ii). In order to prove (iii), fix a number $\delta \in [0, \epsilon]$ and take

any divergent sequence $\{R_j\}_{j \in \mathbb{N}} \subset [R_1, +\infty)$. By standard compactness results (see for instance [11]), the sequence $\{u_{R_j, \delta}\}_{j \in \mathbb{N}}$ converges uniformly on compact subsets of $\overline{A_{R_0}^n}$ to a solution u of the minimal surface equation in $A_{R_0}^n$ with boundary data $u = v$ on $c_{R_0}^n$. Furthermore, (4.4) gives that $v \leq u \leq v + \delta$, and hence $u = v$ by the maximum principle at infinity (see for instance [44]). This proves (iii) and concludes the proof. \square

Proof of Theorem 4.1. If S is compact, then the result follows from Proposition 2.11.

Assume now that S is not compact. Up to passing to the two sheeted orientable covering, we may assume that $S = X(M)$, where M is a noncompact Riemann surface with compact boundary $bM \neq \emptyset$ and $X : M \rightarrow \mathbb{R}^3$ is a complete conformal minimal immersion with finite total curvature. With this notation, we have $bS = X(bM)$.

The assumptions on S imply that M is of finite topology and of parabolic conformal type (in particular, its ends are annular conformal punctures), and the (conformal) Gauss map $N : M \rightarrow \mathbb{S}^2$ of X extends conformally to the ends; see [48]. Given an annular end $E \subset M$, $E \cong \mathbb{D} \setminus \{0\}$, let n_E denote the limit normal vector of $X(E)$ at infinity and let Π_E be the vectorial plane in \mathbb{R}^3 orthogonal to n_E .

It is also well known that the minimal immersion X is a proper map and all the ends are *finite sublinear multigraphs*; see [38]. The latter means that for any annular end E of M , there exists an open solid circular cylinder C , with axis parallel to n_E , such that:

- (i) $E \cap X^{-1}(\overline{C})$ is compact and contains bE ;
- (ii) $(\pi_E \circ X)|_{E \setminus X^{-1}(C)} : E \setminus X^{-1}(C) \rightarrow \Pi_E \setminus C$ is a finite covering, where $\pi_E : \mathbb{R}^3 \rightarrow \Pi_E$ is the orthogonal projection;
- (iii) $\lim_{j \rightarrow \infty} \frac{1}{\|X(p_j)\|} \langle n_E, X(p_j) \rangle = 0$ for any divergent sequence $\{p_j\}_{j \in \mathbb{N}} \subset E$.

Write w_E for the winding number of $X(E)$ at infinity; note that w_E is the degree of the covering $(\pi_E \circ X)|_{E \setminus X^{-1}(C)}$.

Recall that $bS \neq \emptyset$. If $D = \mathbb{R}^3$, there is nothing to prove. Assume now that $D^c \neq \emptyset$ and let

$$d := \text{dist}(S, D^c) < +\infty.$$

It suffices to prove the following claim:

$$(4.5) \quad \text{there exist points } \mathbf{x}_0 \in S \text{ and } \mathbf{y}_0 \in bD \text{ such that } \|\mathbf{x}_0 - \mathbf{y}_0\| = d.$$

Indeed, assume for a moment that this holds. If $\mathbf{x}_0 \in bS$, we are done. Otherwise, $\mathbf{x}_0 \in S \setminus bS$, and we infer from Proposition 2.11 that the surface S is flat and $\mathbf{y}_0 - \mathbf{x}_0 + S \subset bD$. Thus, $d = \text{dist}(bS, D^c)$, which concludes the proof of Theorem 4.1 provided that (4.5) holds.

We now prove the assertion (4.5). We reason by contradiction and assume that

$$(4.6) \quad \text{dist}(\mathbf{x}, D^c) > d \quad \text{for all } \mathbf{x} \in S.$$

Under this assumption, there exists an annular end $X(E) \subset S$ with $\text{dist}(X(E), D^c) = d$; recall that E is conformally equivalent to $\mathbb{D} \setminus \{0\}$. Set

$$E_t := t n_E + X(E) \quad \text{for all } t \geq 0.$$

In particular, $E_0 = X(E)$. Condition (4.6) ensures that

$$(4.7) \quad \bigcup_{t \in [0, d]} E_t \subset D.$$

Set $C_R := \{(x, y, z) \in \mathbb{R}^3 : \|(x, y)\| < R\}$ for $R > 0$. Up to a rigid motion, a shrinking of E , and taking $R_0 > 0$ large enough, we may assume that $n_E = (0, 0, -1)$, $X(bE) \subset bC_{R_0}$, and

$$(4.8) \quad \text{dist}(E_d, D^c) = 0.$$

Given $\delta \geq 0$, we set $\gamma_{R_0}(\delta) := bE_{d+\delta} = (d + \delta)n_E + bE_0 \subset bC_{R_0}$. Since $(\bigcup_{t \in [0, d]} E_t) \cap \overline{C}_R$ is compact for all $R \geq R_0$ (see (i)), condition (4.7) provides numbers $\epsilon > 0$ and $R_1 > R_0$ such that

$$(4.9) \quad \bigcup_{t \in [0, d+\epsilon]} (E_t \cap \overline{C}_{R_1}) \subset D.$$

In particular, $\gamma_{R_0}(\delta) \subset D$ for all $\delta \in [0, \epsilon]$. Set $\gamma_R(\delta) := E_{d+\delta} \cap bC_R$ for all $\delta \in [0, \epsilon]$ and $R > R_0$. For simplicity, write n for the winding number w_E of $X(E)$ as a multigraph over $A_{R_0} = \{(x, y) \in \mathbb{R}^2 : \|(x, y)\| > R_0\}$, and denote by $\phi_{R, \delta} : bA_{R_0, R}^n \rightarrow \mathbb{R}$ the unique analytic function satisfying

$$\gamma_{R_0}(\delta) \cup \gamma_R(0) = \{(p, \phi_{R, \delta}(p)) : p \in bA_{R_0, R}^n\} \text{ for all } \delta \in [0, \epsilon] \text{ and } R > R_0.$$

(See Definition 4.2 for the notation.) Likewise, let $v : \overline{A}_{R_0}^n \rightarrow \mathbb{R}$ denote the unique analytic function such that

$$E_d = \{(p, v(p)) : p \in \overline{A}_{R_0}^n\}.$$

Without loss of generality (increasing R_0 if necessary), we may assume that $\|\nabla v\| < 1/4$ on $\overline{A}_{R_0}^n$; see property (iii) above.

On the other hand, if $\epsilon > 0$ is chosen small enough, then Lemma 4.3 shows that the Dirichlet problem for the minimal surface equation in $A_{R_0, R}^n$ with boundary data $\phi_{R, \delta}$ has a unique solution $u_{R, \delta}$ for all $R \geq R_1$ and $\delta \in [0, \epsilon]$. Furthermore,

$$(4.10) \quad v - \delta \leq u_{R, \delta} \leq v \text{ in } A_{R_0, R}^n \text{ for all } R \geq R_1 \text{ and } \delta \in [0, \epsilon].$$

Fix a pair of numbers $\delta \in [0, \epsilon]$ and $R \geq R_1$, and set $T_{R, \delta} := \{(p, u_{R, \delta}(p)) : p \in \overline{A}_{R_0, R}^n\}$. Note that (4.9) and (4.10) guarantee that $T_{R_1, \delta} \subset D$, whereas (4.7) and (4.9) ensure that $bT_{R, \delta} = \gamma_{R_0}(\delta) \cup \gamma_R(0) \subset D$ for all $R \geq R_1$. Thus, the Kontinuitätssatz for minimal surfaces (Proposition 2.9) implies that $T_{R, \delta} \subset D$ for all $R \geq R_1$. Further, by Lemma 4.3 we have $T_{R, \delta} \rightarrow E_{d+\delta}$ uniformly on compact subsets as $R \rightarrow +\infty$, and hence $E_{d+\delta} \subset \overline{D}$. Since this holds for arbitrary $\delta \in [0, \epsilon]$, we infer that $\bigcup_{t \in [0, d+\epsilon]} E_t \subset \overline{D}$, and hence $\bigcup_{t \in [0, d+\epsilon]} E_t \subset D$. This contradicts (4.8) and thereby proves (4.5). \square

5. NULL HULLS IN \mathbb{C}^n AND MINIMAL HULLS IN \mathbb{R}^n

The approximate solution of the Riemann-Hilbert problem for null discs, furnished by [2, Lemma 3.3], allows us to extend the main results of the paper [21] to null hulls in \mathbb{C}^n and minimal hulls in \mathbb{R}^n , for any $n \geq 3$. We now explain these generalizations. The proofs are similar to those in [21] and are left to the reader.

We denote by $\mathfrak{N}(\mathbb{D}, \Omega)$ the set of all immersed null holomorphic discs $\overline{\mathbb{D}} \rightarrow \mathbb{C}^n$ with range in a domain $\Omega \subset \mathbb{C}^n$, and we write

$$\mathfrak{N}(\mathbb{D}, \Omega, \mathbf{z}) = \{F \in \mathfrak{N}(\mathbb{D}, \Omega) : F(0) = \mathbf{z}\} \quad \text{for } \mathbf{z} \in \Omega.$$

The case $n = 3$ of the next result is [21, Theorem 2.10]; the general case $n \geq 3$ is proved in exactly the same way by using [2, Lemma 3.3] instead of [21, Lemma 2.8].

Theorem 5.1 (Null plurisubharmonic minorant). *Let $\phi: \Omega \rightarrow \mathbb{R} \cup \{-\infty\}$ be an upper semicontinuous function on a domain $\Omega \subset \mathbb{C}^n$ ($n \geq 3$). Then the function*

$$(5.1) \quad u(\mathbf{z}) := \inf \left\{ \int_0^{2\pi} \phi(F(e^{it})) \frac{dt}{2\pi} : F \in \mathfrak{N}(\mathbb{D}, \Omega, \mathbf{z}) \right\}, \quad \mathbf{z} \in \Omega,$$

is null plurisubharmonic on Ω or identically $-\infty$. Moreover, u is the supremum of all null plurisubharmonic functions on Ω which are not greater than ϕ .

Remark 5.2. The disc functional P_ϕ in (5.1), which assigns to any holomorphic disc $F: \overline{\mathbb{D}} \rightarrow \Omega$ the average $P_\phi(F) = \int_0^{2\pi} \phi(F(e^{it})) \frac{dt}{2\pi} \in \mathbb{R} \cup \{-\infty\}$, is called the *Poisson functional* associated to the function ϕ . If we take all holomorphic discs $F: \overline{\mathbb{D}} \rightarrow \Omega$ with $F(0) = \mathbf{z}$ (instead of just null discs) in (5.1), we obtain the biggest plurisubharmonic function on Ω satisfying $u \leq \phi$. This fundamental result of Poletsky [49, 50] and Bu and Schachermayer [13] was generalized by Rosay [52, 53] to all complex manifolds Ω (see also Lárusson and Sigurdsson [39, 41]), and by Drinovec Drnovšek and Forstnerič [20, Theorem 1.1] to all locally irreducible complex space.

Given a domain $\omega \subset \mathbb{R}^n$, we denote by $\mathfrak{M}(\mathbb{D}, \omega)$ the set of all conformal minimal immersions $\overline{\mathbb{D}} \rightarrow \omega$. By using Theorem 5.1, along with the connection between null plurisubharmonic and minimal plurisubharmonic functions (see Lemma 2.14), we obtain the following result. The case $n = 3$ is [21, Theorem 4.5].

Theorem 5.3 (Minimal plurisubharmonic minorant). *Let ω be a domain in \mathbb{R}^n ($n \geq 3$), and let $\phi: \omega \rightarrow \mathbb{R} \cup \{-\infty\}$ be an upper semicontinuous function. Then the function*

$$(5.2) \quad u(\mathbf{x}) := \inf \left\{ \int_0^{2\pi} \phi(F(e^{it})) \frac{dt}{2\pi} : F \in \mathfrak{M}(\mathbb{D}, \omega), F(0) = \mathbf{x} \right\}, \quad \mathbf{x} \in \omega,$$

is minimal plurisubharmonic on ω or identically $-\infty$. Moreover, u is the supremum of the minimal plurisubharmonic functions on ω which are not greater than ϕ .

Definition 5.4. Let K be a compact set in \mathbb{C}^n , $n \geq 3$. The *null hull* of K is the set

$$(5.3) \quad \widehat{K}_{\mathfrak{N}} = \{\mathbf{z} \in \mathbb{C}^n : v(\mathbf{z}) \leq \max_K v \quad \forall v \in \mathfrak{NPsh}(\mathbb{C}^n)\}.$$

Note that $\widehat{K}_{\mathfrak{N}}$ is a special case of a \mathbb{G} -convex hull introduced by Harvey and Lawson in [33, Definition 4.3, p. 2434]. The maximum principle for subharmonic functions implies that for any bounded null holomorphic curve $A \subset \mathbb{C}^n$ with boundary $bA \subset K$ we have $A \subset \widehat{K}_{\mathfrak{N}}$. Since $\text{Psh}(\mathbb{C}^n) \subset \mathfrak{NPsh}(\mathbb{C}^n)$, we clearly have the inclusions

$$(5.4) \quad K \subset \widehat{K}_{\mathfrak{N}} \subset \widehat{K} \subset \text{Co}(K).$$

The polynomial hull \widehat{K} is rarely equal to the convex hull $\text{Co}(K)$, and in general we also have $\widehat{K}_{\mathfrak{N}} \neq \widehat{K}$ (see [21, Example 3.2]).

The following characterization of the null hull agrees with [21, Corollary 3.5] for $n = 3$. The proof in [21] holds in any dimension $n \geq 3$, the nontrivial direction being furnished by Theorem 5.1 in this paper. Recall that $|I|$ denotes the Lebesgue measure of a set $I \subset \mathbb{R}$.

Corollary 5.5. *Let K be a compact set in \mathbb{C}^n ($n \geq 3$), and let $\Omega \subset \mathbb{C}^n$ be a bounded pseudoconvex Runge domain containing K . A point $\mathbf{z} \in \Omega$ belongs to the null hull $\widehat{K}_{\mathfrak{N}}$ of K if and only if there exists a sequence of null discs $F_j \in \mathfrak{N}(\mathbb{D}, \Omega, \mathbf{z})$ such that*

$$(5.5) \quad |\{t \in [0, 2\pi] : \text{dist}(F_j(e^{it}), K) < 1/j\}| \geq 2\pi - 1/j, \quad j = 1, 2, \dots$$

Similarly, the following characterization of the minimal hull generalizes [21, Corollary 4.9] to any dimension $n \geq 3$. The nontrivial direction is furnished by Theorem 5.3.

Corollary 5.6. *Let D be a minimally convex domain in \mathbb{R}^n ($n \geq 3$), let K be a compact set in D , and let $\omega \Subset D$ be a relatively compact domain containing the minimal hull $\widehat{K}_{\mathfrak{M}, D}$ of K . A point $\mathbf{x} \in \omega$ belongs to $\widehat{K}_{\mathfrak{M}, D}$ if and only if there exists a sequence of conformal minimal discs $F_j: \mathbb{D} \rightarrow \omega$ such that, for every $j = 1, 2, \dots$, we have $F_j(0) = \mathbf{x}$ and (5.5).*

Remark 5.7. Recall (cf. Remark 1.6) that a smoothly bounded domain $D \subset \mathbb{R}^n$ is mean-convex if and only if it is $(n - 1)$ -convex. By the maximum principle (see Proposition 2.11), mean-convex domains containing a given compact set $K \subset \mathbb{R}^n$ are natural barriers for minimal hypersurfaces with boundaries in K . The smallest such barrier, if it exists, is called the *mean-convex hull* of K ; clearly it coincides with the $(n - 1)$ -convex hull \widehat{K}_{n-1} (see Definition 2.5). The main technique for finding the mean-convex hull is the *mean curvature flow* of hypersurfaces, introduced by Brakke [12]. For results on this subject we refer, among others, to the papers [14, 23, 24, 28, 36, 37, 45, 46, 55] and the monograph by Colding and Minicozzi [15]. Our proof of Corollary 5.6 relies on completely different ideas, but it applies only to the 2-convex hull (which equals the mean-convex hull only in dimension $n = 3$). We indicate the following natural question.

Problem 5.8. Let K be a compact set with smooth boundary in \mathbb{R}^n for some $n \geq 3$. Given a point $\mathbf{x}_0 \in \widehat{K}_{\mathfrak{N}}$, does there exist a conformal minimal disc $F: \mathbb{D} \rightarrow \mathbb{R}^n$ such that $F(0) = \mathbf{x}_0$ and $F(b\mathbb{D}) \subset K$?

Recall that the *Green current* on \mathbb{C} is defined on any 2-form $\alpha = adx \wedge dy$ by

$$G(\alpha) = -\frac{1}{2\pi} \int_{\mathbb{D}} \log |\cdot| \alpha = -\frac{1}{2\pi} \int_{\zeta \in \mathbb{D}} \log |\zeta| \cdot a(\zeta) dx \wedge dy.$$

Clearly, G is a positive current of bidimension $(1, 1)$ and $dd^c G = \sigma - \delta_0$, where σ is the normalized Lebesgue measure on the circle $\mathbb{T} = b\mathbb{D}$ and $\delta_{\mathbf{z}}$ denotes the point mass at \mathbf{z} . If $F: \mathbb{D} \rightarrow \mathbb{C}^n$ is a holomorphic disc, then F_*G is a positive current of bidimension $(1, 1)$ on \mathbb{C}^n satisfying $dd^c(F_*G) = F_*\sigma - \delta_{F(0)}$. (See Duval and Sibony [22, Example 4.9].)

Assume now that K is a compact set in \mathbb{C}^n , \mathbf{z} is a point in the null hull $\widehat{K}_{\mathfrak{N}}$, and $F_j: \mathbb{D} \rightarrow \mathbb{C}^n$ is a sequence of holomorphic null discs with centers $F_j(0) = \mathbf{z}$, furnished by Corollary 5.5. By Wold [57] (see also [21, Proof of Theorem 6.2]), the sequence of Green currents $T_j = (F_j)_*G$ on \mathbb{C}^n has a weakly convergent subsequence, and the limit current T satisfies $dd^c T = \mu - \delta_{\mathbf{z}}$ where μ is a probability

measure on K . This generalizes the characterization of the null hull of a compact set in \mathbb{C}^3 by null positive Green currents, given by [21, Theorem 6.2], to any dimension $n \geq 3$. Similarly, applying the above argument to the sequence of conformal minimal discs $F_j: \overline{\mathbb{D}} \rightarrow \mathbb{R}^n$ furnished by Corollary 5.6 and using the mass formula in [21, Lemma 5.1], we see that [21, Theorem 6.4, Corollaries 6.5 and 6.10] hold in any dimension $n \geq 3$, with the same proofs.

Remark 5.9. Recently, Sibony [54] found nonnegative directed currents of bidimension $(1, 1)$ describing the Γ -hull \widehat{K}_Γ of a compact set $K \subset \mathbb{C}^n$ in any directed system determined by a closed, fiberwise conical subset Γ of the tangent bundle $T\mathbb{C}^n$. The hull \widehat{K}_Γ is defined by the maximum principle in terms of Γ -plurisubharmonic functions, i.e., functions whose Levi form is nonnegative in directions from Γ . Sibony's characterization holds even if there are no Γ -directed holomorphic discs (i.e., discs whose derivatives lie in Γ); in particular, his Γ -directed current need not be limits of directed Green currents. The null hull falls within this framework; in this case, the fiber $\Gamma_z \subset T\mathbb{C}^n \cong \mathbb{C}^n$ over any point $z \in \mathbb{C}^n$ is the null quadric (2.12), Γ -plurisubharmonic functions are null plurisubharmonic functions, and Γ -discs are null discs. (The classical case of the polynomial hull is due to Duval and Sibony [22] (see also Wold [57]); in this case $\Gamma = T\mathbb{C}^n$.) It seems an interesting question to decide in which systems directed by a complex analytic variety $\Gamma \subset T\mathbb{C}^n$ with conical fibers is it possible to describe the hull \widehat{K}_Γ by sequences of Γ -directed holomorphic discs $F_j: \overline{\mathbb{D}} \rightarrow \mathbb{C}^n$ whose boundaries converge to K in measure (cf. (5.5)). For the polynomial hull, this holds by Poletsky [49, 50] and Bu-Schachermayer [13]. (For generalizations to complex manifolds, see [40, 41, 52, 53]; for complex spaces, see [20].) For the null hull, this holds by [21, Theorem 6.2] (for $n = 3$) and Corollary 5.5 (for $n > 3$). These seem to be the only cases studied so far.

REFERENCES

- [1] Antonio Alarcón, *Compact complete proper minimal immersions in strictly convex bounded regular domains of \mathbb{R}^3* , XVIII International Fall Workshop on Geometry and Physics, AIP Conf. Proc., vol. 1260, Amer. Inst. Phys., Melville, NY, 2010, pp. 105–111. MR3024549
- [2] A. Alarcón, B. Drinovec Drnovšek, F. Forstnerič, and F. J. López, *Every bordered Riemann surface is a complete conformal minimal surface bounded by Jordan curves*, Proc. Lond. Math. Soc. (3) **111** (2015), no. 4, 851–886. MR3407187
- [3] Antonio Alarcón and Franc Forstnerič, *The Calabi-Yau problem, null curves, and Bryant surfaces*, Math. Ann. **363** (2015), no. 3-4, 913–951. MR3412347
- [4] A. Alarcón, F. Forstnerič, and F. J. López, *New complex analytic methods in the study of nonorientable minimal surfaces in \mathbb{R}^n* . *Mem. Amer. Math. Soc.*, in press.
- [5] Antonio Alarcón, Franc Forstnerič, and Francisco J. López, *Embedded minimal surfaces in \mathbb{R}^n* , Math. Z. **283** (2016), no. 1-2, 1–24. MR3489056
- [6] Antonio Alarcón and Francisco J. López, *Minimal surfaces in \mathbb{R}^3 properly projecting into \mathbb{R}^2* , J. Differential Geom. **90** (2012), no. 3, 351–381. MR2916039
- [7] Antonio Alarcón and Francisco J. López, *Null curves in \mathbb{C}^3 and Calabi-Yau conjectures*, Math. Ann. **355** (2013), no. 2, 429–455. MR3010135
- [8] Antonio Alarcón and Francisco J. López, *Properness of associated minimal surfaces*, Trans. Amer. Math. Soc. **366** (2014), no. 10, 5139–5154. MR3240920
- [9] Antonio Alarcón and Francisco J. López, *Approximation theory for nonorientable minimal surfaces and applications*, Geom. Topol. **19** (2015), no. 2, 1015–1062. MR3336277
- [10] Antonio Alarcón and Nikolai Nadirashvili, *Limit sets for complete minimal immersions*, Math. Z. **258** (2008), no. 1, 107–113. MR2350037
- [11] Michael T. Anderson, *Curvature estimates for minimal surfaces in 3-manifolds*, Ann. Sci. École Norm. Sup. (4) **18** (1985), no. 1, 89–105. MR803196

- [12] Kenneth A. Brakke, *The motion of a surface by its mean curvature*, Mathematical Notes, vol. 20, Princeton University Press, Princeton, N.J., 1978. MR485012
- [13] Shang Quan Bu and Walter Schachermayer, *Approximation of Jensen measures by image measures under holomorphic functions and applications*, Trans. Amer. Math. Soc. **331** (1992), no. 2, 585–608. MR1035999
- [14] Yun Gang Chen, Yoshikazu Giga, and Shun'ichi Goto, *Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations*, J. Differential Geom. **33** (1991), no. 3, 749–786. MR1100211
- [15] Tobias Holck Colding and William P. Minicozzi II, *A course in minimal surfaces*, Graduate Studies in Mathematics, vol. 121, American Mathematical Society, Providence, RI, 2011. MR2780140
- [16] Jean-Pierre Demailly, *Cohomology of q -convex spaces in top degrees*, Math. Z. **204** (1990), no. 2, 283–295. MR1055992
- [17] Avner Dor, *A domain in \mathbf{C}^m not containing any proper image of the unit disc*, Math. Z. **222** (1996), no. 4, 615–625. MR1406270
- [18] Barbara Drinovec Drnovšek and Franc Forstnerič, *Holomorphic curves in complex spaces*, Duke Math. J. **139** (2007), no. 2, 203–253. MR2352132
- [19] Barbara Drinovec Drnovšek and Franc Forstnerič, *Strongly pseudoconvex domains as subvarieties of complex manifolds*, Amer. J. Math. **132** (2010), no. 2, 331–360. MR2654777
- [20] Barbara Drinovec Drnovšek and Franc Forstnerič, *The Poletsky-Rosay theorem on singular complex spaces*, Indiana Univ. Math. J. **61** (2012), no. 4, 1407–1423. MR3085613
- [21] Barbara Drinovec Drnovšek and Franc Forstnerič, *Minimal hulls of compact sets in \mathbb{R}^3* , Trans. Amer. Math. Soc. **368** (2016), no. 10, 7477–7506. MR3471098
- [22] Julien Duval and Nessim Sibony, *Polynomial convexity, rational convexity, and currents*, Duke Math. J. **79** (1995), no. 2, 487–513. MR1344768
- [23] Klaus Ecker and Gerhard Huisken, *Mean curvature evolution of entire graphs*, Ann. of Math. (2) **130** (1989), no. 3, 453–471. MR1025164
- [24] Klaus Ecker and Gerhard Huisken, *Interior estimates for hypersurfaces moving by mean curvature*, Invent. Math. **105** (1991), no. 3, 547–569. MR1117150
- [25] Leonor Ferrer, Francisco Martín, and William H. Meeks III, *Existence of proper minimal surfaces of arbitrary topological type*, Adv. Math. **231** (2012), no. 1, 378–413. MR2935393
- [26] F. Forstnerič, *Stein manifolds and holomorphic mappings*, The homotopy principle in complex analysis. Second edition. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], 56. Springer, Cham, 2017. MR3700709
- [27] Franc Forstnerič and Josip Globevnik, *Discs in pseudoconvex domains*, Comment. Math. Helv. **67** (1992), no. 1, 129–145. MR1144617
- [28] M. Gage and R. S. Hamilton, *The heat equation shrinking convex plane curves*, J. Differential Geom. **23** (1986), no. 1, 69–96. MR840401
- [29] David Gilbarg and Neil S. Trudinger, *Elliptic partial differential equations of second order*, reprint of the 1998 edition, Classics in Mathematics, Springer-Verlag, Berlin, 2001. MR1814364
- [30] R. E. Greene and H. Wu, *Embedding of open Riemannian manifolds by harmonic functions* (English, with French summary), Ann. Inst. Fourier (Grenoble) **25** (1975), no. 1, vii, 215–235. MR0382701
- [31] F. Reese Harvey and H. Blaine Lawson Jr., *An introduction to potential theory in calibrated geometry*, Amer. J. Math. **131** (2009), no. 4, 893–944. MR2543918
- [32] F. Reese Harvey and H. Blaine Lawson Jr., *Plurisubharmonicity in a general geometric context*, Geometry and analysis. No. 1, Adv. Lect. Math. (ALM), vol. 17, Int. Press, Somerville, MA, 2011, pp. 363–402. MR2882430
- [33] F. Reese Harvey and H. Blaine Lawson Jr., *Geometric plurisubharmonicity and convexity: an introduction*, Adv. Math. **230** (2012), no. 4-6, 2428–2456. MR2927376
- [34] F. Reese Harvey and H. Blaine Lawson Jr., *p -convexity, p -plurisubharmonicity and the Levi problem*, Indiana Univ. Math. J. **62** (2013), no. 1, 149–169. MR3158505
- [35] Lars Hörmander, *An introduction to complex analysis in several variables*, 3rd ed., North-Holland Mathematical Library, vol. 7, North-Holland Publishing Co., Amsterdam, 1990. MR1045639

- [36] Gerhard Huisken and Tom Ilmanen, *The inverse mean curvature flow and the Riemannian Penrose inequality*, J. Differential Geom. **59** (2001), no. 3, 353–437. MR1916951
- [37] Gerhard Huisken and Tom Ilmanen, *Higher regularity of the inverse mean curvature flow*, J. Differential Geom. **80** (2008), no. 3, 433–451. MR2472479
- [38] Luquésio P. Jorge and William H. Meeks III, *The topology of complete minimal surfaces of finite total Gaussian curvature*, Topology **22** (1983), no. 2, 203–221. MR683761
- [39] Finnur Lárusson and Ragnar Sigurdsson, *Plurisubharmonic functions and analytic discs on manifolds*, J. Reine Angew. Math. **501** (1998), 1–39. MR1637837
- [40] Finnur Lárusson and Ragnar Sigurdsson, *Plurisubharmonic extremal functions, Lelong numbers and coherent ideal sheaves*, Indiana Univ. Math. J. **48** (1999), no. 4, 1513–1534. MR1757081
- [41] Finnur Lárusson and Ragnar Sigurdsson, *Plurisubharmonicity of envelopes of disc functionals on manifolds*, J. Reine Angew. Math. **555** (2003), 27–38. MR1956593
- [42] Francisco Martín, William H. Meeks III, and Nikolai Nadirashvili, *Bounded domains which are universal for minimal surfaces*, Amer. J. Math. **129** (2007), no. 2, 455–461. MR2306042
- [43] William H. Meeks III and Joaquín Pérez, *Conformal properties in classical minimal surface theory*, Surveys in differential geometry. Vol. IX, Surv. Differ. Geom., vol. 9, Int. Press, Somerville, MA, 2004, pp. 275–335. MR2195411
- [44] William H. Meeks III and Harold Rosenberg, *Maximum principles at infinity*, J. Differential Geom. **79** (2008), no. 1, 141–165. MR2401421
- [45] William W. Meeks III and Shing Tung Yau, *The existence of embedded minimal surfaces and the problem of uniqueness*, Math. Z. **179** (1982), no. 2, 151–168. MR645492
- [46] Gwenaél Mercier and Matteo Novaga, *Mean curvature flow with obstacles: existence, uniqueness and regularity of solutions*, Interfaces Free Bound. **17** (2015), no. 3, 399–426. MR3421913
- [47] Johannes C. C. Nitsche, *Vorlesungen über Minimalflächen* (German), Die Grundlehren der mathematischen Wissenschaften, Band 199, Springer-Verlag, Berlin-New York, 1975. MR0448224
- [48] Robert Osserman, *A survey of minimal surfaces*, 2nd ed., Dover Publications, Inc., New York, 1986. MR852409
- [49] Evgeny A. Poletsky, *Plurisubharmonic functions as solutions of variational problems*, Several complex variables and complex geometry, Part 1 (Santa Cruz, CA, 1989), Proc. Sympos. Pure Math., vol. 52, Amer. Math. Soc., Providence, RI, 1991, pp. 163–171. MR1128523
- [50] Evgeny A. Poletsky, *Holomorphic currents*, Indiana Univ. Math. J. **42** (1993), no. 1, 85–144. MR1218708
- [51] R. Michael Range, *Holomorphic functions and integral representations in several complex variables*, Graduate Texts in Mathematics, vol. 108, Springer-Verlag, New York, 1986. MR847923
- [52] Jean-Pierre Rosay, *Approximation of non-holomorphic maps, and Poletsky theory of discs*, J. Korean Math. Soc. **40** (2003), no. 3, 423–434. MR1973910
- [53] Jean-Pierre Rosay, *Poletsky theory of disks on holomorphic manifolds*, Indiana Univ. Math. J. **52** (2003), no. 1, 157–169. MR1970025
- [54] Nessim Sibony, *Pfaff systems, currents and hulls*, Math. Z. **285** (2017), no. 3-4, 1107–1123. MR3623742
- [55] E. Spadaro, *Mean-convex sets and minimal barriers*, preprint, 2011. arXiv:1112.4288.
- [56] Graham H. Williams, *The Dirichlet problem for the minimal surface equation with Lipschitz continuous boundary data*, J. Reine Angew. Math. **354** (1984), 123–140. MR767575
- [57] Erlend Fornæss Wold, *A note on polynomial convexity: Poletsky disks, Jensen measures and positive currents*, J. Geom. Anal. **21** (2011), no. 2, 252–255. MR2772071

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA E INSTITUTO DE MATEMÁTICAS (IEMATH-GR),
UNIVERSIDAD DE GRANADA, E-18071 GRANADA, SPAIN

Email address: `alarcon@ugr.es`

FACULTY OF MATHEMATICS AND PHYSICS, UNIVERSITY OF LJUBLJANA, AND INSTITUTE OF
MATHEMATICS, PHYSICS AND MECHANICS, JADRANSKA 19, SI-1000 LJUBLJANA, SLOVENIA

Email address: `barbara.drinovec@fmf.uni-lj.si`

FACULTY OF MATHEMATICS AND PHYSICS, UNIVERSITY OF LJUBLJANA, AND INSTITUTE OF
MATHEMATICS, PHYSICS AND MECHANICS, JADRANSKA 19, SI-1000 LJUBLJANA, SLOVENIA

Email address: `franc.forstneric@fmf.uni-lj.si`

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA E INSTITUTO DE MATEMÁTICAS (IEMATH-GR),
UNIVERSIDAD DE GRANADA, E-18071 GRANADA, SPAIN

Email address: `fjlopez@ugr.es`