



The Calabi–Yau problem for Riemann surfaces with finite genus and countably many ends

Antonio Alarcón and Franc Forstnerič

Abstract. In this paper, we show that if R is a compact Riemann surface and $M = R \setminus \bigcup_i D_i$ is a domain in R whose complement is a union of countably many pairwise disjoint smoothly bounded closed discs, D_i , then there is a complete conformal minimal immersion $X: M \rightarrow \mathbb{R}^3$, extending to a continuous map $X: \overline{M} \rightarrow \mathbb{R}^3$, such that $X(bM) = \bigcup_i X(bD_i)$ is a union of pairwise disjoint Jordan curves. In particular, M is the complex structure of a complete bounded minimal surface in \mathbb{R}^3 . This extends a recent result for finite bordered Riemann surfaces.

1. Introduction

A classical problem in the theory of minimal surfaces in Euclidean spaces is the *conformal Calabi–Yau problem*, asking which open Riemann surfaces, M , admit a complete conformal minimal immersion $X: M \rightarrow \mathbb{R}^n$ ($n \geq 3$) with bounded image. (Recall that a continuous map $X: M \rightarrow \mathbb{R}^n$ is said to be complete if the image of any divergent curve in M has infinite Euclidean length. If X is an immersion, this is equivalent to asking that the Riemannian metric $X^*(ds^2)$ on M , induced by the Euclidean metric ds^2 on \mathbb{R}^n via X , is a complete metric.) This problem originates in a conjecture of E. Calabi from 1965 that such immersions do not exist (see Kobayashi and Eells [24], p. 170, and Chern [17], p. 212). Groundbreaking counterexamples to Calabi’s conjecture were given by Jorge and Xavier [23] in 1980 (a complete immersed minimal disc in \mathbb{R}^3 with a bounded coordinate function), Nadirashvili [28] in 1996 (a complete bounded immersed minimal disc in \mathbb{R}^3), and many others. In particular, there are examples in the literature of complete bounded minimal surfaces in \mathbb{R}^3 with any topological type (see Ferrer, Martín, and Meeks [19]). The related *asymptotic Calabi–Yau problem* (see S.-T. Yau [39], p. 360) asks about the asymptotic behaviour of such surfaces near their ends. We refer to the recent papers [3], [8] for the history and literature on these problems.

The first main result of this paper is the following.

Mathematics Subject Classification (2010): 53A10, 53C42; 32B15, 32H02.

Keywords: Riemann surface, minimal surface, Calabi–Yau problem.

Theorem 1.1. *Let R be a compact Riemann surface. If $M = R \setminus \bigcup_{i=0}^{\infty} D_i$ is a domain in R whose complement is a countable union of pairwise disjoint, smoothly bounded closed discs D_i (diffeomorphic images of $\overline{\mathbb{D}} = \{z \in \mathbb{C} : |z| \leq 1\}$), then M is the complex structure of a complete bounded minimal surface in \mathbb{R}^3 .*

More precisely, for any $n \geq 3$ there exists a continuous map $X: \overline{M} \rightarrow \mathbb{R}^n$ such that $X: M \rightarrow \mathbb{R}^n$ is a complete conformal minimal immersion and such that $X(bM) = \bigcup_i X(bD_i)$ is the union of pairwise disjoint Jordan curves. If $n = 4$, then $X: M \rightarrow \mathbb{R}^4$ can be chosen an immersion with simple double points, and if $n \geq 5$, then $X: \overline{M} \rightarrow \mathbb{R}^n$ can be chosen an embedding.

The analogous result holds if R is a nonorientable compact conformal surface.

The discs D_i in the theorem are assumed to have boundaries of class \mathcal{C}^r for some $r > 1$, possibly noninteger. The Jordan curves $X(bD_i) \subset \mathbb{R}^n$ are everywhere nonrectifiable, but we show that X can be chosen such that they have Hausdorff dimension one.

The analogue of Theorem 1.1 when M is the complement of finitely many pairwise disjoint discs in a compact Riemann surface was obtained in [3], Theorem 1.1; this result also follows by a simplification of the proof of Theorem 1.1. Such M is a *bordered Riemann surface* whose boundary, bM , consists of finitely many Jordan curves. The surfaces in Theorem 1.1 still have finite genus, but they may have countably many ends.

Theorem 1.1 is proved in Section 3 by an inductive application of Lemma 4.1 in [3] (see Lemma 2.4) which shows how to increase the intrinsic diameter of a conformal minimal immersion $M \rightarrow \mathbb{R}^n$ from a compact bordered Riemann surface by an arbitrarily big amount, while at the same time keeping the map uniformly close to the given one. Lemma 2.2 provides an estimate of the intrinsic radius of the image surface from below during the inductive process.

The proof of Theorem 1.1 gives several additions. In particular, the complete conformal minimal immersion $X: M \rightarrow \mathbb{R}^n$ can be chosen to have vanishing flux. Alternatively, a minor modification of the proof enables us to prescribe the flux of X on any given finite family of classes in the first homology group $H_1(M, \mathbb{Z})$; however, we do not know whether X can be chosen with arbitrarily prescribed flux map. On the other hand, if we do not insist on controlling the flux of X , then we can choose any conformal minimal immersion $X_0: \overline{R \setminus D_0} \rightarrow \mathbb{R}^n$ ($n \geq 3$) and find for any given number $\epsilon > 0$ a map X as in the theorem which is uniformly ϵ -close to X_0 on \overline{M} .

We wish to emphasize that the class of domains in Theorem 1.1 contains the conformal classes of all Riemann surfaces of finite genus with at most countably many ends, none of which are point ends. Indeed, the uniformization theorem of Z.-X. He and O. Schramm (Theorem 0.2 in [22]) says that every open Riemann surface, M' , with finite genus and at most countably many ends is conformally equivalent to a *circle domain* in a compact Riemann surface R , i.e., a domain of the form

$$(1.1) \quad M = R \setminus \bigcup_i D_i$$

whose complement is the union of at most countably many connected components D_i each of which is either a closed geometric disc or a point. Here, a *geometric disc* in a Riemann surface R is a topological disc whose lifts in the universal cover \tilde{R} of R (which is the disc, the Euclidean plane, or the Riemann sphere) are round discs in \tilde{M} . The ends D_i of M which are points are called *point ends*, while the others are called *disc ends*. An *annular end* is a disc end which does not contain any limit points of other ends. A *puncture end*, or simply a *puncture*, is an end which is conformally isomorphic to the punctured disc; it corresponds to an isolated boundary point of a domain of the form (1.1). The type of an end is independent of a particular representation of a given open Riemann surface as a circle domain, and hence the above notions are well defined for open Riemann surfaces in this class. The He–Schramm theorem includes as a special case open Riemann surfaces of finite topological type (i.e., with finitely generated first homology group $H_1(M, \mathbb{Z})$) and says that every such is conformally equivalent to a domain in a compact Riemann surface whose complement consists of finitely many closed geometric discs and points. (This was known earlier, see e.g. the paper [33] by E. L. Stout.)

In light of these results, Theorem 1.1 gives the following immediate corollary. The second statement follows from the fact that a bounded harmonic function extends harmonically across an isolated point, and hence a bounded complete conformal minimal surface does not have any punctures.

Corollary 1.2. *Every open Riemann surface of finite genus and at most countably ends, none of which are point ends, is the conformal structure of a complete bounded immersed minimal surface in \mathbb{R}^3 , and of a complete bounded embedded minimal surface in \mathbb{R}^5 .*

An open Riemann surface of finite topological type admits a bounded complete conformal minimal immersion into \mathbb{R}^n for some (and hence for any) integer $n \geq 3$ if and only if it has no point ends.

On the other hand, Colding and Minicozzi proved (Corollary 0.13 in [18]) that a complete *embedded* minimal surface of finite topology in \mathbb{R}^3 is necessarily proper in \mathbb{R}^3 , hence unbounded; this was extended to surfaces of finite genus and countably many ends by Meeks, Pérez, and Ros [26], Theorem 1.3. Hence, Corollary 1.2 exposes a major dichotomy between the immersed and the embedded conformal Calabi–Yau problem in dimension 3.

Remark 1.3. We recall the following classical results on the boundary regularity of conformal maps. These show that we are free to precompose conformal minimal immersions $\overline{M} \rightarrow \mathbb{R}^n$ from a compact bordered Riemann surface, $\overline{M} = M \cup bM$, by conformal isomorphisms $M' \rightarrow M$, provided that both bordered Riemann surfaces M and M' have boundaries of class \mathcal{C}^r for some $r > 1$; this does not affect the boundary regularity of the maps. Indeed, any conformal isomorphism $\phi: M \rightarrow M'$ between two such surfaces extends to a homeomorphism $\phi: \overline{M} \rightarrow \overline{M}'$ of their closures by the seminal theorem of Carathéodory [16] from 1913. Furthermore, if the boundaries bM and bM' are smooth (of class \mathcal{C}^∞), then the extension $\phi: \overline{M} \rightarrow \overline{M}'$

is a smooth diffeomorphism by Painlevé's theorem from 1887 [31], [32]. Improvements of these results were made by many authors. In particular, it was shown by Warschawski [34] in 1935 that if bM is of class $\mathcal{C}^{k,\alpha}$ for some $k \in \mathbb{Z}_+ = \{0, 1, 2, \dots\}$ and $0 < \alpha < 1$, then ϕ is of the same class $\mathcal{C}^{k,\alpha}$ on \overline{M} . See Goluzin [21] for more information. For the corresponding boundary regularity results for minimal surfaces, see Nitsche [29], [30]. \square

The situation for Riemann surfaces more general than those in Theorem 1.1 is not understood yet, and we mention the following open problems in this direction.

Problem 1.4. (A) *Let M be a domain of the form $M = R \setminus K$ in a compact Riemann surface R , where K is a nonempty compact subset of R . Assume that M admits a nonconstant bounded harmonic function $h: M \rightarrow \mathbb{R}$ which does not extend to a bounded harmonic function in any bigger domain in R . Does M admit a bounded complete conformal minimal immersion into \mathbb{R}^3 ?*

(B) *Is there an example of a complete bounded minimal surface in \mathbb{R}^3 whose underlying complex structure is $\mathbb{C} \setminus K$, where K is a Cantor set in \mathbb{C} ?*

Recall (see [23], [4], [5], [6]) that every nonconstant bounded harmonic function $h: M \rightarrow (a, b) \subset \mathbb{R}$ on an open Riemann surface M is a component function of a complete conformal minimal immersion $X = (X_1, X_2, h): M \rightarrow \mathbb{R}^3$ whose range is therefore contained in the slab $\{x = (x_1, x_2, x_3) \in \mathbb{R}^3 : a < x_3 < b\}$. Note also that the complement of a compact set $K \subset \mathbb{C}$ of positive capacity admits nonconstant bounded holomorphic (hence harmonic) functions. This shows that the above questions are very natural.

A particular case of problem (A) concerns surfaces with point ends on which disc ends cluster. We wish to thank Antonio Ros for having asked whether anything could be said about the conformal Calabi–Yau problem in this case (private communication on May 17, 2019). This seems a difficult problem, and the answer may depend on how the disc ends approach the set of point ends. In Section 3 we prove the following positive result under the assumption that the compact set of point ends is at infinite distance from the interior of M .

Theorem 1.5. *Let R be a compact Riemann surface, and let M be a domain in R of the form*

$$(1.2) \quad M = R \setminus (D \cup E),$$

where E is a compact set in R and $D = \bigcup_{i=0}^{\infty} D_i$ is the union of a countable family of pairwise disjoint closed geometric discs $D_i \subset R \setminus E$. Fix a point $p_0 \in M$ and set $M_i = R \setminus \bigcup_{j=0}^i D_j$ for every $i \in \mathbb{N}$. If

$$(1.3) \quad \lim_{i \rightarrow \infty} \text{dist}_{M_i}(p_0, E) = +\infty,$$

then there exists a continuous map $X: \overline{M} \rightarrow \mathbb{R}^3$ such that $X|_M: M \rightarrow \mathbb{R}^3$ is a complete conformal minimal immersion and $X|_{bD}: bD = \bigcup_{i=0}^{\infty} bD_i \rightarrow \mathbb{R}^3$ is a topological embedding. In particular, M is the complex structure of a complete bounded minimal surface in \mathbb{R}^3 . The analogous result holds if R is nonorientable.

The distance $\text{dist}_{M_i}(p_0, E)$ is measured with respect to a Riemannian metric d on the ambient surface R . In particular, the set E may consist of point ends of M , and it may even be a Cantor set. Our proof fails for point ends which are at finite distance from an interior point, and it remains an open problem to decide what can happen in such case.

We give an example of a domain in \mathbb{CP}^1 satisfying the requirements in Theorem 1.5 that is inspired by the labyrinth constructed by Jorge and Xavier in [23].

Example 1.6. Let $0 < a < b < 1$ be a pair of numbers and let $\lambda > 0$. Choose finitely many numbers $a < s_0 < s_1 < \dots < s_k < b$. For each $j = 1, \dots, k$, let

$$\delta_j = \frac{s_j - s_{j-1}}{3} > 0$$

and

$$K_j = \{z \in \mathbb{C} : s_{j-1} + \delta_j \leq |z| \leq s_j - \delta_j, \quad |\arg((-1)^j z)| \geq \delta_j\},$$

where $\arg(\cdot)$ is the principal branch of the argument with values in $(-\pi, \pi]$. Up to a slight enlargement of each K_j , we can assume that they are smoothly bounded closed discs, still being pairwise disjoint. Let \mathbb{T} denote the unit circle in \mathbb{C} . Setting $K = \bigcup_{j=1}^k K_j$, it turns out that $\text{dist}_{\mathbb{D} \setminus K}(a\mathbb{T}, b\mathbb{T}) > \lambda$ provided the integer $k \geq 1$ is chosen sufficiently large. Assume that this is so and denote the resulting set K by $K_{a,b,\lambda}$.

Now, choose a decreasing sequence $1 > b_1 > a_1 > b_2 > a_2 > \dots$ with $\lim_{i \rightarrow \infty} a_i = 0$. Set $R = \mathbb{CP}^1$, $D_0 = \mathbb{CP}^1 \setminus \mathbb{D}$, and denote by D_1, D_2, \dots the components of $\bigcup_{j=1}^{\infty} K_{a_j, b_j, 1}$, ordered so that $|z_i| > |z_j|$ for all $z_i \in D_i$ and $z_j \in D_j$ for any pair of indices $i < j$. It is clear that the domain $M = R \setminus (D \cup E)$ with $E = \{0\}$ satisfies condition (1.3) for $M_i = R \setminus \bigcup_{j=0}^i D_j$ and any point $p_0 \in M$. \square

The idea in the previous example can also be used to show the following.

Proposition 1.7. *Let E be a proper compact subset of a compact Riemann surface R . Then there is a sequence of closed, smoothly bounded, pairwise disjoint discs $D_i \subset R \setminus E$ ($i \in \mathbb{Z}_+$) such that $M = R \setminus (\bigcup_{i=0}^{\infty} D_i \cup E)$ is a domain satisfying condition (1.3). Hence, the domain M is the complex structure of a complete bounded minimal surface in \mathbb{R}^3 .*

Proof. Choose a point $p_0 \in R \setminus E$ and a Morse exhaustion function $\rho: R \setminus E \rightarrow \mathbb{R}$ with $\rho(p_0) < 0$. There are sequences $0 < a_1 < b_1 < a_2 < b_2 < \dots$ converging to $+\infty$ such that ρ has no critical values in $[a_j, b_j]$ for every $j \in \mathbb{N}$. It follows that the set $A_j = \{p \in M : a_j \leq \rho(p) \leq b_j\}$ is a union of finitely many pairwise disjoint annuli for each j . By placing sufficiently many closed pairwise disjoint geometric discs in the interior of each connected component of A_j , similarly to what has been done in the above example, we make the length of every path crossing A_j longer than 1. (The length is measured with respect to any given Riemannian metric on R .) Doing this for every $j \in \mathbb{N}$ yields a countable sequence of pairwise disjoint closed discs $D_i \subset R \setminus (E \cup \{p_0\})$ such that the length of any path from p_0 to E which avoids all the discs D_i is infinite and (1.3) holds. \square

The techniques developed in [7], [3], [11] furnish an analogue of Lemma 2.4 for immersed holomorphic curves in \mathbb{C}^n ($n \geq 2$), holomorphic null curves in \mathbb{C}^n for $n \geq 3$, and holomorphic Legendrian curves in \mathbb{C}^{2n+1} for $n \geq 1$. Recall that null curves are holomorphic immersions $Z = (Z_1, \dots, Z_n): M \rightarrow \mathbb{C}^n$ from an open Riemann surface M satisfying the nullity condition $(dZ_1)^2 + (dZ_2)^2 + \dots + (dZ_n)^2 = 0$. Every holomorphic curve in \mathbb{C}^n is also a minimal surface by Wirtinger's theorem [37], while the real and the imaginary part of a holomorphic null curve in \mathbb{C}^n are conformal minimal surfaces in \mathbb{R}^n . By following the proof of Theorem 1.1 and using the analogues of Lemma 2.4 for the appropriate classes of holomorphic curves, one obtains the following result.

Theorem 1.8. *Let M be an open Riemann surface as in Theorem 1.1. Then, for any $n \geq 2$ there exists a continuous map $Z: \overline{M} \rightarrow \mathbb{C}^n$ such that $Z: M \rightarrow \mathbb{C}^n$ is a complete holomorphic immersion and $Z(bM) = \bigcup_i Z(bD_i)$ is the union of pairwise disjoint Jordan curves $Z(bD_i)$. If $n \geq 3$, then Z can be chosen an embedding and such that $Z|_M: M \rightarrow \mathbb{C}^n$ is a complete holomorphic null embedding, or (if n is odd) a complete holomorphic Legendrian embedding.*

The analogue of Theorem 1.5 also holds for these classes of maps, and the questions in Problem 1.4 make sense for holomorphic (null, Legendrian) curves.

Theorem 1.8 contributes to the body of results concerning Yang's problem [38] from 1977, asking about the existence and boundary behaviour of bounded complete complex submanifolds of complex Euclidean spaces. For recent developments on this subject, see the papers [2], [9], [13], [14], [15], [20], and the authors' survey [8].

2. Preliminaries

Given a compact smooth manifold M with nonempty boundary bM and an interior point $p_0 \in \overset{\circ}{M} = M \setminus bM$, we denote by $\mathcal{C} = \mathcal{C}(M, p_0)$ the set of paths $\gamma: [0, 1] \rightarrow M$ with $\gamma(0) = p_0$ and $\gamma(1) \in bM$. (The word *path* always stands for a continuous path. In fact, we shall mainly use piecewise \mathcal{C}^1 paths.) Given a continuous map $X: M \rightarrow \mathbb{R}^n$, we define

$$\text{dist}_X(p_0, bM) = \inf \{ \text{length}(X \circ \gamma) : \gamma \in \mathcal{C} \} \in [0, +\infty],$$

where $\text{length}(\lambda)$ denotes the Euclidean length of a path $\lambda: [0, 1] \rightarrow \mathbb{R}^n$, i.e., the supremum of the sums $\sum_{i=1}^m |\lambda(t_i) - \lambda(t_{i-1})|$ over all subdivisions $0 = t_0 < t_1 < \dots < t_m = 1$ of the interval $[0, 1]$. The same definition applies to a compact domain M with countably many boundary components. The number $\text{dist}_X(p_0, bM)$ is called the *intrinsic diameter* of M with respect to the map X and the point p_0 . A change of the base point changes the intrinsic diameter by a constant.

Assume now that $X: M \rightarrow \mathbb{R}^n$ is a smooth immersion and let $g = X^*ds^2$ be the induced Riemannian metric on M . Given a piecewise \mathcal{C}^1 path $\gamma: [0, 1] \rightarrow M$, we have that

$$\text{length}(X \circ \gamma) = \text{length}_g(\gamma) = \int_0^1 \|\dot{\gamma}(t)\|_g dt.$$

It is well known and easily seen that $\text{dist}_X(p_0, bM) = \text{dist}_g(p_0, bM)$ is the infimum of the lengths of piecewise \mathcal{C}^1 paths in the family $\mathcal{C}(M, p_0)$. Indeed, we can replace any path γ in M by a piecewise smooth path which is not longer than γ by taking a suitably fine subdivision $0 = t_0 < t_1 < \dots < t_m = 1$ of $[0, 1]$ and replacing each segment $C_i = \{\gamma(t) : t_{i-1} \leq t \leq t_i\}$ by the geodesic arc connecting the points $\gamma(t_{i-1})$ and $\gamma(t_i)$.

We shall consider two bigger classes $\mathcal{C}_d = \mathcal{C}_d(M, p_0) \subset \mathcal{C}_{qd} = \mathcal{C}_{qd}(M, p_0)$ of piecewise \mathcal{C}^1 paths in M with the given initial point $p_0 \in \overset{\circ}{M}$. The first one, $\mathcal{C}_d(M, p_0)$, consists of *divergent paths* $\gamma: [0, 1] \rightarrow M$ with $\gamma(0) = p_0$, i.e., such that $\gamma(t)$ leaves any compact subset of $\overset{\circ}{M}$ as t approaches 1. (However, the limit of $\gamma(t)$ as $t \rightarrow 1$ need not exist.) The class $\mathcal{C}_{qd}(M, p_0)$ consists of paths $\gamma: [0, 1] \rightarrow M$ such that $\gamma(0) = p_0$ and γ has a cluster point on bM , i.e., there is a sequence

$$(2.1) \quad 0 < t_1 < t_2 < \dots < 1 \quad \text{with} \quad \lim_{j \rightarrow \infty} t_j = 1 \quad \text{and} \quad \lim_{j \rightarrow \infty} \gamma(t_j) = p \in bM.$$

We call such path *quasidivergent*.

The following lemma shows that we get the same intrinsic diameter by using paths in the bigger family \mathcal{C}_{qd} , and hence also by using paths in \mathcal{C}_d .

Lemma 2.1. *Let g be a Riemannian metric on a compact \mathcal{C}^1 manifold M with boundary bM , and let $p_0 \in \overset{\circ}{M}$. For every path $\gamma \in \mathcal{C}_{qd}(M, p_0)$ we have that $\text{length}_g(\gamma) \geq \text{dist}_g(p_0, bM)$.*

Proof. Fix $\epsilon > 0$. Let $U \subset M$ be a neighbourhood of bM such that every point $q \in U$ can be connected to a point $p \in bM$ by an arc of length less than ϵ . Given $\gamma \in \mathcal{C}_{qd}(M, p_0)$, there is $t_0 \in [0, 1)$ such that $\gamma(t_0) \in U$. Choose an arc $C \subset U$ with $\text{length}_g(C) < \epsilon$ connecting $\gamma(t_0)$ to a point $p \in bM$. Let $\lambda: [0, 1] \rightarrow M$ be a path such that $\lambda(t) = \gamma(t)$ for $t \in [0, t_0]$, and $\lambda(t)$ for $t \in [t_0, 1]$ is a parametrization of C with $\lambda(1) = p$. Then,

$$\text{dist}_g(p_0, bM) \leq \text{length}_g(\lambda) < \text{length}_g(\gamma) + \epsilon.$$

Letting $\epsilon \rightarrow 0$ we obtain $\text{length}_g(\gamma) \geq \text{dist}_g(p_0, bM)$. □

The following lemma will enable us to control the intrinsic radius of a conformal minimal immersion from below in the proofs of Theorems 1.1 and 1.5.

Lemma 2.2. *Let M be a compact connected \mathcal{C}^1 manifold with boundary $bM \neq \emptyset$, and let $X: M \rightarrow \mathbb{R}^n$ be a \mathcal{C}^1 immersion. Given a point $p_0 \in \overset{\circ}{M}$ and a number $\eta > 0$, there exists a number $\epsilon > 0$ such that, for every continuous map $Y: M \rightarrow \mathbb{R}^n$ with $\|X - Y\|_{\mathcal{C}^0(M)} := \max\{|X(p) - Y(p)| : p \in M\} < \epsilon$, we have that*

$$\inf\{\text{length}(Y \circ \gamma) : \gamma \in \mathcal{C}_{qd}(M, p_0)\} \geq \text{dist}_X(p_0, bM) - \eta.$$

Proof. This obviously holds if Y is uniformly \mathcal{C}^1 -close to X on $\overset{\circ}{M}$ since small \mathcal{C}^1 perturbations only change lengths of curves by a small amount. Furthermore, any \mathcal{C}^1 structure on a manifold is equivalent to a \mathcal{C}^∞ structure by a theorem of H. Whitney (Lemma 24 in [36]). Hence, we may assume that M is a compact domain with \mathcal{C}^1 boundary in a smooth manifold $\overset{\circ}{M}$ and X is a smooth immersion $X: \overset{\circ}{M} \rightarrow \mathbb{R}^n$; the latter statement uses an approximation theorem of Whitney [35].

Let $N \rightarrow \widetilde{M}$ denote the normal bundle of the immersion $X: \widetilde{M} \rightarrow \mathbb{R}^n$, so $\dim N = n$. We identify \widetilde{M} with the zero section of N . By the tubular neighbourhood theorem, X extends to a smooth immersion $F: N \rightarrow \mathbb{R}^n$ which agrees with X on the zero section \widetilde{M} of N . Then, $g = F^*(ds^2)$ is a smooth Riemannian metric on N whose restriction to \widetilde{M} is the metric $X^*(ds^2)$, and the map $F: (N, g) \rightarrow (\mathbb{R}^n, ds^2)$ is a local isometry. Let dist_g denote the distance function on N induced by the Riemannian metric g .

We claim that there is a neighbourhood $U \subset N$ of M such that

$$(2.2) \quad \text{dist}_{g,U}(p_0, bM) > \text{dist}_{g,M}(p_0, bM) - \eta/2 = \text{dist}_X(p_0, bM) - \eta/2,$$

where $\text{dist}_{g,U}(p_0, bM)$ is the distance from p_0 to bM over all paths in U and $\eta > 0$ is as in the lemma. Here is an elementary proof. After shrinking N and \widetilde{M} around M if necessary, there is a smooth retraction $\rho: N \rightarrow \widetilde{M}$ such that the kernel $\ker(d\rho_x)$ of its differential at any point $x \in \widetilde{M}$ is the g -orthogonal complement of $T_x \widetilde{M}$ in $T_x N$. Since $d\rho_x$ equals the identity on $T_x \widetilde{M}$, it follows that $d\rho_x$ has g -norm 1. Hence, for any $r > 1$ there is a neighbourhood $U \subset N$ of M such that $d\rho_x: T_x N \rightarrow T_{\rho(x)} N$ has g -norm less than r for every point $x \in U$. For every path $\gamma: [0, 1] \rightarrow U$ we then have $\text{length}_g(\rho \circ \gamma) \leq r \cdot \text{length}_g(\gamma)$. Choosing γ to be a path in U connecting $\gamma(0) = p_0$ to a point $\gamma(1) \in bM$, we obtain

$$\text{dist}_{g,M}(p_0, bM) \leq \text{length}_g(\rho \circ \gamma) \leq r \cdot \text{length}_g(\gamma).$$

(The first inequality holds even if $\rho \circ \gamma$ is not contained in M since it then crosses bM at some time $t_0 \in (0, 1)$, and the length of this shorter path is still $\geq \text{dist}_{g,M}(p_0, bM)$.) Taking the infimum over all such paths γ gives

$$\text{dist}_{g,M}(p_0, bM) \leq r \cdot \text{dist}_{g,U}(p_0, bM).$$

If r is chosen close enough to 1, then (2.2) holds, thereby proving the claim.

Since $F: N \rightarrow F(N) \subset \mathbb{R}^n$ is a local isometry and M is compact, there is a number $\epsilon_0 > 0$ such that for every point $p \in M$, the closed ball

$$(2.3) \quad B_g(p, \epsilon_0) := \{q \in N : \text{dist}_g(p, q) \leq \epsilon_0\}$$

is contained in U and F maps $B_g(p, \epsilon_0)$ isometrically onto the closed Euclidean ball $\overline{\mathbb{B}}(X(p), \epsilon_0) \subset \mathbb{R}^n$. By decreasing ϵ_0 if necessary, we may assume that $0 < \epsilon_0 < \eta/2$.

Given a continuous map $Y: M \rightarrow \mathbb{R}^n$ satisfying

$$\max_{p \in M} |Y(p) - X(p)| < \epsilon \leq \epsilon_0,$$

the above implies that there is a unique continuous map $\widetilde{Y}: M \rightarrow U$ such that

$$(2.4) \quad Y = F \circ \widetilde{Y} \quad \text{and} \quad \text{dist}_g(p, \widetilde{Y}(p)) = |X(p) - Y(p)| < \epsilon \quad \text{for all } p \in M.$$

Let $\gamma \in \mathcal{C}_{cd}$ be a quasidivergent path in M with $\gamma(0) = p_0$. Fix a number ϵ with $0 < \epsilon < \epsilon_0/2$. There is a boundary point $p \in bM$ and $t_0 \in (0, 1)$ such that

$$(2.5) \quad \text{dist}_g(\gamma(t_0), p) < \epsilon.$$

By (2.4) we have that $Y \circ \gamma = F \circ \tilde{\gamma}$, where the path $\tilde{\gamma} = \tilde{Y} \circ \gamma: [0, 1] \rightarrow U$ satisfies

$$(2.6) \quad \text{dist}_g(\gamma(t), \tilde{\gamma}(t)) < \epsilon \quad \text{for all } t \in [0, 1].$$

Since $F: (N, g) \rightarrow (\mathbb{R}^n, ds^2)$ is a local isometry, we also have that

$$(2.7) \quad \text{length}_g(\tilde{\gamma}) = \text{length}(F \circ \tilde{\gamma}) = \text{length}(Y \circ \gamma).$$

By (2.5), (2.6), and the triangle inequality, the point $\tilde{\gamma}(t_0) = \tilde{Y}(\gamma(t_0))$ satisfies

$$\text{dist}_g(\tilde{\gamma}(t_0), p) \leq \text{dist}_g(\tilde{\gamma}(t_0), \gamma(t_0)) + \text{dist}_g(\gamma(t_0), p) < \epsilon + \epsilon \leq \epsilon_0 < \eta/2.$$

By adding to the path $\tilde{\gamma}: [0, t_0] \rightarrow N$ an arc in the ball $B_g(p, \epsilon)$ (2.3) of g -length $< \eta/2$ connecting the point $\tilde{\gamma}(t_0)$ to $p \in bM$, we obtain a path $\lambda: [0, 1] \rightarrow U$ connecting $\lambda(0) = p_0$ to $\lambda(1) = p \in bM$ such that

$$\text{length}_g(\lambda) < \text{length}_g(\tilde{\gamma}) + \eta/2.$$

We obviously have $\text{length}_g(\lambda) \geq \text{dist}_{g,U}(p_0, bM)$. Together with (2.2) we obtain

$$\text{length}_g(\tilde{\gamma}) > \text{length}_g(\lambda) - \eta/2 \geq \text{dist}_{g,U}(p_0, bM) - \eta/2 > \text{dist}_X(p_0, bM) - \eta.$$

In view of (2.7), it follows that $\text{length}(Y \circ \gamma) > \text{dist}_X(p_0, bM) - \eta$. \square

The proof of Lemma 2.2 also applies to the distance from an interior point $p_0 \in \overset{\circ}{M}$ to any given nonempty compact subset E of M . The following result to this effect will be used in the proof of Theorem 1.5.

Lemma 2.3. *Let M be a compact connected \mathcal{C}^1 manifold (either closed or with boundary), and let $X: M \rightarrow \mathbb{R}^n$ be a \mathcal{C}^1 immersion. Given a nonempty compact set $E \subset M$, a point $p_0 \in M \setminus E$, and a number $\eta > 0$, there is a number $\epsilon > 0$ such that, for every continuous map $Y: M \rightarrow \mathbb{R}^n$ with $\|X - Y\|_{\mathcal{C}^0(M)} < \epsilon$ and for every path $\gamma: [0, 1] \rightarrow M \setminus E$ with $\gamma(0) = p_0$ such that $\gamma(t)$ has a limit point in E as $t \rightarrow 1$, we have $\text{length}(Y \circ \gamma) \geq \text{dist}_X(p_0, E) - \eta$.*

The following lemma (see Lemma 4.1 in [3]) is the main ingredient in the proof of Theorem 1.1. It enables one to make the intrinsic diameter of an immersed conformal minimal surface arbitrarily big by a \mathcal{C}^0 small deformation.

Lemma 2.4. *Let M be a compact bordered Riemann surface, and let $X: M \rightarrow \mathbb{R}^n$ ($n \geq 3$) be a conformal minimal immersion of class $\mathcal{C}^1(M)$. Given a point $p_0 \in M = M \setminus bM$, an integer $d \in \mathbb{Z}_+$, and numbers $\epsilon > 0$ (small) and $\mu > 0$ (big), there is a continuous map $Y: M \rightarrow \mathbb{R}^n$ whose restriction to $\overset{\circ}{M}$ is a conformal minimal immersion such that the following conditions hold.*

- (i) $|Y(p) - X(p)| < \epsilon$ for all $p \in M$.
- (ii) $\text{dist}_Y(p_0, bM) > \mu$.
- (iii) $Y|_{bM}: bM \rightarrow \mathbb{R}^n$ is injective.
- (iv) $\text{Flux}_Y = \text{Flux}_X$.

Condition (iii) follows from a general position theorem; see Theorem 4.5 in [3]. The analogous result holds if M is a nonorientable compact bordered conformal surface; see Section 6.3 in [12], and in particular Lemma 6.7 in the cited paper.

3. Proof of Theorems 1.1 and 1.5

Proof of Theorem 1.1. Assume that R is a compact Riemann surface and M is a domain in R of the form

$$(3.1) \quad M = R \setminus \bigcup_{i=0}^{\infty} D_i,$$

where $\{D_i\}_{i \in \mathbb{Z}_+}$ is a countable family of closed, pairwise disjoint, smoothly bounded discs in R . We shall construct a continuous map $X: \overline{M} \rightarrow \mathbb{R}^n$ satisfying the conclusion of the theorem and such that the Jordan curves $X(bD_i)$, $i \in \mathbb{Z}_+$, have Hausdorff dimension one. Moreover, we shall ensure that the complete conformal minimal immersion $X: M \rightarrow \mathbb{R}^n$ has vanishing flux.

For every $i = 0, 1, 2, \dots$, we let

$$(3.2) \quad M_i = R \setminus \bigcup_{k=0}^i \mathring{D}_k.$$

This is a compact bordered Riemann surface with boundary $bM_i = \bigcup_{k=0}^i bD_k$, and

$$M_0 \supset M_1 \supset M_2 \supset \dots \supset \bigcap_{i=1}^{\infty} M_i = \overline{M}.$$

By Theorem 4.5 (a) in [3], there exists a conformal minimal immersion $X_0: M_0 \rightarrow \mathbb{R}^n$ of class $\mathcal{C}^1(M_0)$ with vanishing flux such that $X_0|_{bM_0}: bM_0 \rightarrow \mathbb{R}^n$ is injective. Choose a Riemannian distance function d on R , a point $p_0 \in M$, and a pair of numbers $\epsilon_0 > 0$ and $\tau_0 \in \mathbb{N} = \{1, 2, 3, \dots\}$. An inductive application of Lemmas 2.2 and 2.4 furnishes a sequence of conformal minimal immersions $X_i: M_i \rightarrow \mathbb{R}^n$ of class $\mathcal{C}^1(M_i)$, numbers $\epsilon_i > 0$, and integers $\tau_i > i$ satisfying the following conditions for every $i \in \mathbb{N}$.

- (a_i) $\text{dist}_{X_i}(p_0, bM_i) > i$.
- (b_i) $X_i: bM_i \rightarrow \mathbb{R}^n$ is injective.
- (c_i) $\sup_{p \in M_i} |X_i(p) - X_{i-1}(p)| < \epsilon_{i-1}$.
- (d_i) For every continuous map $Y: M_i \rightarrow \mathbb{R}^n$ with $\|Y - X_i\|_{\mathcal{C}(M_i)} < 2\epsilon_i$ we have

$$\inf\{\text{length}(Y \circ \gamma) : \gamma \in \mathcal{C}_{qd}(M_i, p_0)\} > \text{dist}_{X_i}(p_0, bM_i) - 1 > i - 1.$$

- (e_i) We have $0 < \epsilon_i < \frac{1}{2} \min\{\epsilon_{i-1}, \delta_i, \tau_i^{-i}\}$, where

$$\delta_i := \frac{1}{i^2} \inf \left\{ |X_i(p) - X_i(q)| : p, q \in bM_i, d(p, q) > \frac{1}{i} \right\} > 0.$$

- (f_i) We have $\tau_i > \tau_{i-1}$, and for each $k \in \{0, \dots, i\}$ there is a set $A_{i,k} \subset X_i(bD_k)$ consisting of τ_i^{i+1} points such that

$$\max\{\text{dist}(p, A_{i,k}) : p \in X_i(bD_k)\} < 1/\tau_i^i,$$

where $\text{dist}(p, A_{i,k}) = \min\{|p - q| : q \in A_{i,k}\}$ is the Euclidean distance in \mathbb{R}^n .

- (g_i) X_i has vanishing flux.

Let us explain the induction step. Assume that for some $i \in \mathbb{N}$ we have maps X_0, \dots, X_{i-1} and numbers $\epsilon_0, \dots, \epsilon_{i-1}$ and $\tau_0, \dots, \tau_{i-1}$ satisfying these conditions for the respective values of the index. (This holds for $i = 1$ by using X_0, ϵ_0 , and τ_0 , and the above conditions are void except for (a₀), (b₀), and (g₀); the second part of (f₀) also holds true if we choose $\tau_0 \in \mathbb{N}$ sufficiently large.) Lemma 2.4 applied to $X_{i-1}|_{M_i}$ furnishes a conformal minimal immersion $X_i: M_i \rightarrow \mathbb{R}^n$ satisfying (a_i), (b_i), (c_i), and (g_i); note that $X_{i-1}|_{M_i}$ is flux vanishing since so is X_{i-1} by (g_{i-1}). Pick $\tau_i \in \mathbb{N}$ so large that condition (f_i) is satisfied; it suffices to choose

$$\tau_i > \tau_{i-1} + \sum_{k=0}^i \text{length}(X_i(bD_k)).$$

Pick a number $\epsilon_i > 0$ satisfying condition (e_i); such exists since $X_i|_{bM_i}$ is injective by (b_i). Finally, decreasing $\epsilon_i > 0$ if necessary we may assume that condition (d_i) holds as well in view of Lemma 2.2. The induction may proceed.

Conditions (c_i) and (e_i) imply that the sequence X_i converges uniformly on \overline{M} , where M was given in (3.1), to a continuous map $X = \lim_{i \rightarrow \infty}: \overline{M} \rightarrow \mathbb{R}^n$ whose restriction to M is a conformal minimal immersion $X: M \rightarrow \mathbb{R}^n$, provided that each $\epsilon_i > 0$ is chosen sufficiently small. More precisely, for every $p \in \overline{M}$ we have

$$(3.3) \quad |X(p) - X_i(p)| \leq \sum_{k=i}^{\infty} |X_{k+1}(p) - X_k(p)| < \sum_{k=i}^{\infty} \epsilon_k < 2\epsilon_i.$$

We can extend X from \overline{M} to a continuous map $X: M_i \rightarrow \mathbb{R}^n$ such that the above inequality holds for all $p \in M_i$.

Conditions (a_i) and $0 < \epsilon_i < \epsilon_{i-1}/2$ (see (e_i)) ensure that $X: M \rightarrow \mathbb{R}^n$ is complete. Indeed, consider any divergent path $\gamma: [0, 1) \rightarrow M$ with $\gamma(0) = p_0$. There is an increasing sequence $0 < t_1 < t_2 < \dots < 1$ with $\lim_{j \rightarrow \infty} t_j = 1$ such that $\lim_{j \rightarrow \infty} \gamma(t_j) = p \in bM$ (cf. (2.1)). Then, $p \in bD_{i_0}$ for some $i_0 \in \mathbb{Z}_+$, and hence $p \in bM_i$ for all $i \geq i_0$. It follows that γ is a quasidivergent path in the bordered Riemann surface M_i for any $i \geq i_0$. (See Section 2 for this notion.) Conditions (a_i), (d_i), and (3.3) imply for any $i \geq i_0$ that

$$\text{length}(X(\gamma)) > \text{dist}_{X_i}(p_0, bM_i) - 1 > i - 1.$$

Letting $i \rightarrow +\infty$ shows that $\text{length}(X(\gamma)) = +\infty$.

Conditions (b_i), (c_i), and (e_i) imply that the limit map $X: \overline{M} \rightarrow \mathbb{R}^n$ is injective on $bM = \bigcup_{i \in \mathbb{Z}_+} bD_i$ (see the proof of Theorem 1.1 in [3] for the details), whereas (g_i) ensures that X has vanishing flux.

Finally, in order to see that all Jordan curves $X(bD_k)$ ($k \in \mathbb{Z}_+$) have Hausdorff dimension one, pick such a k . By (3.3), (e_i), and (f_i), we have for each $i > k$ that

$$(3.4) \quad \max\{\text{dist}(p, A_{i,k}): p \in X(bD_k)\} < 2/\tau_i^i.$$

Since $A_{i,k}$ consists of precisely τ_i^{i+1} points and $\tau_i^{i+1}(2/\tau_i^i)^{1+1/i} = 2^{1+1/i} \leq 4$ for every integer $i > k$, (3.4) implies that the Hausdorff measure $\mathcal{H}^1(X(bD_k))$ is finite, and hence the Hausdorff dimension of $X(bD_k)$ is at most one (cf. Lemma 2.2 in [25] or Section 4.1 in [1]; see [27] for an introduction to the Hausdorff measure). On the other hand, $X(bD_k)$ is homeomorphic to the circle $\mathbb{T} = \{z \in \mathbb{C}: |z| = 1\}$ and hence its Hausdorff dimension is at least one, so it is one.

Furthermore, by using the general position argument for minimal surfaces at every step of the proof (see Theorem 4.1 in [10]), we can ensure that the limit map $X: M \rightarrow \mathbb{R}^n$ is an immersion with simple double points if $n = 4$, and is an embedding if $n \geq 5$. See the proof of Theorem 1.1 in [3] for the details.

This completes the proof of Theorem 1.1. The same proof applies in the nonorientable case if we replace Lemma 2.4 by Lemma 6.7 in [12]. \square

Proof of Theorem 1.5. For every $i = 0, 1, 2, \dots$, let $M_i = R \setminus \bigcup_{k=0}^i \mathring{D}_k$ be the compact domain (3.2) in R . Choose a conformal minimal immersion $X_0: M_0 \rightarrow \mathbb{R}^3$. Then, the given metric on R is comparable on M_0 to the metric $g_0 = (X_0)^*(ds^2)$ induced by X_0 , and hence condition (1.3) holds for the latter metric as well. We shall use the same argument at every step when changing the metric.

By (1.3), there is $i_1 \in \mathbb{N}$ such that $\text{dist}_{M_{i_1}, g_0}(p_0, E) > 1$. Choose a conformal minimal immersion $X_1: M_{i_1} \rightarrow \mathbb{R}^3$ which approximates X_0 uniformly on M_{i_1} and satisfies the conditions

$$\text{dist}_{M_{i_1}, X_1}(p_0, bM_{i_1}) > 1 \quad \text{and} \quad \text{dist}_{M_{i_1}, X_1}(p_0, E) > 1.$$

The first condition is achieved by Lemma 2.4, while the second holds by Lemma 2.3 provided the approximation is close enough. The metric $g_1 = (X_1)^*(ds^2)$ is comparable on M_{i_1} with g_0 . Hence, by (1.3) there is an integer $i_2 > i_1$ such that $\text{dist}_{M_{i_2}, g_1}(p_0, E) > 2$. The same argument as before gives a conformal minimal immersion $X_2: M_{i_2} \rightarrow \mathbb{R}^3$ approximating X_1 uniformly on M_{i_2} and satisfying

$$\text{dist}_{M_{i_2}, X_2}(p_0, bD_{i_2}) > 2 \quad \text{and} \quad \text{dist}_{M_{i_2}, X_2}(p_0, E) > 2.$$

Continuing inductively we get sequences of integers $i_1 < i_2 < \dots$ and conformal minimal immersions $X_k: M_{i_k} \rightarrow \mathbb{R}^3$ ($k \in \mathbb{N}$) satisfying

$$(3.5) \quad \text{dist}_{M_{i_k}, X_k}(p_0, bM_{i_k}) > k \quad \text{and} \quad \text{dist}_{M_{i_k}, X_k}(p_0, E) > k.$$

Assuming as we may that X_k approximates X_{k-1} sufficiently closely uniformly on M_{i_k} for every $k \in \mathbb{N}$, we can ensure as in the proof of Theorem 1.1 that the sequence X_k converges uniformly on the compact set $M' = \bigcap_{i=0}^{\infty} \overline{M}_i$ to a continuous limit map $X: M' \rightarrow \mathbb{R}^3$ whose restriction to the interior of M' is a complete conformal minimal immersion, and such that $X|_M: M \rightarrow \mathbb{R}^n$ satisfies the conclusion of the theorem. (Here, $M = M' \setminus E$ is given by (1.2).) In particular, the distance from any point of M to the boundary $bM = \bigcup_i bC_i \cup bE$ in the metric $X^*(ds^2)$ is infinite by (3.5) and Lemma 2.3. \square

References

- [1] ALARCÓN, A.: Compact complete minimal immersions in \mathbb{R}^3 . *Trans. Amer. Math. Soc.* **362** (2010), no. 8, 4063–4076.
- [2] ALARCÓN, A.: Complete complex hypersurfaces in the ball come in foliations. To appear in *J. Differential Geom.*
- [3] ALARCÓN, A., DRNOVEC DRNOVŠEK, B., FORSTNERIČ, F. AND LÓPEZ, F. J.: Every bordered Riemann surface is a complete conformal minimal surface bounded by Jordan curves. *Proc. Lond. Math. Soc.* (3) **111** (2015), no. 4, 851–886.

- [4] ALARCÓN, A. AND FERNÁNDEZ, I.: Complete minimal surfaces in \mathbb{R}^3 with a prescribed coordinate function. *Differential Geom. Appl.* **29** (suppl. 1) (2011), S9–S15.
- [5] ALARCÓN, A., FERNÁNDEZ, I. AND LÓPEZ, F. J.: Complete minimal surfaces and harmonic functions. *Comment. Math. Helv.* **87** (2012), no. 4, 891–904.
- [6] ALARCÓN, A., FERNÁNDEZ, I. AND LÓPEZ, F. J.: Harmonic mappings and conformal minimal immersions of Riemann surfaces into \mathbb{R}^N . *Calc. Var. Partial Differential Equations* **47** (2013), no. 1-2, 227–242.
- [7] ALARCÓN, A. AND FORSTNERIČ, F.: The Calabi–Yau problem, null curves, and Bryant surfaces. *Math. Ann.* **363** (2015), no. 3-4, 913–951.
- [8] ALARCÓN, A. AND FORSTNERIČ, F.: New complex analytic methods in the theory of minimal surfaces: a survey. *J. Aust. Math. Soc.* **106** (2019), no. 3, 287–341.
- [9] ALARCÓN, A. AND FORSTNERIČ, F.: A foliation of the ball by complete holomorphic discs. *Math. Z.* **296** (2020), no. 1-2, 169–174.
- [10] ALARCÓN, A., FORSTNERIČ, F. AND LÓPEZ, F. J.: Embedded minimal surfaces in \mathbb{R}^n . *Math. Z.* **283** (2016), no. 1-2, 1–24.
- [11] ALARCÓN, A., FORSTNERIČ, F. AND LÓPEZ, F. J.: Holomorphic Legendrian curves. *Compos. Math.* **153** (2017), no. 9, 1945–1986.
- [12] ALARCÓN, A., FORSTNERIČ, F. AND LÓPEZ, F. J.: New complex analytic methods in the study of non-orientable minimal surfaces in \mathbb{R}^n . *Mem. Amer. Math. Soc.* **264** (2020), no. 1283, vi+77 pp.
- [13] ALARCÓN, A. AND GLOBEVIK, J.: Complete embedded complex curves in the ball of \mathbb{C}^2 can have any topology. *Anal. PDE* **10** (2017), no. 8, 1987–1999.
- [14] ALARCÓN, A., GLOBEVIK, J. AND LÓPEZ, F. J.: A construction of complete complex hypersurfaces in the ball with control on the topology. *J. Reine Angew. Math.* **751** (2019), 289–308.
- [15] ALARCÓN, A. AND LÓPEZ, F. J.: Complete bounded embedded complex curves in \mathbb{C}^2 . *J. Eur. Math. Soc. (JEMS)* **18** (2016), no. 8, 1675–1705.
- [16] CARATHÉODORY, C.: Über die Begrenzung einfach zusammenhängender Gebiete. *Math. Ann.* **73** (1913), 323–370.
- [17] CHERN, S. S.: The geometry of G -structures. *Bull. Amer. Math. Soc.* **72** (1966), 167–219.
- [18] COLDING, T. H. AND MINICOZZI II, W. P.: The Calabi–Yau conjectures for embedded surfaces. *Ann. of Math.* (2) **167** (2008), no. 1, 211–243.
- [19] FERRER, L., MARTÍN, F. AND MEEKS III, W. H.: Existence of proper minimal surfaces of arbitrary topological type. *Adv. Math.* **231** (2012), no. 1, 378–413.
- [20] GLOBEVIK, J.: A complete complex hypersurface in the ball of \mathbb{C}^N . *Ann. of Math.* (2) **182** (2015), no. 3, 1067–1091.
- [21] GOLUZIN, G. M.: *Geometric theory of functions of a complex variable*. Translations of Mathematical Monographs 26, Amer. Math. Soc., Providence, RI, 1969.
- [22] HE, Z.-X. AND SCHRAMM, O.: Fixed points, Koebe uniformization and circle packings. *Ann. of Math.* (2) **137** (1993), no. 2, 369–406.
- [23] JORGE, L. P. DE M. AND XAVIER, F.: A complete minimal surface in \mathbb{R}^3 between two parallel planes. *Ann. of Math.* (2) **112** (1980), no. 1, 203–206.
- [24] KOBAYASHI, S. AND EELLS JR., J.: *Proceedings of the United States-Japan Seminar in Differential Geometry, Kyoto, Japan, 1965*. Nippon Hyoronsha Co., Tokyo, 1966.

- [25] MARTÍN, F. AND NADIRASHVILI, N.: A Jordan curve spanned by a complete minimal surface. *Arch. Ration. Mech. Anal.* **184** (2007), no. 2, 285–301.
- [26] MEEKS III, W. H., PÉREZ, J. AND ROS, A.: The embedded Calabi–Yau conjecture for finite genus. Preprint, arXiv: 1806.03104, June 2018.
- [27] MORGAN, F.: *Geometric measure theory. A beginner’s guide.* Foruth edition. Elsevier/Academic Press, Amsterdam, 2009.
- [28] NADIRASHVILI, N.: Hadamard’s and Calabi–Yau’s conjectures on negatively curved and minimal surfaces. *Invent. Math.* **126** (1996), no. 3, 457–465.
- [29] NITSCHE, J. C. C.: The boundary behavior of minimal surfaces. Kellogg’s theorem and Branch points on the boundary. *Invent. Math.* **8** (1969), 313–333.
- [30] NITSCHE, J. C. C.: Concerning my paper on the boundary behavior of minimal surfaces. *Invent. Math.* **9** (1969/1970), 270.
- [31] PAINLEVÉ, P.: *Thèse d’Analyse. Sur les lignes singulières des fonctions analytiques.* Paris. 136 S. 4°, 1887.
- [32] PAINLEVÉ, P.: Sur la théorie de la représentation conforme. *C.R. Acad. Sci., Paris* **112** (1891), 653–657.
- [33] STOUT, E. L.: Bounded holomorphic functions on finite Riemann surfaces. *Trans. Amer. Math. Soc.* **120** (1965), 255–285.
- [34] WARSCHAWSKI, S. E.: On the higher derivatives at the boundary in conformal mapping. *Trans. Amer. Math. Soc.* **38** (1935), no. 2, 310–340.
- [35] WHITNEY, H.: Analytic extensions of differentiable functions defined in closed sets. *Trans. Amer. Math. Soc.* **36** (1934), no. 1, 63–89.
- [36] WHITNEY, H.: Differentiable manifolds. *Ann. of Math.* (2) **37** (1936), no. 3, 645–680.
- [37] WIRTINGER, W.: Eine Determinantenidentität und ihre Anwendung auf analytische Gebilde in euklidischer und Hermitescher Maßbestimmung. *Monatsh. Math. Phys.* **44** (1936), no. 1, 343–365.
- [38] YANG, P.: Curvatures of complex submanifolds of \mathbb{C}^n . *J. Differential Geom.* **12** (1977/1978), no. 4, 499–511.
- [39] YAU, S.-T.: Review of geometry and analysis. In *Mathematics: frontiers and perspectives*, 353–401. Amer. Math. Soc., Providence, RI, 2000.

Received April 18, 2019; revised October 2, 2019. Published online October 26, 2020.

ANTONIO ALARCÓN: Departamento de Geometría y Topología e Instituto de Matemáticas (IEMath-GR), Universidad de Granada, Campus de Fuentenueva s/n, 18071 Granada, Spain.

E-mail: alarcon@ugr.es

FRANC FORSTNERIČ: Faculty of Mathematics and Physics, University of Ljubljana, Jadranska 19, SI-1000 Ljubljana, Slovenia; and Institute of Mathematics, Physics and Mechanics, Jadranska 19, SI-1000 Ljubljana, Slovenia.

E-mail: franc.forstneric@fmf.uni-lj.si

A.A. is supported by the State Research Agency and European Regional Development Fund via the grant MTM2017-89677-P, MICINN, and the Junta de Andalucía-FEDER grants P18-FR-4049 and A-FQM-139-UGR18, Spain. F.F. is supported by the research program P1-0291 and the research grant J1-9104 from ARRS, Republic of Slovenia.