



# Proper Holomorphic Maps in Euclidean Spaces Avoiding Unbounded Convex Sets

Barbara Drinovec Drnovšek<sup>1,2</sup> · Franc Forstnerič<sup>1,2</sup>

Received: 4 January 2023 / Accepted: 8 February 2023  
© The Author(s) 2023

## Abstract

We show that if  $E$  is a closed convex set in  $\mathbb{C}^n$  ( $n > 1$ ) contained in a closed halfspace  $H$  such that  $E \cap bH$  is nonempty and bounded, then the concave domain  $\Omega = \mathbb{C}^n \setminus E$  contains images of proper holomorphic maps  $f : X \rightarrow \mathbb{C}^n$  from any Stein manifold  $X$  of dimension  $< n$ , with approximation of a given map on closed compact subsets of  $X$ . If in addition  $2 \dim X + 1 \leq n$  then  $f$  can be chosen an embedding, and if  $2 \dim X = n$ , then it can be chosen an immersion. Under a stronger condition on  $E$ , we also obtain the interpolation property for such maps on closed complex subvarieties.

**Keywords** Stein manifold · Holomorphic embedding · Oka manifold · Minimal surface · Convexity

**Mathematics Subject Classification** 32H02 · 32Q56; 52A20 · 53A10

## 1 Introduction

Let  $X$  be a Stein manifold. Denote by  $\mathcal{O}(X, \mathbb{C}^n)$  the Fréchet space of holomorphic maps  $X \rightarrow \mathbb{C}^n$  endowed with the compact-open topology and write  $\mathcal{O}(X, \mathbb{C}) = \mathcal{O}(X)$ . A theorem of Remmert [36], Narasimhan [35], and Bishop [7] states that almost proper maps are residual in  $\mathcal{O}(X, \mathbb{C}^n)$  if  $\dim X = n$ , proper maps are dense if  $\dim X < n$ ,

---

In memoriam Nessim Sibony

---

✉ Franc Forstnerič  
franc.forstneric@fmf.uni-lj.si

Barbara Drinovec Drnovšek  
barbara.drinovec@fmf.uni-lj.si

<sup>1</sup> Faculty of Mathematics and Physics, University of Ljubljana, Jadranska 19, 1000 Ljubljana, Slovenia

<sup>2</sup> Institute of Mathematics, Physics and Mechanics, Jadranska 19, 1000 Ljubljana, Slovenia

proper immersions are dense if  $2 \dim X \leq n$ , and proper embeddings are dense if  $2 \dim X < n$ . A proof is also given in the monograph [29] by Gunning and Rossi.

It is natural to ask how much space proper maps need. We pose the following question.

**Problem 1.1** For which domains  $\Omega \subset \mathbb{C}^n$  are proper holomorphic maps (immersions, embeddings)  $X \rightarrow \mathbb{C}^n$  as above, with images contained in  $\Omega$ , dense in  $\mathcal{O}(X, \Omega)$ ?

It is evident that  $\Omega$  cannot be contained in a halfspace of  $\mathbb{C}^n$  since every holomorphic map from  $\mathbb{C}$  to a halfspace lies in a complex hyperplane. In this paper, we give an affirmative answer for concave domains of which complement  $E = \mathbb{C}^n \setminus \Omega$  satisfies the following condition.

**Definition 1.2** A closed convex set  $E$  in a finite dimensional affine space  $V$  has *bounded convex exhaustion hulls* (BCEH) if for every compact convex set  $K$  in  $V$

$$\text{the set } h(E, K) = \text{Conv}(E \cup K) \setminus E \text{ is bounded.} \quad (1.1)$$

Here,  $\text{Conv}$  denotes the convex hull. The following is our first main result.

**Theorem 1.3** *Let  $E$  be an unbounded closed convex set in  $\mathbb{C}^n$  ( $n > 1$ ) with bounded convex exhaustion hulls. Given a Stein manifold  $X$  with  $\dim X < n$ , a compact  $\mathcal{O}(X)$ -convex set  $K$  in  $X$ , and a holomorphic map  $f_0 : K \rightarrow \mathbb{C}^n$  with  $f_0(bK) \subset \Omega = \mathbb{C}^n \setminus E$ , we can approximate  $f_0$  uniformly on  $K$  by proper holomorphic maps  $f : X \rightarrow \mathbb{C}^n$  satisfying  $f(X \setminus \mathring{K}) \subset \Omega$ . The map  $f$  can be chosen an embedding if  $2 \dim X < n$  and an immersion if  $2 \dim X \leq n$ .*

In this paper, a map  $f : K \rightarrow \mathbb{C}^n$  from a compact set  $K$  is said to be holomorphic if it is the restriction to  $K$  of a holomorphic map on an open neighbourhood of  $K$ .

In particular, if  $f_0(K) \subset \Omega$ , then the theorem gives uniform approximation of  $f_0$  by proper holomorphic maps  $f : X \rightarrow \mathbb{C}^n$  with  $f(X) \subset \Omega$ . If  $bE$  is of class  $\mathcal{C}^1$  and strictly convex near infinity, we obtain an analogue of Theorem 1.3 with additional interpolation on a closed complex subvariety  $X'$  of  $X$  such that  $f_0 : X' \rightarrow \mathbb{C}^n$  is proper holomorphic; see Theorem 4.2. Without the condition on the range, interpolation of proper holomorphic embeddings  $X \hookrightarrow \mathbb{C}^n$  on a closed complex subvariety was obtained by Acquistapace et al. [1] in 1975.

The analogue of the BCEH condition for unbounded closed sets in Stein manifolds, with the convex hull replaced by the holomorphically convex hull, is used in holomorphic approximation theory of Arakelyan and Carleman type; see the survey in [18].

It is evident that a closed convex set  $E \subset \mathbb{R}^n$  has BCEH if and only if there is an increasing sequence  $K_1 \subset K_2 \subset \dots$  of compact convex sets exhausting  $\mathbb{R}^n$  such that the set  $h(E, K_j)$  (see (1.1)) is bounded for every  $j = 1, 2, \dots$ . In particular, BCEH is a condition at infinity which is invariant under perturbations supported on a compact subset. For compact convex sets  $E \subset \mathbb{C}^n$ , Theorem 1.3 was proved in [24]; in this case BCEH trivially holds.

We show in Sect. 3 that a closed convex set  $E$  in  $\mathbb{R}^n$  has BCEH if and only if  $E$  is continuous in the sense of Gale and Klee [26]; see Proposition 3.3. If  $E$  has BCEH,

then  $\text{Conv}(E \cup K)$  is closed for any compact convex set  $K \subset \mathbb{R}^n$  (see [26, Theorem 1.5]). If such  $E$  is unbounded, which is the main case of interest, there are affine coordinates  $(x, y) \in \mathbb{R}^{n-1} \times \mathbb{R}$  such that  $E = E_\phi = \{(x, y) \in \mathbb{R}^n : y \geq \phi(x)\}$  is the epigraph of a convex function  $\phi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}_+ = [0, +\infty)$  growing at least linearly near infinity (see Proposition 3.4). In particular, an unbounded closed convex set  $E \subset \mathbb{C}^n$  with BCEH is of the form

$$E = E_\phi = \{z = (z', z_n) \in \mathbb{C}^n : \Re z_n \geq \phi(z', \Im z_n)\}, \tag{1.2}$$

in some affine complex coordinates  $z = (z', z_n)$  on  $\mathbb{C}^n$ , with  $\phi$  as above. (Here,  $\Re$  and  $\Im$  denote the real and the imaginary part.) For a convex function  $\phi$  of class  $\mathcal{C}^1$ , we give a differential characterization of the BCEH condition on its epigraph  $E_\phi$ ; see Proposition 3.8. The BCEH property holds if the radial derivative of  $\phi$  tends to infinity; see Corollary 3.9. On the other hand, there are convex functions of linear growth of which epigraphs have BCEH; see Example 3.10. By Proposition 3.11, a convex function  $\phi$  with at least linear growth at infinity can be approximated uniformly on compacts by functions  $\psi \leq \phi$  of the same kind of which epigraphs  $E_\psi$  have BCEH. This allows us to extend Theorem 1.3 as follows; see Sect. 4 for the proof.

**Corollary 1.4** *The conclusion of Theorem 1.3 holds for any convex epigraph  $E_\phi$  of the form (1.2) such that  $\phi \geq 0$  and the set  $\{\phi = 0\}$  is nonempty and compact.*

A closed convex set  $E \subset \mathbb{C}^n$  with BCEH does not contain any affine real line (see Proposition 3.4), and for  $n > 1$ , its complement  $\Omega = \mathbb{C}^n \setminus E$  is an Oka domain according to Wold and the second named author; see [25, Theorem 1.8]. This fact plays an important role in our proof of Theorem 1.3, given in Sect. 4. (The precise result from Oka theory which we shall use is stated as Theorem 4.1.) Among closed convex epigraphs (1.2), the class of sets with Oka complement is strictly bigger than the class of sets with BCEH. In particular, the former class contains many sets containing boundary lines, which is impossible for a set with BCEH.

**Problem 1.5** Is there a (not necessarily convex) set  $E_\phi \subset \mathbb{C}^n$  of the form (1.2) with  $\phi \geq 0$  of sublinear growth for which Theorem 1.3 holds? Is there a set of this kind in  $\mathbb{C}^2$  such that  $\mathbb{C}^2 \setminus E_\phi$  contains the image of a proper holomorphic disc  $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\} \rightarrow \mathbb{C}^2$ ?

Theorem 1.3 is the first general result in the literature providing proper holomorphic maps  $X \rightarrow \mathbb{C}^n$  from any Stein manifold of dimension  $< n$  whose images avoid large convex sets in  $\mathbb{C}^n$  close to a halfspace, and with approximation of a given map on a compact holomorphically convex set in  $X$ . Without the approximation condition and assuming that  $\dim X \leq n - 2$ , there are proper holomorphic maps of  $X$  into a complex hyperplane in  $\mathbb{C}^n \setminus E$ .

On the other hand, there are many known results concerning proper holomorphic maps in Euclidean spaces and in more general Stein manifolds whose images avoid certain small closed subsets, such as compact or complete pluripolar ones, and results in which the source manifold is the disc  $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ . Proper holomorphic discs in  $\mathbb{C}^2$  avoiding closed complete pluripolar sets of the form  $E = E' \times \mathbb{C}$ , with  $E' \subset \mathbb{C}$ , were constructed by Alexander [5] in 1977. The first named author showed

in [13] that for every closed complete pluripolar set  $E$  in a Stein manifold  $Y$  with  $\dim Y > 1$  and point  $p \in Y \setminus E$ , there is a proper holomorphic disc  $f : \mathbb{D} \rightarrow Y$  with  $p \in f(\mathbb{D}) \subset Y \setminus E$ . If  $Y = \mathbb{C}^2$ , there also exist embedded holomorphic discs with this property according to Borell et al. [8], and for  $\dim Y \geq 3$ , this holds by the general position argument. Proper holomorphic discs in  $\mathbb{C}^2$  with images contained in certain concave cones were constructed by Globevnik and the second named author [23] in 2001. They also constructed proper holomorphic discs in  $\mathbb{C}^2$  with images in  $(\mathbb{C} \setminus \{0\})^2$ , and hence, proper harmonic discs  $\mathbb{D} \rightarrow \mathbb{R}^2$ , disproving a conjecture by Schoen and Yau [37, p. 18]. (Another construction of such maps was given by Božin [9].) More generally, it was shown by Alarcón and López [4, Corollary 1.1] in 2012 that every open Riemann surface  $X$  admits a proper harmonic map to  $\mathbb{R}^2$  which is the projection of a conformal minimal immersion  $X \rightarrow \mathbb{R}^3$ . The aforementioned result from [23] was used by the first named author in [12] to classify closed convex sets in  $\mathbb{C}^2$  whose complement is filled by images of holomorphic discs which are proper in  $\mathbb{C}^2$ . More recently, Forstnerič and Ritter [24] proved Theorem 1.3 in the case when  $E \subset \mathbb{C}^n$  is a compact polynomially convex set and  $2 \dim X \leq n$  (for immersions) or  $2 \dim X < n$  (for embeddings), and for proper holomorphic maps  $X \rightarrow \mathbb{C}^n$  when  $\dim X < n$  and  $E$  is a compact convex set. A further development in this direction is the analogue of Theorem 1.3 when  $\mathbb{C}^n$  is replaced by a Stein manifold  $Y$  with the density property and  $E \subset Y$  is a compact  $\mathcal{O}(Y)$ -convex set; see [22, Remark 4.5] and the references therein. However, in all mentioned results except those in [12, 23], the avoided sets are thin or compact.

Without insisting on approximation, the theorem of Remmert, Bishop, and Narasimhan is not optimal with respect to the dimension of the target space. Indeed, it was shown by Eliashberg and Gromov [17] in 1992, with an improvement for odd dimensional Stein manifolds by Schürmann [38] in 1997 that a Stein manifold  $X$  of dimension  $m \geq 2$  embeds properly holomorphically in  $\mathbb{C}^n$  with  $n = \lceil \frac{3m}{2} \rceil + 1$ , and for  $m \geq 1$  it immerses properly holomorphically in  $\mathbb{C}^n$  with  $n = \lceil \frac{3m+1}{2} \rceil$ . (See also [20, Sect. 9.3].) However, the construction method in these papers, which relies on the Oka principle for sections of certain stratified holomorphic fibre bundles, does not give the density statement, and we do not know whether Theorem 1.3 holds for maps to these lower dimensional spaces. It is also an open problem whether every open Riemann surface embeds properly holomorphically in  $\mathbb{C}^2$ ; see [20, Secs. 9.10–9.11] and the survey [21].

Theorem 1.3 is proved in Sect. 4. The proof relies on two main ingredients. One is the result of Wold and the second named author [25, Theorem 1.8] which shows in particular that the complement  $\Omega = \mathbb{C}^n \setminus E$  of a closed convex set  $E$  having BCEH is an Oka domain. The second main technique comes from the work of Dor [10, 11], following earlier papers by Stensønes [39] and Hakim [30]. Dor constructed proper holomorphic immersions and embeddings of any smoothly bounded, relatively compact, strongly pseudoconvex domain  $D$  in a Stein manifold  $X$  into any pseudoconvex domain  $\Omega$  in  $\mathbb{C}^n$  under the dimension conditions in Theorem 1.3. Previously, Hakim [30] constructed proper holomorphic maps to balls in codimension one. The main idea is to inductively lift the image of  $bD$  under a holomorphic map  $f : \bar{D} \rightarrow \Omega$  to a given higher superlevel set of a strongly plurisubharmonic exhaustion function

$\rho : \Omega \rightarrow \mathbb{R}_+$  in a controlled way, taking care not to decrease the value of  $\rho \circ f$  very much anywhere on  $D$  during the process. When  $D$  is a finite bordered Riemann surface, this can be achieved by using approximate solutions of a Riemann-Hilbert boundary value problem (see [14]). In higher dimensions, the proof is more subtle and uses carefully controlled holomorphic peak functions on  $\bar{D}$  to push a given map  $f : \bar{D} \rightarrow \Omega$  locally at a point  $z \in f(bD)$  in the direction of the zero set  $S_z$  of the holomorphic (quadratic) Levi polynomial of the exhaustion function  $\rho : \Omega \rightarrow \mathbb{R}$ . At a noncritical point  $z \in \Omega$  of  $\rho$ ,  $S_z$  is a smooth local complex hypersurface and the restricted function  $\rho|_{S_z}$  increases quadratically as we move away from  $z$ . If  $\rho$  is a *strictly convex* function, this can be achieved by pushing the image of  $f(bD)$  in the direction of suitably chosen affine complex hyperplanes. Dor's construction was extended by the authors to maps from strongly pseudoconvex domains in Stein manifolds to an arbitrary Stein manifold  $\Omega$ , and also to  $q$ -convex complex manifolds for suitable values of  $q \in \mathbb{N}$ ; see the papers [14, 15] from 2007 and 2010, respectively. In those papers, we introduced the technique of glueing holomorphic sprays of manifold-valued maps on a strongly pseudoconvex Cartan pair with control up to the boundary (a nonlinear version of the Cousin-I problem) and a systematic approach for avoiding critical points of a  $q$ -convex Morse exhaustion function on  $\Omega$ .

Earlier constructions of this type, using simpler holomorphic peak functions and higher codimension, were given in 1985 by Løw [34] and Forstnerič [19] who showed that every relatively compact strongly pseudoconvex domain  $D$  in a Stein manifold embeds properly holomorphically in a high dimensional Euclidean ball. A related result with interpolation on a suitable subset of the boundary of  $D$  is due to Globevnik [27]. This peak function technique was inspired by the construction of inner functions on the ball of  $\mathbb{C}^n$  by Løw [33] in 1982, based on the work of Hakim and Sibony [31].

We apply this technique to push the boundary  $f_0(bD) \subset \Omega = \mathbb{C}^n \setminus E$  of a holomorphic map  $f_0 : \bar{D} \rightarrow \mathbb{C}^n$  in Theorem 1.3 out of a certain compact convex cap  $C$  attached to  $E$  along a part of  $bC$  contained in  $bE$  and such that the set  $E_1 = E \cup C$  is convex and has bounded convex exhaustion hulls. At the same time, we ensure that the new map  $g : \bar{D} \rightarrow \mathbb{C}^n$  still sends  $\bar{D} \setminus \bar{K}$  to  $\Omega$ . For a precise result, see Proposition 2.1. In the next step, we use that  $\Omega_1 = \mathbb{C}^n \setminus E_1$  is an Oka domain (see Corollary 3.6). Since  $g(bD) \subset \Omega_1$ , we can apply the Oka principle (see Theorem 4.1) to approximate  $g$  by a holomorphic map  $f_1 : X \rightarrow \mathbb{C}^n$  with  $f_1(X \setminus D) \subset \Omega_1$ . Continuing inductively, we obtain a sequence of holomorphic maps  $X \rightarrow \mathbb{C}^n$  converging to a proper map satisfying Theorem 1.3. The details are given in Sect. 4.

The analogues of Theorem 1.3 and Corollary 1.4 also hold for minimal surfaces in  $\mathbb{R}^n$ .

**Theorem 1.6** *Let  $n \geq 3$ , and let  $\phi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}_+$  be a convex function such that the set  $\{\phi = 0\}$  is nonempty and compact. Given an open Riemann surface  $X$ , a compact  $\mathcal{O}(X)$ -convex set  $K$  in  $X$ , and a conformal minimal immersion  $f_0 : U \rightarrow \mathbb{R}^n$  from a neighbourhood of  $K$  with  $f_0(bK) \subset \Omega = \{y < \phi(x)\}$ , we can approximate  $f_0$  uniformly on  $K$  by proper conformal minimal immersions  $f : X \rightarrow \mathbb{R}^n$  (embeddings if  $n \geq 5$ ) satisfying  $f(X \setminus \mathring{K}) \subset \Omega$ .*

If in addition  $\phi$  is of class  $\mathcal{C}^1$ , strictly convex at infinity, and the epigraph  $E_\phi = \{y \geq \phi(x)\}$  has BCEH then one can add to this statement the interpolation of the map on discrete sets, in analogy to Theorem 4.2.

Theorem 1.6 is obtained by following the proof of Theorem 1.3, replacing Proposition 2.1 by the analogous result obtained by the Riemann–Hilbert deformation method for conformal minimal surfaces (see [2] or [3, Chapter 6]). Furthermore, it has recently been shown by the authors [16, Corollary 1.5] that the complement of a closed convex set  $E \subset \mathbb{R}^n$  ( $n \geq 3$ ) is flexible for minimal surfaces (an analogue of the Oka property in complex geometry) if and only if  $E$  is not a halfspace or a slab; clearly this includes all sets with BCEH.

Another method for constructing proper minimal surfaces, which yields the same result in some examples not covered by Theorem 1.6, was developed by Alarcón and López [4] in 2012. They showed that Theorem 1.6 holds for any wedge domain  $\Gamma \times \mathbb{R} \subset \mathbb{R}^3$ , where  $\Gamma \subset \mathbb{R}^2$  is an open cone with angle  $> \pi$ ; see [4, Theorem 5.6]. The complement of this set is convex, but it fails to satisfy the hypotheses of Theorem 1.6 due to the presence of lines in the boundary. An important difference between these two fields, which affects the possible construction methods, is that every open Riemann surface admits a proper harmonic map to the plane  $\mathbb{R}^2$  (see [4, Theorem I]), while only few such surfaces admit proper holomorphic maps to  $\mathbb{C}$ .

The analogue of Problem 1.5 for minimal surfaces asks whether there is a domain in  $\mathbb{R}^3$  of the form  $\{x_3 < \phi(x_1, x_2)\}$ , where  $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}_+$  is a function with sublinear growth, which contains minimal surfaces of hyperbolic type that are proper in  $\mathbb{R}^3$ , or just a proper hyperbolic end of a minimal surface. In particular, it would be interesting to know whether the domain below the upper half of a vertical catenoid has this property. On the other hand, the strong halfspace theorem of Hoffman and Meeks [32] says that the only proper minimal surfaces in  $\mathbb{R}^3$  contained in a halfspace are planes.

## 2 Pushing a Strongly Pseudoconvex Boundary Out of a Strictly Convex Cap

Let  $O$  be a convex domain in  $\mathbb{C}^n$  for some  $n > 1$ . Recall that a continuous function  $\rho : O \rightarrow \mathbb{R}$  is said to be *strictly convex* if for any pair of points  $a, b \in O$ , we have that

$$\rho(ta + (1 - t)b) < t\rho(a) + (1 - t)\rho(b) \text{ for all } 0 < t < 1.$$

Assume now that  $\rho_t : O \rightarrow \mathbb{R}$  ( $t \in [0, 1]$ ) is a continuous family of  $\mathcal{C}^1$  functions satisfying the following conditions:

- (a) For every  $t \in [0, 1]$  the function  $\rho_t$  is strictly convex. Note that  $d\rho_t \neq 0$  on  $M_t := \{\rho_t = 0\}$ .
- (b) If  $0 \leq s < t \leq 1$ , then  $\rho_t \leq 0$  on  $M_s$ .
- (c) There is an open relatively compact subset  $\omega_0$  of  $M_0$  such that for every pair of numbers  $0 \leq s < t \leq 1$ , we have that  $M_t \cap M_0 = M_t \cap M_s = M_0 \setminus \omega_0$ .

This means that the hypersurfaces  $M_t$  coincide on the subset  $M_0 \setminus \omega_0$ , and as  $t \in [0, 1]$  increases the domains  $\omega_t = M_t \setminus M_0 \subset M_t$  are pairwise disjoint and move into the convex direction. Each compact set of the form:

$$C_t = \bigcup_{s \in [0,t]} \bar{\omega}_s \text{ for } t \in [0, 1], \tag{2.1}$$

is called a *strictly convex cap with the base  $\omega_0$* . Note that  $bC_t = \bar{\omega}_0 \cup \bar{\omega}_t$ ,  $C_t$  is strictly convex along  $\omega_t$ , strictly concave along  $\omega_0$ , and it has corners along  $\bar{\omega}_0 \cap \bar{\omega}_t$ . As  $t \in [0, 1]$  increases to 1, the caps  $C_t$  monotonically increase to  $C_1$  and they share the same base  $\omega_0$ . Likewise, for any  $0 \leq s < t \leq 1$ , the set  $C_{s,t} = \bigcup_{u \in [s,t]} \bar{\omega}_u$  is a strictly convex cap with the base  $\omega_s$ . The sets

$$E_t = \{z \in O : \rho_t(z) \leq 0\} \text{ for } t \in [0, 1], \tag{2.2}$$

are strictly convex along  $bE_t = \{\rho_t = 0\}$ , they form a continuously increasing family in  $t$ , and

$$E_t = E_0 \cup C_t \text{ for every } t \in [0, 1].$$

Under these assumptions, we have the following result.

**Proposition 2.1** *Let  $D$  be a smoothly bounded, relatively compact, strongly pseudoconvex domain in a Stein manifold  $X$  with  $\dim X < n$ . Let the sets  $E_t \subset O \subset \mathbb{C}^n$  ( $t \in [0, 1]$ ) be given by (2.2), and let  $f_0 : \bar{D} \rightarrow O$  be a map of class  $\mathcal{A}(\bar{D})$  such that  $f_0(bD) \cap E_0 = \emptyset$ . Given a compact set  $K \subset D$  such that  $f_0(\bar{D} \setminus K) \cap E_0 = \emptyset$  and a number  $\epsilon > 0$ , there is a map  $f : \bar{D} \rightarrow O$  of class  $\mathcal{A}(\bar{D})$  satisfying the following conditions:*

- (i)  $f(bD) \cap E_1 = \emptyset$ ,
- (ii)  $f(\bar{D} \setminus K) \cap E_0 = \emptyset$ , and
- (iii)  $\max_{x \in K} |f(x) - f_0(x)| < \epsilon$ .

Recall that a map  $f : \bar{D} \rightarrow O$  is said to be of class  $\mathcal{A}(\bar{D})$  if it is continuous on  $\bar{D}$  and holomorphic on  $D$ . In our application of Proposition 2.1 in the proof of Theorem 1.3, the set  $O$  will be a ball (or the entire Euclidean space) and the hypersurfaces  $M_t = \{\rho_t = 0\} = bE_t$  will be convex graphs over the coordinate hyperplane  $\mathbb{C}^{n-1} \times \mathbb{R} \subset \mathbb{C}^n$ .

In the proof of Proposition 2.1 we shall need the following lemma.

**Lemma 2.2** *Assume that  $O$  is a convex open subset of  $\mathbb{C}^n$  for  $n > 1$ ,  $L$  is a compact subset of  $O$ , and  $\rho : O \rightarrow \mathbb{R}$  is a  $\mathcal{C}^1$  smooth strictly convex function. Then there is a number  $\delta > 0$  with the following property. If  $D$  is a smoothly bounded strongly pseudoconvex domain in a Stein manifold  $X$  of dimension  $\dim X = m < n$ ,  $K$  is a compact subset of  $D$ , and  $f : \bar{D} \rightarrow O$  is a map of class  $\mathcal{A}(\bar{D})$  such that*

$$\rho(f(z)) > -\delta \text{ for all } z \in bD \text{ and } \rho(f(z)) > 0 \text{ if } z \in bD \text{ and } f(z) \notin L, \tag{2.3}$$

*then given  $\eta > 0$ , there is a map  $g : \bar{D} \rightarrow O$  of class  $\mathcal{A}(\bar{D})$  satisfying the following conditions:*

- (i)  $\rho(g(z)) > 0$  for  $z \in bD$ ,
- (ii)  $\rho(g(z)) > \delta$  for those  $z \in bD$  for which  $g(z) \in L$ ,
- (iii)  $\rho(g(z)) > \rho(f(z)) - \eta$  for  $z \in \overline{D \setminus K}$ , and
- (iv)  $|f(z) - g(z)| < \eta$  for  $z \in K$ .

For  $m = 1$ , i.e. when  $D$  is a finite bordered Riemann surface, this is a simplified version of [14, Lemmas 6.2 and 6.3], which is proved by using approximate solutions of a Riemann–Hilbert boundary value problem. This method was employed in several earlier papers mentioned in [14]. When  $\rho$  is strictly convex,  $\mathcal{C}^1$  smoothness suffices since in the proof we may take a continuous family of tangential linear discs to the sublevel set of  $\rho$ .

For  $m \geq 2$ , Lemma 2.2 is a simplified and slightly modified version of [15, Lemma 5.3]. Besides the fact that we are considering single maps  $\bar{D} \rightarrow O$  instead of sprays of maps, the only difference is that the assumption in [15, Lemma 5.3] that the set  $\{\rho = 0\}$  is compact is replaced by the assumption (2.3) saying that  $\rho(f(z))$  for  $z \in bD$  may be negative only if  $f(z)$  lies in the compact set  $L \subset O$ . This hypothesis ensures that the lifting for a relatively big amount (the role of the constant  $\delta$ ) only needs to be made on a compact subset of  $O$ , while elsewhere it suffices to pay attention not to decrease  $\rho \circ f$  by more than a given amount and to approximate sufficiently closely on  $K$  (the role of the constant  $\eta$ ). The proof requires only a minor adaptation of [15, proof of Lemma 5.3], using its local version [15, Lemma 5.2] in a finite induction with respect to a covering of  $bD$  by small open sets on which there are good systems of local holomorphic peak functions. In fact, Lemma 2.2 corresponds to a simplified version of [15, Sublemma 5.4], which explains how to lift the image of  $bD$  with respect to  $\rho$  for a sufficiently large amount at those points in  $bD$  which the map  $f$  sends to a certain coordinate chart  $U_i$  in the target manifold. In our situation, the role of  $U_i$  is played by an open relatively compact neighbourhood of the set  $L \cap \{\rho = 0\}$  in  $O$ , and there is no need to use the rest of the proof of [15, Lemma 5.3].

**Proof of Proposition 2.1** For  $t \in [0, 1]$ , let  $\delta_t > 0$  be a number for which the conclusion of Lemma 2.2 holds for the function  $\rho_t$  and the compact set  $L = C_1$  (see (2.1)). The open sets

$$U_t = \{z \in O : -\delta_t < \rho_t(z) < \delta_t\} \text{ for } t \in [0, 1],$$

form an open covering of  $C_1$ , so there exists a finite subcovering  $\{U_{t_j}\}$  for  $0 \leq t_1 < t_2 < \dots < t_k \leq 1$ . Applying Lemma 2.2, we inductively find maps  $f_1, \dots, f_k \in \mathcal{A}(\bar{D})$  such that for every  $j = 1, \dots, k$ , we have that

- (a)  $f_j(bD) \cap E_{t_j} = \emptyset$  (where  $E_t$  is given by (2.2)),
- (b)  $f_j(\overline{D \setminus K}) \cap \bar{E}_0 = \emptyset$ , and
- (c)  $|f_j - f_{j-1}| < \epsilon/k$  on  $K$ .

Note that conditions (a) and (b) hold for  $f_0$  and (c) is void. Assume inductively that for some  $j \in \{1, \dots, k\}$  we have maps  $f_0, \dots, f_{j-1}$  satisfying these conditions. Applying Lemma 2.2 with  $f = f_{j-1}$  and taking  $f_j = g$ , condition (a) follows from part (i) in Lemma 2.2, (b) follows from (ii) provided that the number  $\eta > 0$  in Lemma 2.2 is chosen small enough, and (c) follows from (iii) in Lemma 2.2 provided that  $\eta \leq \epsilon/k$ .



This gives the map  $f_j$  satisfying conditions (a)–(c) and the induction may continue. The map  $f = f_k$  then satisfies the proposition.  $\square$

**Remark 2.3** Proposition 2.1 also holds, with the same proof, if  $\rho_t$  ( $t \in [0, 1]$ ) are strongly plurisubharmonic functions of class  $\mathcal{C}^2$  satisfying  $d\rho_t \neq 0$  on  $M_t = \{\rho_t = 0\}$ . Indeed, the results from [15], which are used in the proof, pertain to this case. In the present paper we shall only use the convex case under  $\mathcal{C}^1$  smoothness, which comes naturally in the construction.

### 3 Closed Convex Sets with BCEH

In the context of convex analysis, closed unbounded convex sets that share several important properties with compact convex sets were studied by Gale and Klee [26] in 1959. They introduced the class of continuous sets, and we show that this class coincides with the class of sets having BCEH, introduced in Definition 1.2; see Proposition 3.3. We then develop further properties of these sets which are relevant to the proof of our main theorems.

By a ray in  $\mathbb{R}^n$ , we shall mean a closed affine halfline. Let  $E$  be a closed convex subset of  $\mathbb{R}^n$ . A boundary ray of  $E$  is a ray contained in the boundary of  $E$ . An asymptote of  $E$  is a ray  $L \subset \mathbb{R}^n \setminus E$  such that  $\text{dist}(L, E) = \inf\{|x - y| : x \in L, y \in E\} = 0$ . The function

$$\sigma : \{u \in \mathbb{R}^n : |u| = 1\} \rightarrow \mathbb{R} \cup \{+\infty\}, \quad \sigma(u) = \sup\{x \cdot u : x \in E\},$$

is called the support function of  $E$ . (Here,  $x \cdot u$  denotes the Euclidean inner product.) A closed convex set  $E$  is said to be continuous in the sense of Gale and Klee [26] if the support function of  $E$  is continuous. Note that every compact convex set is continuous.

The following result is a part of [26, Theorem 1.3] due to Gale and Klee; we only list those conditions that will be used. The last item (iv) uses also [26, Theorem 1.5].

**Theorem 3.1** For a closed convex subset  $E$  in  $\mathbb{R}^n$  the following conditions are equivalent:

- (i)  $E$  is continuous.
- (ii)  $E$  has no boundary ray nor asymptote.
- (iii) For each point  $p \in \mathbb{R}^n$  the convex hull  $\text{Conv}(E \cup \{p\})$  is closed.
- (iv) For every compact convex set  $K \subset \mathbb{R}^n$  the set  $\text{Conv}(E \cup K)$  is closed.

Condition (iii) implies that the closed convex hull  $\overline{\text{Conv}}(E \cup \{p\})$  is the union of the line segments connecting  $p$  to the points in  $E$ . It also shows that an unbounded continuous closed convex subset  $E$  of  $\mathbb{R}^n$  is not contained in any affine hyperplane.

Let us record the following observation which will be used in the sequel.

**Lemma 3.2** Let  $E \subset \mathbb{R}^n$  be a closed convex set,  $p \in \mathbb{R}^n \setminus E$ , and  $L \subset \mathbb{R}^n$  be an affine subspace containing  $p$ . Then,  $\text{Conv}(E \cup \{p\}) \cap L = \text{Conv}((E \cap L) \cup \{p\})$ .

**Proof** Set  $E' = E \cap L$ . It is obvious that  $\text{Conv}(E' \cup \{p\}) \subset \text{Conv}(E \cup \{p\}) \cap L$ . Conversely, since  $E$  is convex, every point  $q \in \text{Conv}(E \cup \{p\})$  belongs to a line

segment from  $p$  to a point  $q' \in E$ . If in addition  $q \in L$  and  $q \neq p$  then  $q' \in E'$ , and hence  $q \in \text{Conv}(E' \cup \{p\})$ .  $\square$

**Proposition 3.3** *A closed convex set  $E \subset \mathbb{R}^n$  has BCEH if and only if it is continuous in the sense of Gale and Klee [26].*

**Proof** Since all closed bounded convex sets have BCEH and are continuous, it suffices to consider the case when the set  $E$  is unbounded.

If  $E$  is not continuous then by Theorem 3.1 it has a boundary ray or an asymptote. Denote it by  $L$ , and let  $\ell$  be the affine line containing  $L$ . Pick any affine 2-plane  $H \subset \mathbb{R}^n$  containing  $\ell$  such that  $H \not\subset E$  and a point  $p \in H \setminus (\ell \cup E)$ . By considering rays from  $p$  to points  $q \in E$  approaching  $L$  and going to infinity (if  $L$  is a boundary ray, we can choose points  $q \in L$ ), we see that the closure of the set  $h(E, p) = \text{Conv}(E \cup \{p\}) \setminus E$  contains the parallel translate  $L' \subset H$  of  $L$  passing through  $p$ , so  $h(E, p)$  is unbounded and hence  $E$  does not have BCEH.

Assume now that  $E$  is continuous and let us prove that it has BCEH. We need to show that for any closed ball  $B \subset \mathbb{R}^n$  the set  $h(E, B) = \text{Conv}(E \cup B) \setminus E$  is bounded. Assume to the contrary that there is a sequence  $x_m \in h(E, B)$  with  $|x_m| \rightarrow \infty$  as  $m \rightarrow \infty$ . Since the sets  $E$  and  $B$  are convex, we have that

$$x_m = t_m b_m + (1 - t_m) e_m \text{ for } t_m \in [0, 1], b_m \in B, e_m \in E, \text{ and } m \in \mathbb{N}.$$

Note that  $(1 - t_m)|e_m| \rightarrow \infty$  as  $m \rightarrow \infty$ . By compactness of the respective sets we may assume, passing to a subsequence, that  $e_m \neq 0$  for all  $m$  and the sequences  $t_m$ ,  $b_m$ , and  $\frac{1}{|e_m|} e_m$  are convergent. Denote their respective limits by  $t$ ,  $b$ , and  $f$ . We have that

$$x_m = t_m b_m + (1 - t_m) e_m = b_m + (1 - t_m) |e_m| \left( \frac{e_m}{|e_m|} - \frac{b_m}{|e_m|} \right) = b_m + (1 - t_m) |e_m| f_m,$$

where  $f_m = \left( \frac{e_m}{|e_m|} - \frac{b_m}{|e_m|} \right)$ . Note that  $\lim_{m \rightarrow \infty} f_m = f$ . Pick a number  $\alpha \geq 0$  and set  $p = b + \alpha f$ . If  $m$  is large enough then  $(1 - t_m)|e_m| > \alpha$ , so the point  $y_m = b_m + \alpha f_m$  lies on the line segment connecting  $b_m$  and  $x_m$ . Since  $x_m \in \text{Conv}(E \cup \{b_m\})$ , it follows that  $y_m \in \text{Conv}(E \cup \{b_m\})$ . Note that the sequence  $y_m$  converges to  $p$ . Since  $E$  is continuous,  $\text{Conv}(E \cup \{b\})$  is closed by Theorem 3.1, so  $p = \lim_{m \rightarrow \infty} y_m \in \text{Conv}(E \cup \{b\})$ . Since this holds for every  $\alpha \geq 0$ , the ray  $L = \{b + \alpha f : \alpha \in [0, \infty)\}$  lies in  $\text{Conv}(E \cup \{b\})$ . By Lemma 3.2 there is  $\alpha_0 \in [0, \infty)$  such that the ray  $L' = \{b + \alpha f : \alpha \geq \alpha_0\}$  lies in  $E$ . Since  $E$  is continuous,  $L$  is not a boundary ray of  $E$  by Theorem 3.1, thus  $L$  contains a point  $q = b + \alpha_1 f \in E \setminus bE$  for some  $\alpha_1 \geq \alpha_0$ . Choose a neighbourhood  $U_q \subset E$  of  $q$ . For any large enough  $m$  we then have  $p_m := b_m + \alpha_1 f_m \in U_q$ . Let  $L_m = \{b_m + \alpha f_m : \alpha \geq 0\}$ . Note that  $L_m \cap \text{Conv}(E \cup \{b_m\}) = \text{Conv}((L_m \cap E) \cup \{b_m\})$  by Lemma 3.2. However, for  $m$  large enough the point  $x_m \in L_m$  lies on the opposite side of  $p_m$  than  $b_m$ , so  $x_m$  belongs to  $L_m \cap \text{Conv}(E \cup \{b_m\})$  but not to  $\text{Conv}((L_m \cap E) \cup \{b_m\})$ . This contradiction proves that  $E$  has BCEH.  $\square$

Given a function  $\phi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ , the *epigraph* of  $\phi$  is the set

$$E = E_\phi = \{(x, y) \in \mathbb{R}^{n-1} \times \mathbb{R} : y \geq \phi(x)\}. \tag{3.1}$$

Note that a function is convex if and only if its epigraph is convex.

**Proposition 3.4** *If  $E \subsetneq \mathbb{R}^n$  is a closed unbounded convex set with BCEH then*

- (i)  *$E$  does not contain any affine real line, and*
- (ii) *for every affine line  $\ell$  intersecting  $E$  in a ray and any hyperplane  $H$  transverse to  $\ell$ ,  $E$  is the epigraph of a convex function on  $H$ . In particular, there are affine coordinates  $(x, y)$  on  $\mathbb{R}^n$  in which  $E$  is of the form (3.1) for a convex function  $\phi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}_+$  satisfying*

$$\liminf_{|x| \rightarrow +\infty} \frac{\phi(x)}{|x|} > 0. \tag{3.2}$$

The condition (3.2) says that  $\phi$  grows at least linearly at infinity. We show in Example 3.10 that there are convex epigraphs  $E_\phi$  having BCEH such that the function  $\phi$  has linear growth.

**Proof** (i) Assume that  $\ell \subset E$  is an affine line and let us prove that  $E$  does not have BCEH. Since  $E$  is a proper subset of  $\mathbb{R}^n$ , there is a parallel translate  $\ell'$  of  $\ell$  which is not contained in  $E$ , and hence  $\ell' \setminus E$  contains a ray  $L$ . Let  $p$  be the endpoint of  $L$ , and let  $p' \in L$  be an arbitrary other point. Since  $E \cap L = \emptyset$ , there is a ball  $B$  around  $p'$  such that  $\text{Conv}(B \cup \{p\}) \cap E = \emptyset$ . Clearly, there is a point  $q \in B$  such that the ray  $L_q$  with the endpoint  $p$  and containing  $q$  intersects the line  $\ell$ , so the line segment from  $p$  to  $q$  belongs to  $\text{Conv}(E \cup \{p\}) \setminus E = h(E, p)$ . By moving  $p' \in L$  to infinity we see that  $h(E, p)$  is unbounded, so  $E$  does not have BCEH.

(ii) Since  $E$  is unbounded, it contains a ray  $L$ . Denote by  $\ell$  the affine line containing  $L$ . Let  $\ell'$  be any parallel translate of  $\ell$ . Since  $E$  contains no affine lines by part (i), there is a point  $p \in \ell' \setminus E$ . The closed convex hull of the union of  $L$  and  $p$  contains the parallel translate  $L' \subset \ell'$  of  $L$  passing through  $p$ . Since  $E$  has BCEH, we conclude that  $L' \subset \text{Conv}(E \cup \{p\})$  and  $L' \setminus E$  is bounded. Since  $E \cap L'$  is convex,  $L' \cap E$  is a closed ray with the endpoint on  $bE$ . This shows that  $E$  is a union of closed rays contained in parallel translates of the line  $\ell$ , so it is an epigraph of a convex function defined on any hyperplane  $H \subset \mathbb{R}^n$  transverse to  $\ell$ . Choosing  $H$  such that  $H \cap E = \emptyset$  there are affine coordinates  $(x, y)$  on  $\mathbb{R}^n$  with  $H = \{y = 0\}$  and  $\ell = \{x = 0\}$ . In these coordinates,  $E$  is of the form (3.1) for a positive convex function  $\phi$ . Finally, if condition (3.2) fails then there is a sequence  $(x_k, y_k) \in E$  with  $|x_k| \rightarrow +\infty$  and  $y_k/|x_k| \rightarrow 0$  as  $k \rightarrow \infty$ . The union of the line segments  $L_k$  connecting  $p = (0, -1) \in \mathbb{R}^{n-1} \times \mathbb{R}$  to  $(x_k, y_k)$ , intersected with the lower halfspace  $y \leq 0$ , is then an unbounded subset of  $h(E, p) = \text{Conv}(E \cup \{p\}) \setminus E$ , contradicting the assumption that  $E$  has BCEH. □

**Remark 3.5** The growth condition (3.2) for an epigraph can always be achieved in suitable linear coordinates (even without the BCEH property) if there is a supporting hyperplane  $H \subset \mathbb{R}^n$  for  $E$  such that the set  $E \cap H$  is nonempty and compact. Indeed, we may then choose coordinates  $(x, y)$  on  $\mathbb{R}^n$  such that  $H = \{y = 0\}$ ,  $E \subset \{y \geq 0\}$ , and  $0 \in E$ . If the condition (3.2) fails, there is a sequence  $(x_k, y_k) \in E$  with  $|x_k| \rightarrow +\infty$  and  $y_k/|x_k| \rightarrow 0$  as  $k \rightarrow \infty$ . After passing to a subsequence, a ray in  $E \cap H$  lies in the closure of the union of the line segments  $L_k \subset E$  connecting the origin to  $(x_k, y_k)$ , contradicting the assumption that the latter set is compact.

**Corollary 3.6** *If  $E$  is a closed convex set in  $\mathbb{C}^n$  ( $n > 1$ ) having BCEH then  $\mathbb{C}^n \setminus E$  is Oka.*

**Proof** By Proposition 3.4 the set  $E$  does not contain any affine real line, and hence  $\mathbb{C}^n \setminus E$  is Oka by [25, Theorem 1.8]. □

The following lemma shows that the BCEH condition is stable under uniform approximation.

**Lemma 3.7** *Assume that  $\phi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$  is a convex function whose epigraph  $E_\phi$  (3.1) has BCEH. Then for any  $\epsilon > 0$  and convex function  $\psi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$  satisfying  $|\phi - \psi| < \epsilon$  the epigraph  $E_\psi$  also has BCEH.*

**Proof** If  $E_\psi$  fails to have BCEH then by Theorem 3.1 and Proposition 3.3 it has a boundary ray or an asymptote,  $L$ . Since  $\text{dist}(L, E_\psi) = 0$  and  $E_\psi$  is convex,  $\text{dist}(x, E_\psi)$  converges to zero as  $x \in L$  goes to infinity. Thus, by making  $L$  shorter if necessary, we have that  $L \subset E_{\phi-2\epsilon} \setminus E_{\phi+2\epsilon}$ . Hence,  $L$  lies out of  $E_{\phi+2\epsilon}$  but the vertical translation of  $L$  for  $4\epsilon$  pushes it in  $E_{\phi+2\epsilon}$ . Since  $E_{\phi+2\epsilon}$ , being a translate of  $E_\phi$ , has BCEH, this contradicts Proposition 3.4 (ii). The contradiction shows that  $E_\psi$  has BCEH as claimed. □

We now give a differential characterization of the BCEH property of an epigraph (3.1).

**Proposition 3.8** *If  $\phi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$  is a convex function of class  $\mathcal{C}^1$  satisfying condition (3.2), then the epigraph  $E = \{(x, y) \in \mathbb{R}^n : y \geq \phi(x)\}$  has BCEH if and only if*

$$\lim_{|x| \rightarrow \infty} |x| \left( 1 - \frac{\phi(x)}{x \cdot \nabla \phi(x)} \right) = +\infty. \tag{3.3}$$

**Proof** We first consider the case  $n = 2$ . Then,  $x$  is a single variable and (3.3) is equivalent to

$$\lim_{x \rightarrow +\infty} \left( x - \frac{\phi(x)}{\phi'(x)} \right) = +\infty \quad \text{and} \quad \lim_{x \rightarrow -\infty} \left( x - \frac{\phi(x)}{\phi'(x)} \right) = -\infty. \tag{3.4}$$

For every  $x \in \mathbb{R}$  such that  $\phi'(x) \neq 0$  the number

$$\xi(x) = x - \frac{\phi(x)}{\phi'(x)}, \tag{3.5}$$

is the first coordinate of the intersection of the tangent line to the graph of  $\phi$  at the point  $(x, \phi(x))$  with the first coordinate axis  $y = 0$ . By (3.2) and convexity of  $\phi$  we have that  $|\phi'(x)|$  is bounded away from zero for all sufficiently big  $|x|$ . This shows that conditions (3.4) are invariant under translations, so we may assume that  $\phi \geq 0$  and  $\phi(0) = 0$ . It is easily seen that the function  $\xi$  is increasing. If  $\phi$  is of class  $\mathcal{C}^2$ , we have that  $\xi'(x) = \phi(x)\phi''(x)/\phi'(x)^2 \geq 0$ .

Assume now that conditions (3.4) hold. Pick a pair of sequences  $a_j < b_j$  in  $\mathbb{R}$  with  $\lim_{j \rightarrow \infty} a_j = -\infty$  and  $\lim_{j \rightarrow \infty} b_j = +\infty$ . The intervals  $I_j = [\xi(a_j), \xi(b_j)]$  then increase to  $\mathbb{R}$  as  $j \rightarrow \infty$ . We identify  $I_j$  with  $I_j \times \{0\} \subset \mathbb{R}^2$ . Since  $\phi$  is convex, its epigraph lies above the tangent line at any point. It follows that the set  $h(E, I_j)$  (see (1.1)) is the bounded region in  $\mathbb{R} \times \mathbb{R}_+$  whose boundary consists of  $I_j$ , the two line segments  $L_j$  and  $L'_j$  connecting the endpoints  $(\xi(a_j), 0)$  and  $(\xi(b_j), 0)$  of  $I_j$  to the respective points  $A_j = (a_j, \phi(a_j))$  and  $B_j = (b_j, \phi(b_j))$  on  $bE$ , and the graph of  $\phi$  over  $[a_j, b_j]$ . The supporting lines of  $L_j$  and  $L'_j$  intersect at a point  $C_j$  in the lower halfspace  $y < 0$ , and we obtain a closed triangle  $\Delta_j$  with the endpoints  $A_j, B_j$ , and  $C_j$ . Note that  $\Delta_j \cap (\mathbb{R} \times \{0\}) = I_j$ . Since  $\phi$  grows at least linearly (see (3.2)), the triangles  $\Delta_j \subset \mathbb{R}^2$  exhaust  $\mathbb{R}^2$  as  $j \rightarrow \infty$ , and the set  $h(E, \Delta_j)$  (1.1) is bounded for every  $j$ . Hence,  $E$  has BCEH. This argument furthermore shows that for any point  $p = (0, -c) \notin E$  there is a unique pair of tangent lines to  $bE$  passing through  $p$  such that, denoting by  $q_1, q_2 \in bE$  the respective points where these lines intersect  $bE$ , the convex hull  $\text{Conv}(E \cup \{p\})$  is the union of  $E$  and the triangle with vertices  $p, q_1, q_2$ .

Conversely, if (3.3) fails then it is easily seen that  $E$  has a boundary ray or an asymptote, so it does not have BCEH. We leave the details to the reader.

The case with  $n \geq 3$  now follows easily. Pick a unit vector  $v \in \mathbb{R}^{n-1}, |v| = 1$ , and let  $L_v$  denote the 2-plane in  $\mathbb{R}^n$  passing through the origin and spanned by  $v$  and  $e_n = (0, \dots, 0, 1)$ . Then,  $E_v := E \cap L_v = \{(t, y) \in \mathbb{R}^2 : y \geq \phi(tv)\}$  and the first condition in (3.4) reads

$$\lim_{t \rightarrow +\infty} \left( t - \frac{\phi(tv)}{\sum_{j=1}^{n-1} v_j \frac{\partial \phi}{\partial x_j}(tv)} \right) = +\infty. \tag{3.6}$$

Writing  $x = tv$  with  $t \geq 0$  and  $v = x/|x|$ , this is clearly equivalent to (3.3). As before, let  $p = (0, \dots, 0, -c) \notin E$ . If (3.3) holds then  $\text{Conv}(E_v \cup \{p\}) \subset L_v$  is obtained by adding to  $E_v$  the triangle in  $L_v$  obtained by the two tangent lines to  $bE_v$  passing through  $p$  as described in the case  $n = 2$ . The sizes of these triangles are uniformly bounded with respect to the direction vector  $|v| = 1$ , and condition (3.2) implies that these triangles increase to  $L_v$  as  $c \rightarrow +\infty$ , uniformly with respect to  $v$ . Since  $\bigcup_{|v|=1} L_v = \mathbb{R}^n$ , Lemma 3.2 shows that

$$\text{Conv}(E \cup \{p\}) = \bigcup_{|v|=1} \text{Conv}(E_v \cup \{p\}),$$

and hence  $E$  has BCEH. The converse is seen as in the special case  $n = 2$ . □

**Corollary 3.9** *If  $\phi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}_+$  is a convex function of class  $\mathcal{C}^1$  such that*

$$\lim_{|x| \rightarrow +\infty} \frac{x \cdot \nabla \phi(x)}{|x|} = +\infty,$$

*then the epigraph  $E = \{(x, y) \in \mathbb{R}^n : y \geq \phi(x)\}$  has BCEH.*

**Proof** By restricting to planes as in the above proof, it suffices to consider the case  $n = 2$ . We may assume that  $\phi \geq 0$  and  $\phi(0) = 0$ . Since  $\phi$  is convex,  $g(x) = \phi'(x)$  is an increasing function and the above condition reads  $\lim_{x \rightarrow \pm\infty} g(x) = \pm\infty$ . For any  $x_0 > 0$  and  $x \geq x_0$  we have that

$$\xi(x) := x - \frac{1}{g(x)} \int_0^x g(t) dt = \int_0^x \left(1 - \frac{g(t)}{g(x)}\right) dt \geq \int_0^{x_0} \left(1 - \frac{g(t)}{g(x)}\right) dt.$$

Letting  $x \rightarrow +\infty$  we have that  $\frac{g(t)}{g(x)} \rightarrow 0$  uniformly on  $t \in [0, x_0]$ , and hence the last integral converges to  $x_0$ . Letting  $x_0 \rightarrow \infty$  we see that  $\lim_{x \rightarrow +\infty} \xi(x) = +\infty$ . The analogous argument applies when  $x \rightarrow -\infty$ . Hence, conditions (3.3) hold and therefore  $E$  has BCEH. □

**Example 3.10** There exist convex epigraphs (3.1) having BCEH where the function  $\phi$  grows linearly, although it cannot be too close to linear near infinity in the absence of boundary rays and asymptotes. We give such an example in  $\mathbb{R}^2$ . Let  $g : \mathbb{R} \rightarrow (-1, 1)$  be an odd, continuous, increasing function with  $\lim_{x \rightarrow +\infty} g(x) = 1$  and  $\int_0^\infty (1 - g(x)) dx = +\infty$ . (An explicit example is  $g(x) = \frac{2}{\pi} \text{Arctan } x$ .) Its integral  $\phi(x) = \int_0^x g(t) dt$  for  $x \in \mathbb{R}$  then clearly satisfies  $\phi(x) \geq 0$ ,  $\phi'(x) = g(x) \in (-1, +1)$  (hence  $\phi$  grows linearly), and  $\phi$  is convex. We now show that (3.3) holds. Let  $x > 0$  be large enough so that  $g(x) > 0$ . We have that

$$\xi(x) = x - \frac{1}{g(x)} \int_0^x g(t) dt = \int_0^x \left(1 - \frac{g(t)}{g(x)}\right) dt.$$

Fix  $x_0 > 0$  and let  $x \geq x_0$ . Then,  $\xi(x) \geq \int_0^{x_0} (1 - g(t)/g(x)) dt$ . Since  $\lim_{x \rightarrow +\infty} g(x) = 1$  and  $\xi$  is increasing for large enough  $|x|$ , it follows that  $\lim_{x \rightarrow +\infty} \xi(x) \geq \int_0^{x_0} (1 - g(t)) dt$ . Sending  $x_0 \rightarrow +\infty$  gives  $\lim_{x \rightarrow +\infty} \xi(x) \geq \int_0^\infty (1 - g(t)) dt = +\infty$ . Similarly we see that  $\lim_{x \rightarrow -\infty} \xi(x) = -\infty$ . Thus, (3.3) holds, and hence the epigraph of  $\phi$  has BCEH.

By using the idea in the above example we now prove the following approximation result, which extends Theorem 1.3 to a much bigger class of convex epigraphs (see Corollary 1.4).

**Proposition 3.11** *Assume that  $\phi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}_+$  is a convex function such that the set  $\{\phi = 0\}$  is nonempty and compact. Given numbers  $\epsilon > 0$  (small) and  $R > 0$  (big) there is a smooth convex function  $\psi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$  such that  $\psi < \phi$  on  $\mathbb{R}^{n-1}$ ,  $\phi(x) - \psi(x) < \epsilon$  for all  $|x| \leq R$ , and the epigraph  $E_\psi = \{y \geq \psi\}$  has BCEH.*

**Proof** By Remark 3.5 the function  $\phi$  grows at least linearly near infinity (see (3.2)). Set

$$A = \liminf_{|x| \rightarrow \infty} \frac{\phi(x)}{|x|} > 0. \tag{3.7}$$

Since the set  $\phi = 0$  does not contain any affine line, Azagra’s result [6, Theorem 1 and Proposition 1] implies that for every  $\epsilon > 0$  there is a smooth strictly convex function  $\psi$  on  $\mathbb{R}^{n-1}$  satisfying  $\phi - \epsilon < \psi < \phi$ . Replacing  $\phi$  by  $\psi - \min_x \psi(x) \geq 0$  we may therefore assume that  $\phi$  is smooth. By increasing the number  $R > 0$  if necessary, we may assume that

$$\frac{\phi(x)}{|x|} \geq \frac{A}{2} \text{ for all } |x| \geq R. \tag{3.8}$$

Pick a number  $r \in (0, 1)$  close to 1 such that

$$(1 - r) \sup_{|x| \leq R} \phi(x) < \epsilon. \tag{3.9}$$

Choose a smooth increasing function  $h : \mathbb{R} \rightarrow \mathbb{R}_+$  such that

$$h(t) = 0 \text{ for } t \leq R, \quad \lim_{t \rightarrow +\infty} h(t) = 1, \quad \text{and} \quad \int_0^\infty (1 - h(t))dt = +\infty.$$

(We can take a smoothing of the Arctan function used in Example 3.10.) Set

$$H(x) = \int_0^{|x|} h(s)ds \text{ for } x \in \mathbb{R}^{n-1}.$$

Clearly,  $H \geq 0$  is a radially symmetric smooth convex function that vanishes on  $|x| \leq R$  and satisfies  $H(x) \leq |x|$  for all  $x \in \mathbb{R}^{n-1}$ . With  $A$  and  $r$  as in (3.7) and (3.9) we set

$$\delta = \frac{A(1 - r)}{2}.$$

We claim that the function

$$\psi(x) = r\phi(x) + \delta H(x) \text{ for } x \in \mathbb{R}^{n-1},$$

satisfies the conditions in the theorem. Clearly,  $\psi \geq r\phi$  is a smooth convex function. For  $|x| \leq R$  we have  $H(x) = 0$ , so  $\psi(x) = r\phi(x) \leq \phi(x)$  and  $\phi(x) - \psi(x) = (1 - r)\phi(x) < \epsilon$  by (3.9). If  $|x| > R$  then  $\phi(x)/|x| \geq A/2$  by (3.8) and  $H(x) < |x|$ , which implies

$$\frac{\psi(x)}{|x|} \leq r \frac{\phi(x)}{|x|} + \delta \leq \frac{\phi(x)}{|x|}.$$

Indeed, we have that  $\frac{\phi(x)}{|x|} - r \frac{\phi(x)}{|x|} = (1 - r) \frac{\phi(x)}{|x|} \geq \frac{A(1-r)}{2} = \delta$ . Hence,  $\psi \leq \phi$  on  $\mathbb{R}^{n-1}$ .

It remains to show that the epigraph  $E_\psi$  satisfies BCEH. We shall verify (3.3), which is equivalent to (3.6) with uniform convergence with respect to the vector  $v = x/|x|$ . Write

$$g_v(t) = r \frac{\partial \phi(tv)}{\partial t}, \quad k(t) = \delta h(t), \quad \tilde{g}_v(t) = \frac{\partial \psi(tv)}{\partial t} = g_v(t) + k(t).$$

The quantity in (3.6) associated to the function  $\psi$  is given by

$$\begin{aligned} \xi_v(t) &= t - \frac{\psi(tv)}{\tilde{g}_v(t)} = \int_0^t \left( 1 - \frac{g_v(s) + k(s)}{g_v(t) + k(t)} \right) ds \\ &= \int_0^t \frac{g_v(t) - g_v(s)}{g_v(t) + k(t)} ds + \int_0^t \frac{k(t) - k(s)}{g_v(t) + k(t)} ds \\ &\geq \int_0^t \frac{g_v(t) - g_v(s)}{g_v(t) + \delta} ds + \int_0^t \frac{k(t) - k(s)}{g_v(t) + \delta} ds, \end{aligned}$$

where the last inequality holds since the functions  $g_v$  and  $k$  are nonnegative and increasing and  $k < \delta$ . Pick  $c > 0$ . We will show that for large enough  $t > 0$  and any unit vector  $v \in \mathbb{R}^{n-1}$  the above expression is bigger than or equal to  $c$ . Choose positive numbers  $t_0, a, t_1$  as follows:

$$t_0 = 3c, \quad a = \max\{3 \max_{|v|=1} g_v(t_0), 3\delta\}, \quad \int_0^{t_1} (k(t_1) - k(s)) ds > ac.$$

Such  $t_1$  exists since  $\lim_{t \rightarrow +\infty} \int_0^t (k(t) - k(s)) ds = \delta \int_0^\infty (1 - h(s)) ds = +\infty$ . Since the integrands in the bound for  $\xi_v(t)$  are nonnegative, we have for  $t \geq \max\{t_0, t_1\}$  and  $|v| = 1$  that

$$\xi_v(t) \geq \int_0^{t_0} \frac{g_v(t) - g_v(s)}{g_v(t) + \delta} ds + \int_0^{t_1} \frac{k(t) - k(s)}{g_v(t) + \delta} ds. \tag{3.10}$$

Assume that for some such  $(t, v)$  we have that  $g_v(t) + \delta \geq a$ . Since  $a \geq 3\delta$ , it follows that  $g_v(t) \geq 2\delta$  and hence

$$\frac{g_v(t)}{g_v(t) + \delta} \geq \frac{2}{3}.$$

Furthermore, from  $a \geq 3 \max_{|v|=1} g_v(t_0)$  we get for  $0 \leq s \leq t_0$  that

$$\frac{g_v(s)}{g_v(t) + \delta} \leq \frac{g_v(t_0)}{a} \leq \frac{1}{3}.$$

These two inequalities imply that the first integral in (3.10) is bounded below by  $t_0/3 \geq c$ . If on the other hand  $g_v(t) + \delta < a$  then the denominator of the second integral in (3.10) is at most  $a$ , so the integral is  $\geq c$  by the choice of  $t_1$ . This shows that  $\xi_v(t) \geq c$  for all  $|v| = 1$  and  $t \geq \max\{t_0, t_1\}$ . Since  $c$  was arbitrary, condition (3.3) holds and hence  $E_\psi$  has BCEH. □



The following observation will be used in the proof of Theorem 1.3.

**Proposition 3.12** *Denote by  $\mathbb{B}$  the open unit ball in  $\mathbb{R}^n$ . Let  $E_\phi \subset \mathbb{R}^n$  be a closed convex set of the form (3.1) with  $\mathcal{C}^1$  boundary having BCEH, where the function  $\phi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$  is bounded from below and strictly convex near infinity. Then there is an  $r_0 > 0$  such that for every  $r \geq r_0$  the convex hull  $\text{Conv}(E_\phi \cup r\overline{\mathbb{B}}) = \{y \geq \psi(x)\}$  is a closed convex set with BCEH, and  $\psi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$  is a convex function of class  $\mathcal{C}^1$  such that  $\psi \leq \phi$  and these functions agree near infinity. Furthermore, if  $r \geq r_0$  is large enough then the function  $\phi_t : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$  defined by*

$$\phi_t(x) = (1 - t)\phi(x) + t\psi(x), \quad x \in \mathbb{R}^{n-1}, \tag{3.11}$$

*is strictly convex for every  $t \in (0, 1)$ , and for any  $0 < t_0 < t_1 < 1$  the closure of the set*

$$\{(x, y) \in \mathbb{R}^n : \phi_{t_1}(x) < y < \phi_{t_0}(x)\},$$

*is a strictly convex cap with the base in the strictly convex hypersurface  $\{y = \phi_{t_0}(x)\}$ .*

**Proof** Consider the function on  $\mathbb{R}^{n-1}$  given by

$$\tilde{\phi}_r(x) = \begin{cases} \min\{\phi(x), -\sqrt{r^2 - |x|^2}\}, & |x| < r, \\ \phi(x), & |x| \geq r. \end{cases}$$

(Note that  $\tilde{\phi}_r$  may be discontinuous at the points of the sphere  $|x| = r$ .) The convex hull of its epigraph  $E_{\tilde{\phi}_r}$  equals  $\text{Conv}(E \cup r\overline{\mathbb{B}})$ , which is closed by Theorem 3.1 (iv), and the set  $h(E, r\overline{\mathbb{B}}) = \text{Conv}(E \cup r\overline{\mathbb{B}}) \setminus E$  is bounded since  $E$  has BCEH. By smoothing  $\tilde{\phi}_r$  we get a function  $\tilde{\psi}_r$  of class  $\mathcal{C}^1$  which agrees with  $\phi$  near infinity such that  $\text{Conv}(E_{\tilde{\psi}_r}) = \text{Conv}(E \cup r\overline{\mathbb{B}})$ . By [28, Theorem 3.2] we conclude that  $\text{Conv}(E \cup r\overline{\mathbb{B}})$  has  $\mathcal{C}^1$  boundary, so it is the epigraph  $E_{\psi_r}$  of a convex function  $\psi_r : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$  of class  $\mathcal{C}^1$  which agrees with  $\phi$  near infinity.

Since  $\phi$  grows at least linearly, there is a function  $\tau(r)$  defined for  $r \in \mathbb{R}_+$  large enough such that  $\psi_r(x) = -\sqrt{r^2 - |x|^2}$  for  $|x| \leq \tau(r)$  and  $\tau(r) \rightarrow +\infty$  as  $r \rightarrow +\infty$ . By choosing  $r$  large enough, the compact set of points where the function  $\phi$  fails to be strictly convex is contained in the ball  $|x| < \tau(r)$ . Since on this ball we have that  $\psi_r(x) = -\sqrt{r^2 - |x|^2}$  which is strictly convex, the convex combinations  $\phi_t$  in (3.11) of  $\phi$  and  $\psi = \psi_r$  are strictly convex on  $\mathbb{R}^{n-1}$  for all  $0 < t < 1$ . For such  $r$ , the last statement in the proposition is evident. (Note that the strictly convex functions  $\rho_t(x, y) = \exp(\psi_t(x) - y) - 1$  for  $t \in (0, 1)$  correspond to those used in Sect. 2.)  $\square$

### 4 Proof of Theorem 1.3

For the definition and the main theorem on Oka manifolds, see [20, Definition 5.4.1 and Theorem 5.4.4]. We shall use the following version of the Oka principle; see [22, Theorem 1.3].

**Theorem 4.1** *Assume that  $X$  is a Stein manifold,  $K$  is a compact  $\mathcal{O}(X)$ -convex set in  $X$ ,  $X'$  is a closed complex subvariety of  $X$ ,  $\Omega$  is an Oka domain in a complex manifold  $Y$ ,  $f : X \rightarrow Y$  is a continuous map which is holomorphic on a neighbourhood of  $K$ ,  $f|_{X'} : X' \rightarrow Y$  is holomorphic, and  $f(X \setminus \mathring{K}) \subset \Omega$ . Then there is a homotopy  $\{f_t\}_{t \in [0,1]}$  of continuous maps  $f_t : X \rightarrow Y$  connecting  $f = f_0$  to a holomorphic map  $f_1 : X \rightarrow Y$  such that for every  $t \in [0, 1]$  the map  $f_t$  is holomorphic on a neighbourhood of  $K$ , it agrees with  $f$  on  $X'$ , it approximates  $f$  uniformly on  $K$  and uniformly in  $t \in [0, 1]$  as closely as desired, and  $f_t(X \setminus \mathring{K}) \subset \Omega$ .*

**Proof of Theorem 1.3** By Proposition 3.4 there are complex coordinates  $z = (z', z_n)$  on  $\mathbb{C}^n$  such that the given set  $E$  is an epigraph of the form (1.2). We shall write  $z = (x, y)$  where  $x = (z', \Re z_n) \in \mathbb{C}^{n-1} \times \mathbb{R} \cong \mathbb{R}^{2n-1}$  and  $y = \Im z_n \in \mathbb{R}$ , so  $E = E_\phi = \{y \geq \phi(x)\}$  where  $\phi \geq 0$  is a convex function as in Proposition 3.4. Let the set  $K \subset X$  and the map  $f_0 : K \rightarrow \mathbb{C}^n$  be as in the theorem; in particular,  $f_0(bK) \subset \mathbb{C}^n \setminus E$ . Thus, there are an open neighbourhood  $U \subset X$  of  $K$  and  $\epsilon > 0$  such that  $f_0$  is holomorphic in  $U$  and  $f_0(U \setminus \mathring{K}) \subset \mathbb{C}^n \setminus E_{\phi-\epsilon}$ . By Azagra [6, Theorem 1.8] there is a real analytic strictly convex function  $\phi_0 : \mathbb{R}^{2n-1} \rightarrow \mathbb{R}$  such that  $\phi - \epsilon < \phi_0 < \phi$ . Its epigraph  $E_0 = \{(x, y) \in \mathbb{C}^n : y \geq \phi_0(x)\}$  is a closed strictly convex set with real analytic boundary which has BCEH by Lemma 3.7, and  $f_0(U \setminus \mathring{K}) \subset \mathbb{C}^n \setminus E_0$ .

Let  $\mathbb{B}$  denote the open unit ball in  $\mathbb{C}^n$  centred at 0. Recall the notation  $h(E, K)$  in (1.1). Pick a number  $r_0 > 0$ . We can find an increasing sequence  $r_k > 0$  diverging to infinity such that

$$\overline{h(E_0, r_k \mathbb{B})} \subset r_{k+1} \mathbb{B} \quad \text{for } k = 0, 1, 2, \dots \tag{4.1}$$

Indeed, since  $E_0$  has BCEH, the set  $h(E_0, r_k \mathbb{B})$  is bounded for each  $k$ , and hence (4.1) holds if the number  $r_{k+1}$  is chosen large enough. Set

$$E_{k+1} = \text{Conv}(E_0 \cup r_k \mathbb{B}) = E_0 \cup h(E_0, r_k \mathbb{B}) \quad \text{for } k = 0, 1, 2, \dots$$

We clearly have that  $E_0 \subset E_1 \subset \dots \subset \bigcup_{k=0}^\infty E_k = \mathbb{C}^n$ . Furthermore, (4.1) shows that for  $j = 0, 1, \dots, k + 1$  we have that  $E_0 \subset E_j \subset E_0 \cup r_{k+1} \mathbb{B}$  and hence

$$E_{k+2} = \text{Conv}(E_j \cup r_{k+1} \mathbb{B}) \quad \text{for } j = 0, 1, \dots, k + 1. \tag{4.2}$$

Proposition 3.12 shows that for each  $k = 1, 2, \dots$  we have  $E_k = \{y \geq \phi_k(x)\}$  where  $\phi_k$  a convex function of class  $\mathcal{C}^1$  which agrees with  $\phi_0$  near infinity, and  $E_k$  has BCEH. Hence,

$$\Omega_k = \mathbb{C}^n \setminus E_k = \{(x, y) \in \mathbb{C}^n : y < \phi_k(x)\},$$

is an Oka domain for every  $k = 0, 1, \dots$  by Corollary 3.6. In view of  $E_{k+2} = \text{Conv}(E_k \cup r_{k+1} \mathbb{B})$  (see (4.2)), Proposition 3.12 also shows that if  $r_{k+1}$  is chosen large enough then the function

$$\psi_t = (1 - t)\phi_k + t\phi_{k+2} : \mathbb{C}^{n-1} \times \mathbb{R} \rightarrow \mathbb{R}, \tag{4.3}$$

is strictly convex for every  $t \in (0, 1)$ , and for each  $0 < t_0 < t_1 < 1$  the closure of the set

$$C = \{(x, y) : \psi_{t_1} < y < \psi_{t_0}\}, \tag{4.4}$$

is a strictly convex cap as described in Sect. 2. (Note that the strictly convex functions  $\rho_t(x, y) = \exp(\psi_t(x) - y) - 1$  for  $t \in (0, 1)$  correspond to those used in Sect. 2.)

Choose an exhaustion  $D_0 \subset D_1 \subset \dots \subset \bigcup_{k=0}^\infty D_k = X$  by smoothly bounded, relatively compact, strongly pseudoconvex domains with  $\mathcal{O}(X)$ -convex closures such that  $K \subset D_0 \subset \bar{D}_0 \subset U$ . For consistency of notation we set  $D_{-1} = K$ . We now construct a sequence of holomorphic maps  $f_k : \bar{D}_k \rightarrow \mathbb{C}^n$  satisfying the following conditions for  $k = 0, 1, 2, \dots$ :

- (a)  $f_k(\overline{D_k \setminus D_{k-1}}) \subset \Omega_k = \mathbb{C}^n \setminus E_k$ ,
- (b)  $f_{k+1}(\overline{D_k \setminus D_{k-1}}) \subset \Omega_k$ , and
- (c)  $f_{k+1}$  approximates  $f_k$  uniformly on  $\bar{D}_{k-1}$  as closely as desired.

For  $k = 0$  the initial map  $f_0$  in Theorem 1.3 satisfies condition (a) while conditions (b) and (c) are void. Assuming inductively that we found maps  $f_0, \dots, f_k$  satisfying these conditions, the construction of the next map  $f_{k+1}$  is made in two steps as follows.

By compactness of the set  $f_k(bD_k) \subset \Omega_k = \{y < \phi_k(x)\}$  we can choose  $t_0 \in (0, 1)$  small enough such that  $f(bD_k) \subset \{y < \psi_{t_0}(x)\}$ , where the function  $\psi_t$  ( $t \in [0, 1]$ ) is given by (4.3). By (4.1) we can also choose  $t_1 \in (t_0, 1)$  sufficiently close to 1 such that

$$E_{k+1} \subset \{(x, y) : y \geq \psi_{t_1}(x)\}.$$

Proposition 2.1 applied to the map  $f_k : \bar{D}_k \rightarrow \mathbb{C}^n$ , the set  $E_k$ , and the strictly convex cap  $C$  (4.4) (which corresponds to  $C_1$  in Proposition 2.1) gives holomorphic map  $g_k : \bar{D}_k \rightarrow \mathbb{C}^n$  approximating  $f_k$  on  $\bar{D}_{k-1}$  and satisfying

$$g_k(bD_k) \subset \{(x, y) : y < \psi_{t_1}(x)\} \subset \mathbb{C}^n \setminus E_{k+1} = \Omega_{k+1} \quad \text{and} \quad g_k(\overline{D_k \setminus D_{k-1}}) \subset \Omega_k. \tag{4.5}$$

In the second step, we use that  $\Omega_{k+1}$  is an Oka domain. Since  $\Omega_{k+1}$  is contractible and  $g_k(bD_k) \subset \Omega_{k+1}$  by (4.5),  $g_k$  extends from  $\bar{D}_k$  to a continuous map  $X \rightarrow \mathbb{C}^n$  sending  $X \setminus D_k$  to  $\Omega_{k+1}$ . Theorem 4.1 applied to  $g_k$  gives a holomorphic map  $f_{k+1} : \bar{D}_{k+1} \rightarrow \mathbb{C}^n$  approximating  $g_k$  on  $\bar{D}_k$  and satisfying  $f_{k+1}(\overline{D_{k+1} \setminus D_k}) \subset \Omega_{k+1}$  (which is condition (a) for  $k + 1$ ) and  $f_{k+1}(\overline{D_k \setminus D_{k-1}}) \subset \Omega_k$  (condition (b)). Since  $f_{k+1}$  approximates  $g_k$  on  $\bar{D}_k$  and  $g_k$  approximates  $f_k$  on  $\bar{D}_{k-1}$ ,  $f_{k+1}$  also satisfies condition (c). This completes the induction step.

If the approximations are close enough then the sequence  $f_k$  converges uniformly on compacts in  $X$  to a holomorphic  $f : X \rightarrow \mathbb{C}^n$ . Conditions (a)–(c) and the fact that the sets  $E_k$  exhaust  $\mathbb{C}^n$  imply that  $f$  is a proper holomorphic map satisfying  $f(X \setminus \mathring{K}) \subset \Omega_0 = \mathbb{C}^n \setminus E_0$ . To construct proper holomorphic immersions and embeddings in suitable dimensions given in the theorem, we use the general position argument at every step to ensure that every map  $f_k$  in the sequence is an immersion or an embedding. (See e.g. [20, Corollary 8.9.3].) If the convergence is fast enough then the same holds for the limit map  $f$  by a standard argument. □

**Proof of Corollary 1.4** Given a holomorphic map  $f_0 : K \rightarrow \mathbb{C}^n$  with  $f_0(bK) \subset \mathbb{C}^n \setminus E_\phi$  as in Theorem 1.3, Proposition 3.11 furnishes a closed convex set  $E_\psi \supset E_\phi$  with BCEH such that  $f_0(bK) \subset \mathbb{C}^n \setminus E_\psi$ . Applying Theorem 1.3 with  $E_\psi$  gives the desired conclusion.  $\square$

We have the following analogue of Theorem 1.3 with interpolation on a closed complex subvariety of  $X$ . Unlike in the above corollary, approximation of  $E$  from the outside by convex sets enjoying BCEH cannot be used since the subvariety  $f(X')$  may have zero distance to  $bE$ . This results extends the case of [24, Theorem 15] when  $E$  is a compact convex set.

**Theorem 4.2** *Let  $E$  be a closed convex set in  $\mathbb{C}^n$  ( $n > 1$ ) with  $\mathcal{C}^1$  boundary which is strictly convex near infinity and has bounded convex exhaustion hulls. Let  $X$  be a Stein manifold,  $K \subset X$  be a compact  $\mathcal{O}(X)$ -convex set,  $U \subset X$  be an open set containing  $K$ ,  $X'$  be a closed complex subvariety of  $X$ , and  $f_0 : U \cup X' \rightarrow \mathbb{C}^n$  be a holomorphic map such that  $f_0|_{X'} : X' \rightarrow \mathbb{C}^n$  is proper holomorphic and  $f_0(bK \cup (X' \setminus K)) \cap E = \emptyset$ . Given  $\epsilon > 0$  there exists a proper holomorphic map  $f : X \rightarrow \mathbb{C}^n$  satisfying the following conditions:*

$$(a) f(X \setminus \overset{\circ}{K}) \subset \mathbb{C}^n \setminus E, \quad (b) \|f - f_0\|_K < \epsilon, \quad (c) f|_{X'} = f_0|_{X'}.$$

*If  $2 \dim X \leq n$  then  $f$  can be chosen an immersion (and an embedding if  $2 \dim X + 1 \leq n$ ) provided that  $f_0|_{X'}$  is one.*

**Proof** This is proved by a small modification of the proof of Theorem 1.3, similar to the one in [24, proof of Theorem 15]. The initial step in the proof, approximating  $E$  from the outside by a strictly convex set, is unnecessary since  $bE$  is strictly convex near infinity. The main (and essentially the only) change comes in the choice of the exhaustion  $D_k$  of the Stein manifold  $X$ . In the inductive step when constructing the map  $f_{k+1}$ , we must assume in addition that  $f_k(bD_k \cap X') \subset \Omega_{k+1} = \mathbb{C}^n \setminus E_{k+1}$ . Then, we push the image of  $bD_k$  out of  $E_{k+1}$  by the same method as before, using Proposition 2.1 but ensuring that the modifications are kept fixed on  $X'$  and small near  $bD_k \cap X'$ . This is possible since the method from [15] is applied locally near  $bD_k$  (away from  $bD_k \cap X'$ ), and these local modifications are glued together by preserving the value of the map on  $X'$ . We refer to [24, proof of Theorem 15] for a more precise description. This gives the next holomorphic map  $f_{k+1} : X \rightarrow \mathbb{C}^n$  satisfying  $f_{k+1}(X \setminus D_k) \subset \Omega_{k+1}$ ,  $f_{k+1}|_{X'} = f_k|_{X'}$ , and conditions (b) and (c) in the proof of Theorem 1.3. We then choose the next domain  $D_{k+1} \subset X$  big enough such that  $f_{k+1}(bD_{k+1} \cap X') \subset \Omega_{k+2} = \mathbb{C}^n \setminus E_{k+2}$ . This is possible since the map  $f_{k+1}|_{X'} = f_0|_{X'} : X' \rightarrow \mathbb{C}^n$  is proper,  $f_0(X' \setminus \overset{\circ}{K}) \subset \Omega = \mathbb{C}^n \setminus E$ , and the domain  $\Omega_{k+2}$  agrees with  $\Omega$  near infinity by the construction. Clearly the induction step is now complete. Assuming that the approximations are close enough, the sequence  $f_k$  converges to a limit holomorphic map  $f : X \rightarrow \mathbb{C}^n$  satisfying the stated conditions.  $\square$

**Acknowledgements** The first named author is supported by Grants P1-0291, J1-3005, and N1-0137 from ARRS, Republic of Slovenia. The second named author is supported by the European Union (ERC Advanced grant HPDR, 101053085) and Grants P1-0291, J1-3005, and N1-0237 from ARRS, Republic of Slovenia.

The authors wish to thank Antonio Alarcón for helpful discussions and information concerning the case pertaining to minimal surfaces.

**Funding** Open access publishing supported by the Slovenian Research Agency and Central Technical Library in Ljubljana.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Acquistapace, F., Broglia, F., Tognoli, A.: A relative embedding theorem for Stein spaces. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **2**(4), 507–522 (1975)
2. Alarcón, A., Drinovec Drnovšek, B., Forstnerič, F., López, F.J.: Every bordered Riemann surface is a complete conformal minimal surface bounded by Jordan curves. *Proc. Lond. Math. Soc.* **111**(4), 851–886 (2015)
3. Alarcón, A., Forstnerič, F., López, F.J.: *Minimal Surfaces from a Complex Analytic Viewpoint*. Springer Monographs in Mathematics, Cham (2021)
4. Alarcón, A., López, F.J.: Minimal surfaces in  $\mathbb{R}^3$  properly projecting into  $\mathbb{R}^2$ . *J. Differ. Geom.* **90**(3), 351–381 (2012)
5. Alexander, H.: Explicit imbedding of the (punctured) disc into  $\mathbb{C}^2$ . *Comment. Math. Helv.* **52**(4), 539–544 (1977)
6. Azagra, D.: Global and fine approximation of convex functions. *Proc. Lond. Math. Soc.* **107**(4), 799–824 (2013)
7. Bishop, E.: Mappings of partially analytic spaces. *Am. J. Math.* **83**, 209–242 (1961)
8. Borell, S., Kutzschebauch, F., Wold, E.F.: Proper holomorphic disks in the complement of varieties in  $\mathbb{C}^2$ . *Math. Res. Lett.* **15**(4), 821–826 (2008)
9. Božin, V.: Note on harmonic maps. *Internat. Math. Res. Not.* **1999**(19), 1081–1085 (1999)
10. Dor, A.: Approximation by proper holomorphic maps into convex domains. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **20**(1), 147–162 (1993)
11. Dor, A.: Immersions and embeddings in domains of holomorphy. *Trans. Am. Math. Soc.* **347**(8), 2813–2849 (1995)
12. Drinovec Drnovšek, B.: Proper holomorphic discs avoiding closed convex sets. *Math. Z.* **241**(3), 593–596 (2002)
13. Drinovec Drnovšek, B.: Proper discs in Stein manifolds avoiding complete pluripolar sets. *Math. Res. Lett.* **11**(5–6), 575–581 (2004)
14. Drinovec Drnovšek, B., Forstnerič, F.: Holomorphic curves in complex spaces. *Duke Math. J.* **139**(2), 203–253 (2007)
15. Drinovec Drnovšek, B., Forstnerič, F.: Strongly pseudoconvex domains as subvarieties of complex manifolds. *Am. J. Math.* **132**(2), 331–360 (2010)
16. Drinovec Drnovšek, B., Forstnerič, F.: Flexible domains for minimal surfaces in Euclidean spaces. *J. Math. Anal. Appl.* **517**, 126653 (2023)
17. Eliashberg, Y., Gromov, M.: Embeddings of Stein manifolds of dimension  $n$  into the affine space of dimension  $3n/2 + 1$ . *Ann. Math.* **136**(1), 123–135 (1992)
18. Fornæss, J.E., Forstnerič, F., Wold, E.: Holomorphic approximation: the legacy of Weierstrass, Runge, Oka-Weil, and Mergelyan. In: *Advancements in Complex Analysis. From Theory to Practice*, pp 133–192. Springer, Cham (2020)
19. Forstnerič, F.: Embedding strictly pseudoconvex domains into balls. *Trans. Am. Math. Soc.* **295**(1), 347–368 (1986)

20. Forstnerič, F.: Stein manifolds and holomorphic mappings (The homotopy principle in complex analysis), volume 56 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge*. Springer, Cham, 2nd edn, (2017)
21. Forstnerič, F.: Holomorphic embeddings and immersions of Stein manifolds: a survey. In: *Geometric Complex Analysis*, volume 246 of *Springer Proceedings in Mathematics and Statistics*. pp. 145–169. Springer, Singapore (2018)
22. Forstnerič, F.: Recent developments on Oka manifolds. *Indag. Math.* **34**(2), 367–417 (2023)
23. Forstnerič, F., Globevnik, J.: Proper holomorphic discs in  $\mathbb{C}^2$ . *Math. Res. Lett.* **8**(3), 257–274 (2001)
24. Forstnerič, F., Ritter, T.: Oka properties of ball complements. *Math. Z.* **277**(1–2), 325–338 (2014)
25. Forstnerič, F., Wold, E.F.: Oka domains in Euclidean spaces. *Int. Math. Res. Not.* (2023). <https://doi.org/10.1093/imrn/rnac347>
26. Gale, D., Klee, V.: Continuous convex sets. *Math. Scand.* **7**, 379–391 (1959)
27. Globevnik, J.: Boundary interpolation by proper holomorphic maps. *Math. Z.* **194**(3), 365–373 (1987)
28. Griewank, A., Rabier, P.J.: On the smoothness of convex envelopes. *Trans. Am. Math. Soc.* **322**(2), 691–709 (1990)
29. Gunning, R.C., Rossi, H.: *Analytic functions of several complex variables*. AMS Chelsea Publishing, Providence, RI, (2009). Reprint of the 1965 original
30. Hakim, M.: Applications holomorphes propres continues de domaines strictement pseudoconvexes de  $\mathbb{C}^n$  dans la boule unité de  $\mathbb{C}^{n+1}$ . (On the extension of proper holomorphic mappings from strictly pseudoconvex domains in  $\mathbb{C}^n$  into the unit ball of  $\mathbb{C}^{n+1}$ ). *Duke Math. J.* **60**(1), 115–133 (1990)
31. Hakim, M., Sibony, N.: Fonctions holomorphes bornées sur la boule unité de  $\mathbb{C}^n$ . *Invent. Math.* **67**, 213–222 (1982)
32. Hoffman, D., Meeks, W.H., III.: The strong halfspace theorem for minimal surfaces. *Invent. Math.* **101**(2), 373–377 (1990)
33. Løw, E.: A construction of inner functions on the unit ball in  $C^p$ . *Invent. Math.* **67**, 223–229 (1982)
34. Løw, E.: Embeddings and proper holomorphic maps of strictly pseudoconvex domains into polydiscs and balls. *Math. Z.* **190**(3), 401–410 (1985)
35. Narasimhan, R.: Imbedding of holomorphically complete complex spaces. *Am. J. Math.* **82**, 917–934 (1960)
36. Remmert, R.: Sur les espaces analytiques holomorphiquement séparables et holomorphiquement convexes. *C. R. Acad. Sci. Paris* **243**, 118–121 (1956)
37. Schoen, R., Yau, S.T.: *Lectures on harmonic maps*. Conference Proceedings and Lecture Notes in Geometry and Topology, II. International Press, Cambridge (1997)
38. Schürmann, J.: Embeddings of Stein spaces into affine spaces of minimal dimension. *Math. Ann.* **307**(3), 381–399 (1997)
39. Stensønes, B.: Proper holomorphic mappings from strongly pseudoconvex domains in  $\mathbb{C}^2$  to the unit polydisc in  $\mathbb{C}^3$ . *Math. Scand.* **65**(1), 129–139 (1989)

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.