



Moving through Cartesian products, coronas and joins in general position

Sandi Klavžar^{a,b,c}, Aditi Krishnakumar^d, Dorota Kuziak^e, Ethan Shallcross^d, James Tuite^{d,f}, Ismael G. Yero^{g,*}

^a Faculty of Mathematics and Physics, University of Ljubljana, Slovenia

^b Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia

^c Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia

^d School of Mathematics and Statistics, Open University, Milton Keynes, UK

^e Departamento de Estadística e IO, Universidad de Cádiz, Algeciras Campus, Spain

^f Department of Informatics and Statistics, Klaipėda University, Lithuania

^g Departamento de Matemáticas, Universidad de Cádiz, Algeciras Campus, Spain

ARTICLE INFO

Article history:

Received 1 May 2025

Received in revised form 16 October 2025

Accepted 22 October 2025

Keywords:

General position set

Mobile general position set

Mobile general position number

Robot navigation

Cartesian product graph

ABSTRACT

The general position problem asks for large sets of vertices such that no three vertices of the set lie on a common shortest path. Recently a dynamic version of this problem was defined, called the *mobile general position problem*, in which a collection of robots must visit all the vertices of the graph whilst remaining in general position. In this paper we investigate this problem in the context of Cartesian products, corona products and joins, giving upper and lower bounds for general graphs and exact values for families including grids, cylinders, Hamming graphs and prisms of trees.

© 2025 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The *general position problem* originated as a geometric puzzle of Dudeney [14], but was first investigated in the context of graph theory in [9,26]. A survey of the problem is given in [7]. The arXiv version [25] of the paper [26] gave the following motivation for the problem. Suppose that a collection of robots is stationed on the vertices of a graph. They communicate with each other by sending signals along shortest paths. To avoid their communication being disrupted, we wish that no robot lies on a shortest path between two other robots. Subject to this condition, what is the greatest possible number of robots that we can place on the graph? In fact, the related *mutual-visibility problem* was initially researched in terms of its applications in robotic navigation and communication (see [2,3,12] for a partial overview) and was only recently considered in a pure mathematics context [13].

However, this picture lacks an important feature of real world robotic navigation: the general position problem is ‘static’, whereas in applications the robots will typically need to move around the network. Watching the mobile delivery robots created by Starship Technologies[®] [1] inspired the authors of [17] to consider a dynamic version of the general position problem, in which robots move through the vertices of a graph whilst remaining in general position, and such that every vertex is visited by a robot at least once (it is assumed that one robot moves to an adjacent vertex at each

* Corresponding author.

E-mail addresses: sandi.klavzar@fmf.uni-lj.si (S. Klavžar), aditi.krishnakumar@open.ac.uk (A. Krishnakumar), dorota.kuziak@uca.es (D. Kuziak), es9854@ou.ac.uk (E. Shallcross), james.tuite@open.ac.uk (J. Tuite), ismael.gonzalez@uca.es (I.G. Yero).




Fig. 1. A (non-optimal) general position set (left) and a gp-set (right) of the Petersen graph are shown in red. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

step and no vertex can contain more than one robot). The paper [17] considered this problem for block graphs, rooted products, Kneser graphs, unicyclic graphs, complete multipartite graphs and line graphs of complete graphs. A mobile version of the closely related mutual-visibility problem was recently treated in [11]. In addition, the paper [4] considers the version of the mobile general position problem with the stricter condition that every vertex must be visited by every robot.

The general position problem has been investigated for a wide variety of graphs, but there is a particularly extensive literature on general position sets in Cartesian products, see [19–21,31,32]. General position sets in other graph products were discussed in [15]. Cartesian products have also been considered in the setting of variants of the general position number, such as the lower general position number [22], the mutual-visibility number [10,13], the monophonic position number [8], the lower mutual-visibility number [5], general position polynomials [16], edge general position numbers [28], total mutual-visibility [24], the variety of general position problems [30] and general position and mutual-visibility colourings [6,18]. In this paper, we examine the mobile general position problem in Cartesian products, together with the coronas and joins of graphs.

The plan of the paper is as follows. In Section 1.1 we introduce the formal definitions of the main concepts that we shall use in our exposition. In Section 2 we give bounds for the mobile general position number of Cartesian products and discuss their sharpness. In Section 3 we determine this number for Cartesian products involving paths, including grids, some cylinders and prisms of trees. Section 4 discusses the mobile general position problem for corona products and joins of graphs. We conclude with some open problems in Section 5.

1.1. Formal definitions and preliminaries

We will write $[n]$ for $\{1, \dots, n\}$ and $[m, n]$ for $\{m, m+1, \dots, n-1, n\}$ when $m \leq n$. A graph $G = (V(G), E(G))$ consists of a set $V(G)$ of vertices that are connected by a set of edges $E(G)$. All graphs that we consider are simple and undirected. We will write $u \sim v$ to indicate that u and v are adjacent in G , and will denote the set of neighbours of u by $N_G(u)$, or simply by $N(u)$ if the graph is clear from the context. The degree $\deg(u)$ of a vertex u is $|N(u)|$. A vertex of degree one is a *leaf*, and the number of leaves in a graph G will be denoted by $\ell(G)$. A complete graph with n vertices will be written as K_n and a complete bipartite graph with partite sets of size n and m as $K_{n,m}$.

A path P_r in G is a sequence $u_1, u_2, \dots, u_r$ of distinct vertices such that $u_i \sim u_{i+1}$ for $i \in [r-1]$ and the length of this path is $r-1$. A cycle C_r is a sequence $u_1, u_2, \dots, u_r$ such that $u_i \sim u_{i+1}$ for $i \in [r-1]$ and also $u_1 \sim u_r$. The distance between vertices $u, v \in V(G)$ is the length of a shortest u, v -path in G . We will typically identify the vertices of a cycle C_n with $\mathbb{Z}_n$ and the vertices of a path P_n with $[n]$ in the natural manner. A subgraph X of G is *convex* if for any $u, v \in V(X)$ we have $V(P) \subseteq V(X)$ for any shortest u, v -path P in G . The subgraph $G[X]$ induced by a subset $X \subseteq V(G)$ is the subgraph with vertex set X such that $u, v \in X$ are adjacent in $G[X]$ if and only if they are adjacent in G .

If $S \subseteq V(G)$ has the property that no three vertices of S lie on a common shortest path of G , then we say that S is a *general position set* of G . The largest possible number of vertices in a general position set of G is the *general position number* of G , denoted by $\text{gp}(G)$. Any largest general position set is referred to as a *gp-set*. Fig. 1 shows two general position sets in the well-known Petersen graph. The red vertices on the left form a maximal but non-optimal general position set of order four, whilst the six red vertices on the right constitute a gp-set of the Petersen graph, and so the Petersen graph has general position number six. Note that in both cases no shortest path between any pair of red vertices passes through a third red vertex.

If a robot is located at a vertex u and $u \sim v$, then we indicate the movement of the robot from u to v along the edge uv by $u \rightsquigarrow v$ and refer to this as a *move*. Suppose that we assign exactly one robot to each vertex of a general position set S . If a robot is stationed at a vertex u of S , then the move $u \rightsquigarrow v$ is called a *legal move* if (i) $v \notin S$ (thereby avoiding having more than one robot per vertex at any stage) and (ii) the new set $(S \setminus \{u\}) \cup \{v\}$ is also a general position set.




Fig. 2. A sequence of legal moves (shown by red arrows) for a Mob_{gp} -set of the Petersen graph. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

A configuration of robots on a general position set of G is called a *mobile general position set* if there is a sequence of legal moves starting from S such that every vertex of G is visited at least once by some robot. The *mobile general position number*, written $\text{Mob}_{\text{gp}}(G)$, is the largest number of robots in a mobile general position set of G . We will refer to a largest possible configuration of robots in mobile general position as a Mob_{gp} -set.

Fig. 2 shows a few configurations of a Mob_{gp} -set for the Petersen graph. The red arrows indicate the legal moves that lead from one configuration to the next. Notice that the last configuration is equivalent to the first up to symmetry. By repeating (symmetrical equivalents of) this sequence of legal moves every vertex of the Petersen graph can be visited. It is shown in [17] that this mobile general position has the largest possible cardinality amongst all such sets, i.e. it is a Mob_{gp} -set.

In [30] the variety of general position problems in graphs was introduced, including general position sets, outer general position sets, dual general position sets, and total general position sets. For our purposes we recall the following definitions. If $X \subseteq V(G)$, then $u, v \in V(G)$ are X -positionable if for any shortest u, v -path P we have $V(P) \cap X \subseteq \{u, v\}$. Hence X is a general position set if all pairs $u, v \in X$ are X -positionable. If it also holds that every pair u, v with $u \in X$ and $v \in V(G) \setminus X$ is X -positionable, then X is an *outer general position set*. The cardinality of a largest outer general position set of G is denoted by $\text{gp}_o(G)$ and is called the *outer general position number*. It is shown in [30] that outer general position sets coincide with sets of mutually maximally distant vertices. (The latter concept was introduced in [29], see also the related survey [23].) In particular, in a block graph the outer general position sets are the sets of simplicial vertices; in the case of a tree this yields $\text{gp}(T) = \text{gp}_o(T) = \ell(T)$.

2. Bounds for Cartesian products

Recall that the *Cartesian product* $G \square H$ of graphs G and H satisfies $V(G \square H) = V(G) \times V(H)$ and $(g, h)(g', h') \in E(G \square H)$ if either $gg' \in E(G)$ and $h = h'$, or $g = g'$ and $hh' \in E(H)$. A G -layer is a subgraph of $G \square H$ induced by $V(G) \times \{h\}$ for some $h \in V(H)$, which will be denoted by G^h , with a similar definition for H -layers gH , where $g \in V(G)$. Likewise, if P is a path $u_1, \dots, u_r$ in G and $h \in V(H)$, then we will denote the path $(u_1, h), (u_2, h), \dots, (u_r, h)$ in $G \square H$ by P^h (with an analogous definition of gQ for a path Q in H and $g \in V(G)$).

We begin by deriving some bounds on $\text{Mob}_{\text{gp}}(G \square H)$. A trivial upper bound is $\text{Mob}_{\text{gp}}(G \square H) \leq \text{gp}(G \square H)$. Proposition 2.1 gives two lower bounds in terms of the mobile and outer general position numbers of the factors.

Proposition 2.1. *For any connected graphs G and H of order at least two, the following hold.*

- (i) $\text{Mob}_{\text{gp}}(G \square H) \geq \max\{\text{Mob}_{\text{gp}}(G), \text{Mob}_{\text{gp}}(H)\}$.
- (ii) $\text{Mob}_{\text{gp}}(G \square H) \geq \max\{\text{gp}_o(G), \text{gp}_o(H)\}$.

Proof. (i) Let S be a Mob_{gp} -set of G and let $h \in V(H)$. We first position the robots at the vertices of the set $S \times \{h\}$. As G^h is a convex subgraph of $G \square H$, robots initially stationed at the vertices of $S \times \{h\}$ can visit every vertex of G^h by a sequence of legal moves, all the time remaining inside G^h and in general position in $G \square H$. Now, whenever a robot visits a vertex (g, h) in this layer, this robot can visit all the vertices in the H -layer gH corresponding to g by a sequence of legal moves and then return to (g, h) . Hence $\text{Mob}_{\text{gp}}(G \square H) \geq \text{Mob}_{\text{gp}}(G)$. By symmetry, $\text{Mob}_{\text{gp}}(G \square H) \geq \text{Mob}_{\text{gp}}(H)$.

(ii) Let $S = \{u_1, \dots, u_r\}$ be an outer general position set of G of cardinality $\text{gp}_o(G)$ and start with robots positioned at each vertex of $S \times \{h\}$ for some $h \in V(H)$. Let R_i be the robot at (u_i, h) for $i \in [r]$. Also, for each $i \in [r]$, let G_i be the connected component containing u_i in $G \setminus (S \setminus \{u_i\})$.

Let $h' \in V(H) \setminus \{h\}$ and let Q be a shortest h, h' -path in H . The robot R_i can follow the path ${}^{u_i}Q$ from (u_i, h) to reach the vertex (u_i, h') by legal moves. At this point, R_i can visit all the vertices of $V(G_i) \times \{h'\}$ by a sequence of legal moves. To see this, notice that the shortest paths between the robots remaining in G^h lie within the layer G^h , whilst the shortest paths from R_i to any R_j , $i \neq j$, do not pass through a third robot by the outer general position property. Afterwards R_i can return to (u_i, h) by performing these legal moves in the reverse order. As this holds for any of the robots and any

$h' \in V(H) \setminus \{h\}$, this allows us to perform a sequence of legal moves so that any vertex of $G \square H$ outside G^h is visited, since $\bigcup_{i=1}^r V(G_i) = V(G)$. Having done this, we return all the robots to their original positions in $S \times \{h\}$ using legal moves.

Finally, let $h' \in N_H(h)$. For $i \in [r]$ we move the robot R_i from (u_i, h) to (u_i, h') in sequence; as S is in general position each of these moves is legal. Afterwards the robots will occupy the set $S \times \{h'\}$. The previous reasoning applied to $G^{h'}$ now shows that the robots can visit each vertex of $V(G) \times \{h\}$ by legal moves. Thus $\text{Mob}_{\text{gp}}(G \square H) \geq \text{gp}_0(G)$ and, by a symmetric argument, $\text{Mob}_{\text{gp}}(G \square H) \geq \text{gp}_0(H)$. $\square$

We now show that both lower bounds in Proposition 2.1 are sharp by considering the Cartesian products $K_r \square P_s$. We recall that [31, Theorem 3.2] implies that $\text{gp}(K_r \square P_s) = r + 1$ for $s \geq 3$. Observe that for $r, s \geq 2$ it holds that $\max\{\text{Mob}_{\text{gp}}(K_r), \text{Mob}_{\text{gp}}(P_s)\} = \max\{\text{gp}_0(K_r), \text{gp}_0(P_s)\} = r$.

Proposition 2.2. For all positive integers $r, s \geq 2$,

$$\text{Mob}_{\text{gp}}(K_r \square P_s) = r.$$

Proof. Let $r, s \geq 2$. By Proposition 2.1 we have $\text{Mob}_{\text{gp}}(K_r \square P_s) \geq r$. We now show that $\text{Mob}_{\text{gp}}(K_r \square P_s) \leq r$. Let $V(P_s) = [s]$ and suppose for a contradiction that there exists a mobile general position set S of $K_r \square P_s$ with $|S| > r$. Then choose $x \in V(K_r)$, such that $(x, i), (x, j) \in V(K_r \square P_s)$ are occupied by robots R_1, R_2 , where $i < j$.

First, notice that for any $y \in V(K_r)$ and $j \leq k \leq s$ the vertex (y, k) cannot be occupied by a robot in the initial configuration S , for otherwise (x, j) would lie on a shortest path between (x, i) and (y, k) . Similarly, every other vertex (y, k) with $k \leq i$ is unoccupied. The same reasoning shows that any other P_s -layer can contain at most one robot, since if there are robots at (y, k) and (y, k') , where $y \in V(K_r) \setminus \{x\}$ and $i < k < k' < j$, then (y, k') would lie on a shortest $(y, k), (x, j)$ -path. Thus, we must have $|S| = r + 1$ and each layer P_s contains exactly one robot for $y \neq x$. Moreover, each of the remaining $r - 1$ robots different from R_1 and R_2 are located at vertices (y, k) such that $i < k < j$.

It now follows that neither R_1 nor R_2 can cross to another P_s -layer by a sequence of legal moves. Otherwise, suppose that robots R_1 and R_2 are stationed at (x, i') and (x, j') respectively just before R_1 moves to a different P_s -layer by the legal move $(x, i') \rightsquigarrow (y, i'), y \in V(K_r) \setminus \{x\}$. Since there is a robot at some vertex (y, k') with $i' < k' < j'$, after this move the robot at (y, k') would lie on a shortest path from (y, i') to (x, j') . As a result, no robot can visit any vertex in $(V(K_r) \setminus \{x\}) \times \{1, s\}$, a contradiction. Thus at most r robots can traverse $K_r \square P_s$ in general position. $\square$

To see that the lower bounds of Proposition 2.1 are independent in general, consider the following examples. If $n \geq 2$, then $\text{Mob}_{\text{gp}}(K_{n,n}) = 2$ and $\text{gp}_0(K_{n,n}) = n$. Hence the bound (i) yields $\text{Mob}_{\text{gp}}(K_{n,n} \square K_{n,n}) \geq 2$, whilst (ii) yields $\text{Mob}_{\text{gp}}(K_{n,n} \square K_{n,n}) \geq n$. On the other hand, if $n \geq 7$, then $\text{Mob}_{\text{gp}}(C_n) = 3$ and $\text{gp}_0(C_n) = 2$, hence the bound (i) is better for $\text{Mob}_{\text{gp}}(C_n \square C_n)$ if $n \geq 7$. Moreover, just after Theorem 2.4 we will demonstrate that the mobile general position number of a graph can be arbitrarily larger than its outer general position number, so that bound (i) can also be arbitrarily larger than bound (ii).

From [15, Theorem 3.2] we recall that if $n \geq 2$ and $m \geq 2$, then $\text{gp}(K_n \square K_m) = n + m - 2$. We now sharpen this result by demonstrating that a gp-set of $K_n \square K_m$ as constructed in [15, Theorem 3.2] is essentially unique as soon as $n \geq 3$ and $m \geq 3$.

Lemma 2.3. Let $n \geq 3, m \geq 3, V(K_n) = [n], V(K_m) = [m]$, and let X be a gp-set of $K_n \square K_m$. Then there exist $i \in [n]$ and $j \in [m]$ such that

$$X = (V((K_n)^j) \cup V((K_n)^i)) \setminus \{(i, j)\}.$$

Proof. Let X be a gp-set of $K_n \square K_m$. Then, as stated above, $|X| = n + m - 2$. Note that the set X contains at most two vertices from every induced copy of C_4 of $K_n \square K_m$. This fact implies the following:


Claim A. if $(i, j), (i, j') \in X$, where $j \neq j'$, then $X \cap (V((K_n)^j) \cup V((K_n)^{j'})) = \{(i, j), (i, j')\}$.

If X contains all the vertices of some K_n -layer, then by Claim A, X contains no other vertices, implying that $|X| = n < n + m - 2$, which is not possible. Let $j \in [m]$ be such that $t = |X \cap V((K_n)^j)|$ is as large as possible. Note that $t \geq 2$, for otherwise we would have $|X| \leq m$. Moreover, by the above, $t \leq n - 1$. If $t = n - 1$ and $i \in [n]$ is the index for which $(i, j) \notin X$, then using Claim A again, we have that $X \subseteq V((K_n)^j) \cup V((K_n)^i)$. Since $|X| = n + m - 2$, we conclude that X has the required structure in this case. Suppose finally that $t = n - k \geq 2$, where $k \geq 2$. We may assume without loss of generality that $X \cap V((K_n)^j) = \{(1, j), \dots, (n - k, j)\}$. Then, using Claim A once more,

$$X \cap \bigcup_{\ell=1}^{n-k} V((K_n)^\ell) = \{(1, j), \dots, (n - k, j)\}.$$

Notice that the subgraph H of $K_n \square K_m$ induced by the vertex set

$$\{n - k + 1, \dots, n\} \times (\{1, \dots, j - 1\} \cup \{j + 1, \dots, m\})$$

Fig. 3. Canonical gp-set in $K_7 \square K_5$.

is isomorphic to $K_k \square K_{m-1}$. Since it is a convex subgraph of $K_n \square K_m$, the intersection $X \cap V(H)$ is a general position set of H . Therefore,

$$|X \cap V(H)| \leq \text{gp}(K_k \square K_{m-1}) = k + (m - 1) - 2 = k + m - 3.$$

By our assumption on t we have $X \cap \{(n - k + 1, j), \dots, (n, j)\} = \emptyset$. Thus we can conclude that $|X| \leq (n - k) + (k + m - 3) = n + m - 3$, a contradiction.

We have thus proved that $|X| = n + m - 2$ holds only in the case when $t = n - 1$ and X has the structure as claimed. $\square$

Lemma 2.3 is illustrated in Fig. 3, where the gp-set of $K_7 \square K_5$ corresponding to the vertex (i, j) is shown. We now use this result to find the mobile general position number of $K_n \square K_m$.

Theorem 2.4. If $n \geq m \geq 1$, then

$$\text{Mob}_{\text{gp}}(K_n \square K_m) = \begin{cases} n; & m \in [2], \\ n + m - 3; & m \geq 3. \end{cases}$$

Proof. If $m = 1$, then $K_n \square K_1 \cong K_n$, thus $\text{gp}(K_n \square K_1) = \text{Mob}_{\text{gp}}(K_n \square K_1) = n$. The case $m = 2$ follows from Proposition 2.2.

Let $m \geq 3$. By Lemma 2.3, every gp-set of $K_n \square K_m$ has the canonical form as illustrated in Fig. 3. It is straightforward to check that no robot placed in such a set can make a legal move. This implies that $\text{Mob}_{\text{gp}}(K_n \square K_m) < \text{gp}(K_n \square K_m) = n + m - 2$.

To complete the proof, we are going to show that there exists a mobile general position set of cardinality $n + m - 3$. Station $n + m - 3$ robots on the vertices of $S = \{(2, 1), (3, 1), \dots, (n, 1)\} \cup \{(1, 3), (1, 4), \dots, (1, m)\}$. This set is a general position set, since it is a subset of the canonical gp-set of $K_n \square K_m$, as illustrated in Fig. 3. Then we perform the following sequence of moves:

- $(2, 1) \rightsquigarrow (2, 2), (3, 1) \rightsquigarrow (3, 2), \dots, (n, 1) \rightsquigarrow (n, 2),$
- $(1, 3) \rightsquigarrow (1, 1).$

Observe that each of these moves is legal. Thus, the new set occupied by the robots is a general position set of $K_n \square K_m$. Next, this process of legal moves can be repeated $m - 2$ times, i.e. for $2 \leq j \leq m - 1$ perform the sequence

$$(2, j) \rightsquigarrow (2, j + 1), (3, j) \rightsquigarrow (3, j + 1), \dots, (n, j) \rightsquigarrow (n, j + 1), (1, j + 2) \rightsquigarrow (1, j - 1),$$

(skipping the final undefined move). In this way, the robots initially positioned at $(2, 1), (3, 1), \dots, (n, 1)$ will visit the vertices of the set $\{2, 3, \dots, n\} \times [m]$ by legal moves, while the remaining vertices can be visited by legal moves by the robots initially positioned at $(1, 3), (1, 4), \dots, (1, m)$. This concludes our argument for the existence of a mobile general position set of cardinality $n + m - 3$. $\square$

By Theorem 2.4 we have $\text{Mob}_{\text{gp}}(K_n \square K_n) = 2n - 3$, whilst it follows from [30, Corollary 4.4] that $\text{gp}_0(K_n \square K_n) = n$, so that the mobile general position number of a graph can be arbitrarily larger than the outer general position number. Theorem 2.4 also demonstrates that the mobile general position number of a Cartesian product can be arbitrarily larger than both bounds in Proposition 2.1.

Finally, we give an example (Cartesian products of stars) that shows that the mobile general position number of a non-trivial Cartesian product can be arbitrarily smaller than its general position number.

Proposition 2.5. For all $r \geq 1$, there exist graphs G, H for which

$$\text{gp}(G \square H) - \text{Mob}_{\text{gp}}(G \square H) = r.$$

Proof. For $k \geq 2$, let $V(K_{1,k}) = \{0\} \cup [k]$, with 0 being the vertex of degree k . It follows from [32, Theorem 1] that $\text{gp}(K_{1,k} \square K_{1,k}) = 2k$. We will show that $\text{Mob}_{\text{gp}}(K_{1,k} \square K_{1,k}) = k + 1$, so that $\text{gp}(K_{1,k} \square K_{1,k}) - \text{Mob}_{\text{gp}}(K_{1,k} \square K_{1,k}) = k - 1$ and the result follows on setting $k = r + 1$.

Let S be any mobile general position set of $K_{1,k} \square K_{1,k}$. Any pair of adjacent vertices in the Cartesian product of trees is a maximal general position set [22], so if $|S| > 2$ the robots must always occupy an independent set. We can start at the stage that a robot is stationed at $(0, 0)$. If there are robots at vertices (u_1, v_1) and (u_2, v_2) with $u_1, v_1, u_2, v_2 \in [k]$, $u_1 \neq u_2$ and $v_1 \neq v_2$, then there would be a shortest $(u_1, v_1), (u_2, v_2)$ -path through $(0, 0)$; hence all the remaining robots lie in a set $\{i\} \times [k]$ or $[k] \times \{j\}$ for some $i, j \in [k]$. In either case, we have $|S| \leq k + 1$.

Finally, we show that $\text{Mob}_{\text{gp}}(K_{1,k} \square K_{1,k}) \geq k + 1$. Consider the set $S = \{(0, 0)\} \cup ([k] \times \{1\})$. First, move $(0, 0) \rightsquigarrow (0, 2)$, followed by $(i, 1) \rightsquigarrow (i, 0)$ for $i \in [2, k]$ and then $(0, 2) \rightsquigarrow (1, 2)$. By relabelling as necessary, we see that each vertex except for $(0, 1)$ may be visited using such a sequence. For the remaining vertex, from the initial configuration $S = \{(0, 0)\} \cup ([k] \times \{1\})$ perform $(0, 0) \rightsquigarrow (0, 2)$, $(i, 1) \rightsquigarrow (i, 0)$ for $i \in [2, k]$, and finally $(1, 1) \rightsquigarrow (0, 1)$. Therefore, S is a mobile general position set. $\square$

3. Cartesian products with paths

In this section we continue our exposition with exact values of the mobile general position number for some Cartesian products involving paths, including prism graphs, i.e. products $G \square P_2$. The result for prisms of complete graphs is contained in Proposition 2.2 and Theorem 2.4. We begin with the exact value of $\text{Mob}_{\text{gp}}(T \square K_2)$, where T is a tree. It follows from Proposition 2.1 that for any graph G the mobile general position number of a prism satisfies $\text{Mob}_{\text{gp}}(G) \leq \text{Mob}_{\text{gp}}(G \square K_2) \leq 2 \text{gp}(G)$. Since the mobile general position number of a tree is just two, Theorem 3.1 shows that $\text{Mob}_{\text{gp}}(G \square K_2)$ can be arbitrarily larger than $\text{Mob}_{\text{gp}}(G)$.

Theorem 3.1. For any tree T with order at least three, $\text{Mob}_{\text{gp}}(T \square K_2) = \ell(T)$.

Proof. By Proposition 2.2, we can assume that $\ell(T) \geq 3$. As remarked in Section 1, $\text{gp}_0(T) = \ell(T)$, so by Proposition 2.1(ii) we have $\text{Mob}_{\text{gp}}(T \square K_2) \geq \ell(T)$. We label the vertices of K_2 by 0,1. Suppose that at least $\ell(T) + 1$ robots can traverse $T \square K_2$ in general position. Since $\text{gp}(T) = \ell(T)$ and each T -layer in $T \square K_2$ is a convex subgraph, neither T -layer can contain $> \ell(T)$ robots at any stage, that is, each layer $V(T) \times \{0\}$ and $V(T) \times \{1\}$ must contain at least one robot at any time. Trivially we can assume that at least one layer contains two or more robots.

Suppose that the layer T^1 contains at least two robots. If not all of the robots in T^1 are already stationed at leaves of T , then we may suppose that a robot is at a vertex $(u, 1)$, where u is a cut-vertex of T . Let $T_1, \dots, T_k$ be the components of $T - u$. As the set of robots is in general position, one of the sets $V(T_i) \times \{1\}$ must contain the remaining robots of T^1 ; without loss of generality, suppose that these other robots are in $V(T_1) \times \{1\}$. Considering the shortest paths to the robots in $V(T_1) \times \{1\}$, we see that there cannot be robots positioned at any vertex from $(\{u\} \cup \bigcup_{i=2}^k V(T_i)) \times \{0\}$. Therefore, the robot at $(u, 1)$ can be moved by a sequence of legal moves to $(v, 1)$, where v is a leaf of T lying in T_2 . In this fashion, if both layers T^i , $i \in \{0, 1\}$, contained at least two robots, then all of these robots could be moved to vertices corresponding to leaves of T . However, if w is any leaf of T , then we cannot have robots at both $(w, 0)$ and $(w, 1)$, as this constitutes a maximal general position set of $T \square K_2$. Therefore, in this case, we conclude that there are at most $\ell(T)$ robots in $T \square K_2$, a contradiction.

It follows that there must be a layer, say T^0 , that contains just one robot R , and $\ell(T)$ robots lie in T^1 , which we can assume to start at the leaves of T^1 . By the preceding argument, R cannot move to the layer T^1 and no robot in T^1 can move to T^0 . If T is a path P_n , then we are left with three robots: two positioned at vertices corresponding to leaves of the layer P_n^1 , and the third robot located at an internal vertex of the path layer P_n^0 . It is now readily observed that no robot can visit the vertices corresponding to leaves of P_n^0 . Hence, we may assume that T is not a path. Let z be any vertex of T with degree at least three. No robot in T^1 can visit $(z, 1)$ without creating three-in-a-line within T^1 , and robot R cannot leave T^0 to visit $(z, 1)$, a contradiction. We conclude that $T \square K_2$ can hold at most $\ell(T)$ robots. $\square$

By Theorem 3.1 we have $\text{Mob}_{\text{gp}}(P_n \square P_2) = 2$ for $n \geq 2$. We next complement this result by considering products of two paths each of order at least three.

Theorem 3.2. If $n, m \geq 3$, then $\text{Mob}_{\text{gp}}(P_n \square P_m) = 3$.

Proof. Let $V(P_k) = [k]$, so that $V(P_n \square P_m) = [n] \times [m]$. Consider an arbitrary general position set S of $P_n \square P_m$ with $|S| = 4$. Then from the proof of [19, Theorem 2.1] we deduce that none of the corner vertices $(1, 1)$, $(1, m)$, $(n, 1)$ and (n, m) belongs to S . Hence, no sequence of legal moves for any configuration of four robots in general position in $P_n \square P_m$ can visit any of the vertices $(1, 1)$, $(1, m)$, $(n, 1)$ and (n, m) . Thus, $\text{Mob}_{\text{gp}}(P_n \square P_m) \leq 3$.

To prove that $\text{Mob}_{\text{gp}}(P_n \square P_m) \geq 3$, we start with three robots positioned at the general position set $S = \{(1, 1), (n, 1), (2, m)\}$. We next describe a sequence of legal moves for the three robots.

- $(1, 1)$ moves to all the vertices from $\{1\} \times [m - 1]$ and returns back to $(1, 1)$.
- $(n, 1)$ moves to all the vertices from $\{n\} \times [m - 1]$ and returns back to $(n, 1)$.




Fig. 4. The legal moves in the infinite grid.

- $(2, m)$ moves to all the vertices from $[2, n - 1] \times [2, m]$ and returns back to $(2, m)$.
- $(n, 1) \rightsquigarrow (n, 2)$. After this, $(1, 1)$ moves to vertices $(2, 1), \dots, (n - 1, 1)$. Notice that at this point the robots are at vertices $(2, m), (n - 1, 1)$ and $(n, 2)$. Moreover, by this stage, all the vertices apart from $(1, m)$ and (n, m) have already been visited.
- $(2, m) \rightsquigarrow (1, m)$ and $(n, 2) \rightsquigarrow (n, 3) \rightsquigarrow \dots \rightsquigarrow (n, m)$.

Notice that all these moves are legal, which demonstrates that $\text{Mob}_{\text{gp}}(P_n \square P_m) \geq 3$ and hence $\text{Mob}_{\text{gp}}(P_n \square P_m) = 3$ when $n, m \geq 3$. $\square$

By contrast, for infinite grids $P_\infty \square P_\infty$ we have equality with the general position number.

Theorem 3.3. *If P_∞ is the two-way infinite path, then $\text{Mob}_{\text{gp}}(P_\infty \square P_\infty) = 4$.*

Proof. We first recall from [27, Corollary 3.2] that $\text{gp}(P_\infty \square P_\infty) = 4$. Hence, it remains to show that $\text{Mob}_{\text{gp}}(P_\infty \square P_\infty) \geq 4$. To do so, set $V(P_\infty) = \mathbb{Z}$ and let $(i, j) \in V(P_\infty \square P_\infty)$. We claim that the set $N(i, j) = \{(i - 1, j), (i + 1, j), (i, j - 1), (i, j + 1)\}$ is a mobile general position set of $P_\infty \square P_\infty$. First, from the proof of [27, Corollary 3.2], we know that any such set $N(i, j)$ is in general position. Next, observe that the sequence of moves $(i + 1, j) \rightsquigarrow (i + 2, j), (i, j + 1) \rightsquigarrow (i + 1, j + 1), (i, j - 1) \rightsquigarrow (i + 1, j - 1)$ and $(i - 1, j) \rightsquigarrow (i, j)$ is a sequence of legal moves from robots positioned at the set $N(i, j)$. This sequence moves the four robots one coordinate to the right, leaving robots at $N(i + 1, j)$. Fig. 4 shows a gp-set of the grid $P_\infty \square P_\infty$, and one round of moves as just described. The order of the moves is shown by the numeric order in the figure. We can analogously move the four robots in each of the remaining three directions in the infinite grid. In this way, every vertex of the infinite grid is eventually occupied by some robot. Thus, the conclusion follows. $\square$

We now find the exact value of the mobile general position number for some cylinder graphs $C_r \square P_s$. The general position numbers of the cylinder graphs are given in [19] as

$$\text{gp}(C_r \square P_s) = \begin{cases} 3; & r = 3, s = 2, \\ 5; & r = 7 \text{ or } r \geq 9, \text{ and } s \geq 5, \\ 4; & \text{otherwise.} \end{cases}$$

Note that Proposition 2.2 gives $\text{Mob}_{\text{gp}}(C_3 \square P_s) = 3$ for $s \geq 2$. We begin with the prism graphs $C_r \square P_2$.



Theorem 3.4. *If $n \geq 3$, then*

$$\text{Mob}_{\text{gp}}(C_n \square K_2) = \begin{cases} 3; & n = 3, \\ 2; & n = 4, \\ 4; & \text{otherwise.} \end{cases}$$

Proof. The case $C_3 \square K_2 = K_3 \square K_2$ has already been treated above. Up to symmetry, there are unique general position sets of $C_4 \square K_2$ of cardinalities three and four, both of which are independent sets. However, in both cases no robot can move whilst maintaining the independence property, so that $\text{Mob}_{\text{gp}}(C_4 \square K_2) \leq 2$, and the equality trivially holds.

We assume for the remainder of the proof that $n \geq 5$. It follows from [19, Theorem 3.2] that $\text{gp}(C_n \square K_2) = 4$. Set $V(C_n) = \{v_i : i \in \mathbb{Z}_n\}$ and $V(K_2) = [2]$. We separate the argument into two cases.

Case 1: n is odd.

Fig. 5. Moving robots in $C_5 \square K_2$.Fig. 6. Moving robots in $C_6 \square K_2$.

Consider a set of four robots located at $S = \{(v_0, 1), (v_{\lceil n/2 \rceil}, 1), (v_1, 2), (v_{\lceil n/2 \rceil+1}, 2)\}$. Then S is a general position set. Moreover, consider the following sequence of four moves for the robots:

- $(v_1, 2) \rightsquigarrow (v_2, 2)$;
- $(v_0, 1) \rightsquigarrow (v_1, 1)$;
- $(v_{\lceil n/2 \rceil+1}, 2) \rightsquigarrow (v_{\lceil n/2 \rceil+2}, 2)$;
- $(v_{\lceil n/2 \rceil}, 1) \rightsquigarrow (v_{\lceil n/2 \rceil+1}, 1)$.

Fig. 5 shows this process for the case $C_5 \square K_2$.

Case 2: n is even.

Suppose now that the robots are positioned at $S = \{(v_0, 1), (v_{n/2}, 1), (v_1, 2), (v_{n/2+1}, 2)\}$. Again S is a general position set. Moreover, consider the following sequence of moves:

- $(v_1, 2) \rightsquigarrow (v_2, 2)$;
- $(v_{n/2+1}, 2) \rightsquigarrow (v_{n/2+2}, 2)$;
- $(v_0, 1) \rightsquigarrow (v_1, 1)$;
- $(v_{n/2}, 1) \rightsquigarrow (v_{n/2+1}, 1)$.

Fig. 6 shows this process for the case $C_6 \square K_2$.

In both cases above, we note that these four moves are legal. Since the obtained sets are symmetric with respect to the original ones, by repeating these procedures the robots will eventually visit all the vertices of $C_n \square K_2$. It follows that each S is a mobile general position set, and hence $\text{Mob}_{\text{gp}}(C_n \square K_2) \geq 4$. $\square$

We now introduce a technical lemma that allows us to extend results on short cylinders to longer cylinders.

Lemma 3.5. *If H is a connected graph with girth at least $2r$ and radius at least $r - 1$, then for any graph G it holds that*

$$\text{Mob}_{\text{gp}}(G \square H) \geq \text{Mob}_{\text{gp}}(G \square P_r).$$

Proof. Let Q be any path $v_1, v_2, \dots, v_r$ of length $r - 1$ in H . The subgraph of $G \square H$ induced by $V(G) \times V(Q)$ is isomorphic to $G \square P_r$, and as the girth of H is at least $2r$, the subgraph is convex. Thus, $\text{Mob}_{\text{gp}}(G \square P_r)$ robots can traverse the vertices of $V(G) \times V(Q)$ in general position without leaving the subgraph. Now, let $v_{r+1} \in N_H(v_r) \setminus \{v_{r-1}\}$. Suppose that the robots have visited all the vertices of $V(G) \times V(Q)$ by a sequence of legal moves. Next, move all robots in $V(G) \times \{v_r\}$ to $V(G) \times \{v_{r+1}\}$ in turn by legal moves of the form $(u, v_r) \rightsquigarrow (u, v_{r+1})$, where $u \in V(G)$. Then repeat this process to move the robots in $V(G) \times \{v_j\}$ to $V(G) \times \{v_{j+1}\}$ for $j = r - 1, r - 2, \dots, 1$. As H has girth at least $2r$, the robots remain in general position throughout this process. As H is connected and the radius of H is at least $r - 1$, any layer G^h can be visited in this way. $\square$

Notice that Lemma 3.5 generalises the inequality $\text{Mob}_{\text{gp}}(G) \leq \text{Mob}_{\text{gp}}(G \square K_2)$. We first focus on cylinders with cycles of length four.

Proposition 3.6. *If $s \geq 3$ is an integer, then $\text{Mob}_{\text{gp}}(C_4 \square P_s) = 3$.*

Proof. Let $s \geq 3$, $V(C_4) = \mathbb{Z}_4$, and $V(P_s) = [s]$. We first show that $\text{Mob}_{\text{gp}}(C_4 \square P_s) \leq 3$. Suppose for a contradiction that there exists a mobile general position set S of $C_4 \square P_s$ with $|S| \geq 4$.

Clearly, no three robots from S can lie in the same P_s -layer. Suppose that two robots R_1 and R_2 in S lie in the same P_s -layer of $C_4 \square P_s$; without loss of generality, R_1 and R_2 are stationed at vertices $(0, i)$ and $(0, j)$ respectively, where $1 \leq i < j \leq s$. There is a shortest path from any vertex (u, v) with $v \in [i]$ to R_2 through R_1 , and likewise for vertices with second coordinate at least j . Hence, there are no robots on $\mathbb{Z}_4 \times ([1, i] \cup [j, s])$ apart from R_1 and R_2 . Call the other two robots R_3 and R_4 . By the above analysis, R_3 and R_4 cannot have the same first coordinate. Hence we can assume that robot R_3 lies at $(1, k)$, where $i < k < j$. Any vertex in $(\{1, 2\} \times [i + 1, j - 1]) \setminus \{(1, k)\}$ has a shortest path to either R_1 or R_2 through R_3 , so R_4 must be at a vertex $(3, l)$, where $i < l < j$. However, in this configuration each robot is only free to move within its P_s -layer, and so no robot can visit any vertex in 2P_s , a contradiction. Consequently, there is one robot on each P_s -layer, and none of the robots can move to another P_s -layer.

Observe that any pair of adjacent vertices constitutes a maximal general position set of $C_4 \square P_s$, so the robots must at all times occupy an independent set. Therefore, if we suppose that R_1 is the robot located at the vertex with smallest second coordinate in the initial configuration, say at vertex $(0, i)$, then no robot in the layers 1P_s or 3P_s can ever move to a position with second coordinate smaller than that of R_1 , and hence the vertices in $\{1, 3\} \times \{1\}$ cannot be visited by legal moves. Thus, $\text{Mob}_{\text{gp}}(C_4 \square P_s) \leq 3$.

To show the lower bound, consider $C_4 \square P_3$. We start with robots at vertices $(0, 1)$, $(1, 2)$ and $(0, 3)$. Then for $i = 0, 1, 2$ in succession we perform the sequence of three legal moves $(i + 1, 2) \rightsquigarrow (i + 2, 2)$, $(i, 3) \rightsquigarrow (i + 1, 3)$ and $(i, 1) \rightsquigarrow (i + 1, 1)$ in this order. Lemma 3.5 now gives the result for cylinders $C_4 \square P_s$ for $s \geq 4$. $\square$

Theorem 3.4 for prisms along with Lemma 3.5 implies that $\text{Mob}_{\text{gp}}(C_r \square P_s) \geq 4$ for $r \geq 5$, $s \geq 2$. Combined with the upper bound involving $\text{gp}(C_r \square P_s)$ we see that $\text{Mob}_{\text{gp}}(C_r \square P_s) = 4$ for $r \in \{5, 6, 8\}$ and $s \geq 2$, or for $r \geq 5$ and $s \leq 4$. In the cases $r = 7$ or $r \geq 9$ and $s \geq 5$ the mobile general position number of $C_r \square P_s$ must be either four or five.

Proposition 3.7. *If $r = 9$ or $r \geq 11$, and $s \geq 5$, then $\text{Mob}_{\text{gp}}(C_r \square P_s) = 5$.*

Proof. It is known from [19, Theorem 3.2] that $\text{gp}(C_r \square P_s) = 5$ for $r = 9$ or $r \geq 11$, and $s \geq 5$. It only remains to show that $\text{Mob}_{\text{gp}}(C_r \square P_s) \geq 5$. We show that the result is true for $C_r \square P_s$, and the full claim then follows for larger values of s by Lemma 3.5. Set $V(C_r) = \mathbb{Z}_r$ and $V(P_s) = [5]$.

If $r \geq 11$ and $i \in \mathbb{Z}_r$, then we consider the set

$$S_{i,0} = \{(i + 1, 1), (i + 4, 2), (i + \lfloor r/2 \rfloor + 2, 3), (i, 4), (i + 3, 5)\}.$$

For $i = 0, 1, \dots, r - 2$ we define the following sequence of five moves:

- $(i + 4, 2) \rightsquigarrow (i + 5, 2)$ to give $S_{i,1}$,
- $(i + 3, 5) \rightsquigarrow (i + 4, 5)$ to give $S_{i,2}$,
- $(i + \lfloor r/2 \rfloor + 2, 3) \rightsquigarrow (i + \lfloor r/2 \rfloor + 3, 3)$ to give $S_{i,3}$,
- $(i + 1, 1) \rightsquigarrow (i + 2, 1)$ to give $S_{i,4}$,
- $(i, 4) \rightsquigarrow (i + 1, 4)$.

The final move brings us to the configuration $S_{i+1,0}$. We start with the five robots positioned at the set $S_{0,0}$ and perform these sequences of moves for $i = 0, 1, \dots, r - 2$. Each of these moves is legal. To see this, notice that the robots remain in general position at each stage, which is easily verified for $S_{0,j}$, $j \in [5]$, and by then observing that the automorphism that maps (u, v) to $(u + i, v)$ for all $u \in \mathbb{Z}_r$, $s \in [5]$ transforms $S_{0,j}$ to $S_{i,j}$ for $j \in [5]$. Moreover, by the end of the process, all vertices have been visited.

Similarly, for $r = 9$ we start with robots positioned at the set

$$\{(1, 1), (4, 2), (\lfloor s/2 \rfloor + 2, 3), (0, 4), (3, 5)\}$$

and perform the sequence of moves:

- $(\lfloor s/2 \rfloor + 2, 3) \rightsquigarrow (\lfloor s/2 \rfloor + 3, 3)$,
- $(1, 1) \rightsquigarrow (2, 1)$,
- $(0, 4) \rightsquigarrow (1, 4)$,
- $(4, 2) \rightsquigarrow (5, 2)$,
- $(3, 5) \rightsquigarrow (4, 5)$.

By repeating these moves the robots visit all of the vertices of $C_9 \square P_5$ by legal moves. $\square$

Hence, the only unknown values are $\text{Mob}_{\text{gp}}(C_7 \square P_s)$ and $\text{Mob}_{\text{gp}}(C_{10} \square P_s)$ for $s \geq 5$. We conjecture that the answer is four in these cases.

By combining Lemma 3.5 with Proposition 3.7 we obtain a lower bound for the mobile general position number of sufficiently large torus graphs.

Corollary 3.8. For $r = 9$ or $r > 10$, and $s \geq 10$, $\text{Mob}_{\text{gp}}(C_r \square C_s) \geq 5$.

It is shown in [21] that if $r, s \geq 7$ and r and s do not both lie in $\{8, 10, 12\}$, then $\text{gp}(C_r \square C_s) = 7$. Computer search shows that the torus $C_9 \square C_8$ has mobile general position number seven, so this upper bound can be achieved.

4. Corona products and joins

In this section we consider moving robots in general position through corona products and joins. We first define these two graph operations.

Given two graphs G and H with $V(G) = \{v_1, \dots, v_n\}$, the *corona product* graph $G \odot H$ is formed by taking one copy of G and n disjoint copies of H , call them $H^1, \dots, H^n$, and for each $i \in [n]$ adding all the possible edges between $v_i \in V(G)$ and every vertex of H^i . For $i \in [n]$ we will write $\tilde{H}_i$ for the subgraph of $G \odot H$ induced by $V(H^i) \cup \{v_i\}$. Also, the *join* $G \vee H$ of graphs G and H is obtained from the disjoint union of G and H by adding all possible edges between G and H .

4.1. Corona product graphs

The first paper [17] on the mobile general position problem briefly considered mobile general position sets in rooted products. This suggests investigating the problem in corona products, which can also be viewed as a kind of rooted product. The general position number of corona product graphs was studied in [15]. We now bound the value of the mobile general position number of the corona product $G \odot H$.

Theorem 4.1. For any two graphs G and H ,

$$\max\{\text{Mob}_{\text{gp}}(G), \text{Mob}_{\text{gp}}(H \vee K_1)\} \leq \text{Mob}_{\text{gp}}(G \odot H) \leq \max\{n(G), \text{gp}(H \vee K_1)\}.$$

Proof. Let S be a mobile general position set of $H \vee K_1$ and let S_1 be its copy in $\tilde{H}_1$. We claim that S_1 is a mobile general position set of $G \odot H$. As H_1 is an isometric subgraph of $G \odot H$, first the robots from S_1 can visit each vertex of H_1 . Next, as soon as one robot visits the vertex v_1 , this robot can visit all the vertices of $V(G \odot H) \setminus V(\tilde{H}_1)$ before returning to v_1 . It follows that $\text{Mob}_{\text{gp}}(G \odot H) \geq \text{Mob}_{\text{gp}}(H \vee K_1)$.

Now, let S be a mobile general position set of G and let S' be the copy of S in $G \odot H$. Then each vertex $v_i \in V(G)$ can be visited by a robot from S' . Moreover, as soon as a robot moves to some vertex v_i , this robot can visit all the vertices from H^i and then return to v_i . Hence S' is a mobile general position set of $G \odot H$ and $\text{Mob}_{\text{gp}}(G \odot H) \geq \text{Mob}_{\text{gp}}(G)$. We conclude that $\text{Mob}_{\text{gp}}(G \odot H) \geq \max\{\text{Mob}_{\text{gp}}(H \vee K_1), \text{Mob}_{\text{gp}}(G)\}$.

To prove the upper bound, let S be a mobile general position set of $G \odot H$. If $|S| \leq n(G)$, then there is nothing to prove. Assume next that $|S| \geq n(G) + 1$. Then by the pigeonhole principle we have $|S \cap V(\tilde{H}_i)| \geq 2$ for some $i \in [n(G)]$. Hence either at some point there is already a robot in H^i and a second robot enters $\tilde{H}_i$ via v_i , or else there are always at least two robots in $V(\tilde{H}_i)$ and a further robot must visit v_i . Denote the positions of the robots at this moment by S' . In S' there is a robot R_1 in $V(H^i)$ and a robot R_2 at the cut-vertex v_i . As any path from R_1 to a robot on $V(G \odot H) \setminus V(\tilde{H}_i)$ would pass through R_2 , it follows that $S' \subseteq V(\tilde{H}_i)$ and S' is a general position set of $\tilde{H}_i$. Hence, under the assumption that $|S| \geq n(G) + 1$, we must have $\text{Mob}_{\text{gp}}(G \odot H) = |S'| \leq \text{gp}(H \vee K_1)$. $\square$

When both G and H are complete graphs, $G \odot H$ is a block graph, hence the following consequence can also be deduced from [17, Theorem 2.3].

Corollary 4.2. If $r, s \geq 1$, then $\text{Mob}_{\text{gp}}(K_r \odot K_s) = \max\{r, s + 1\}$.

Note that Corollary 4.2 demonstrates the sharpness of all the bounds in Theorem 4.1. For another sharpness example, in which the upper and lower bounds do not coincide, consider $G = K_2$ and $H = C_4$. Then we have that $\text{Mob}_{\text{gp}}(K_2) = 2$, $\text{Mob}_{\text{gp}}(C_4 \vee K_1) = 2$, and $\text{gp}(C_4 \vee K_1) = 3$. It can be noted that $\text{Mob}_{\text{gp}}(K_2 \odot C_4) = 3 = \text{gp}(C_4 \vee K_1)$. For another infinite family, let G be an arbitrary tree, and H the edgeless graph of order at least two. Since in this case $G \odot H$ is a tree, we have $\text{Mob}_{\text{gp}}(G \odot H) = 2$ (see [17, Theorem 2.3]), which is also the value of the lower bound in Theorem 4.1.

We complete this section by presenting a result which shows that none of the bounds of Theorem 4.1 is sharp in general.

Theorem 4.3. If $n \geq 3$, then $\text{Mob}_{\text{gp}}(C_n \odot K_1) = \lceil \frac{n}{2} \rceil + 1$.

Proof. Set $V(C_n) = \mathbb{Z}_n$ and, for each $i \in \mathbb{Z}_n$, let i' be the leaf in $C_n \odot K_1$ attached to i . We first show that $\text{Mob}_{\text{gp}}(C_n \odot K_1) \geq \lceil \frac{n}{2} \rceil + 1$. Set $Y_i = \{i', (i + 1)', \dots, (\lceil \frac{n}{2} \rceil + i)'\}$ for any $i \in \mathbb{Z}_n$. We claim that Y_0 is a mobile general position set. It is easily seen that each Y_i is a general position set of $C_n \odot K_1$.

Starting from some fixed Y_i , we move the robot R placed at $(\lceil \frac{n}{2} \rceil + i)'$ through the vertices of the set $[(\lceil \frac{n}{2} \rceil + i), (i - 1)]$ and their attached leaves, and finally leave the robot at $(i - 1)'$. After these moves we are left with the robots occupying the set Y_{i-1} and at each stage the robots remained in general position, since the paths between robots in $Y_i \setminus \{(\lceil \frac{n}{2} \rceil + i)'\}$

pass through $[i, \lceil \frac{n}{2} \rceil + i - 1]$. Therefore starting at Y_0 and repeating this process for $i = 0, -1, \dots, -\lceil \frac{n}{2} \rceil$ results in all vertices being visited by a robot and Y_0 is a mobile general position set as claimed.

We now prove that $\text{Mob}_{\text{gp}}(C_n \odot K_1) \leq \lceil \frac{n}{2} \rceil + 1$. Note that each set $\{i, i'\}$, $i \in \mathbb{Z}_n$, is a maximal general position set, so we must have $|X \cap \{i, i'\}| \leq 1$ for each $i \in \mathbb{Z}_n$ and any Mob_{gp} -set X . Consider the moment that a robot visits the vertex 0. Let j, k be the smallest and largest values in $[1, n-1]$, respectively, such that there is a robot in $\{j, j'\}$ and $\{k, k'\}$. To avoid the robot in $\{k, k'\}$ having a shortest path to the robot in $\{j, j'\}$ through the robot at 0 we must have $k - j \leq \lceil \frac{n}{2} \rceil - 1$. As each set $\{i, i'\}$ contains at most one robot for $i \in [j, k] \cup \{0\}$ and no robots for $i \in [k+1, n-1] \cup [1, j-1]$, this gives an upper bound of $\lceil \frac{n}{2} \rceil + 1$ robots in the graph. $\square$

Notice that for $C_n \odot K_1$, the lower bound of Theorem 4.1 is three if $n \geq 3$ and $n \notin \{4, 6\}$, and it is two if $n \in \{4, 6\}$, whereas the upper bound is n . Since $C_n \odot K_1$ is a unicyclic graph, we may recall that mobile general position sets of unicyclic graphs were discussed in [17].

4.2. Joins of graphs

We now give bounds for the mobile general position number of joins $G \vee H$. Observe that if both G and H are cliques, then $G \vee H$ is also a clique and the question is trivial, so we will assume that at least one of G and H is not a clique.

Theorem 4.4. *If G and H are (not necessarily connected) graphs with clique number at least two, and G and H are not both cliques, then*

$$\min\{\omega(G), \omega(H)\} + 1 \leq \text{Mob}_{\text{gp}}(G \vee H) \leq \omega(G) + \omega(H) - 1.$$

For any graph G with order $n \geq 2$,

$$2 \leq \text{Mob}_{\text{gp}}(G \vee K_1) \leq \omega(G) + 1.$$

Proof. Assume that both G and H have clique number at least two and that at least two robots are traversing $G \vee H$ in general position. At some point there must be a robot in G and a robot in H . Hence, at this point, the set of occupied vertices in G and the occupied vertices in H must both be cliques in G and H , respectively, giving the upper bound $\text{Mob}_{\text{gp}}(G \vee H) \leq \omega(G) + \omega(H)$. However, if $\text{Mob}_{\text{gp}}(G \vee H) = \omega(G) + \omega(H)$, then no robot has a legal move, since any move would result in a clique in one of G and H and a non-clique in the other, so in fact $\text{Mob}_{\text{gp}}(G \vee H) \leq \omega(G) + \omega(H) - 1$.

For the lower bound, assume that $\omega(H) \leq \omega(G)$. We can start with robots at a maximum clique W_H of H and one robot in G . The robot in G can visit every vertex of G , since during this process the occupied vertices always form a clique in $G \vee H$. At the end, this robot moves into a maximum clique W_G of G . After that, all the robots from W_H but one move into W_G . At that time, only one robot remains in H and, by the same argument, it can visit every vertex of H .

The inequalities for $G \vee K_1$ can be derived in a similar manner. $\square$

Note that if G and H both have clique number two, then the upper and lower bounds of Theorem 4.4 coincide. For a triangle-free graph G , $\text{Mob}_{\text{gp}}(G \vee K_1)$ could be either two or three. It is easily seen that for cycles we have $\text{Mob}_{\text{gp}}(C_4 \vee K_1) = 2$ and $\text{Mob}_{\text{gp}}(C_n \vee K_1) = 3$ for $n \geq 5$. The first example shows that the lower bound for $\text{Mob}_{\text{gp}}(G \vee K_1)$ is tight, but it is an open question whether this can happen for graphs with large clique number.

Corollary 4.5. *If G and H both have clique number two, then $\text{Mob}_{\text{gp}}(G \vee H) = 3$.*

To show that the upper bound in Theorem 4.4 is tight, consider the join $K_r^- \vee K_s^-$, where K_n^- represents a complete graph minus one edge. If x_1, x_2 is the pair of non-adjacent vertices in K_r^- and y_1, y_2 is the pair of non-adjacent vertices of K_s^- , then $(V(K_r^-) \setminus \{x_2\}) \cup (V(K_s^-) \setminus \{y_1, y_2\})$ is a mobile general position set, as the set of occupied vertices forms a clique in $K_r^- \vee K_s^-$ and the robot at x_1 can follow the route $x_1 \rightsquigarrow y_1 \rightsquigarrow x_2 \rightsquigarrow y_2$ to visit the remaining vertices. This matches the upper bound. More generally, the same argument works when G and H are both joins of cliques with empty graphs.

To demonstrate sharpness of the lower bound, for $r \geq 2$ take the join $K_r \vee K_{r+1}^+$, where K_{r+1}^+ is the complete graph K_{r+1} with an added leaf x . Suppose that a set of at least $r+2$ robots can traverse this graph in general position, and focus on the moment that there is a robot at x . As $r+2$ robots cannot be stationed on K_{r+1}^+ , there must be a robot on K_r and the positions occupied on K_r and K_{r+1}^+ must both induce cliques. Hence every vertex of K_r must contain a robot and in K_{r+1}^+ there is a robot at x and its support vertex x' . However, there are no legal moves in this configuration. Thus, $\text{Mob}_{\text{gp}}(K_r \vee K_{r+1}^+) \leq r+1 = \min\{\omega(K_r), \omega(K_{r+1}^+)\} + 1$.

For an example of graphs with arbitrarily large clique number that meet the upper bound in $\text{Mob}_{\text{gp}}(G \vee K_1) \leq \omega(G) + 1$, consider the *birdcage* graph B_n formed as follows. Let U be a clique on vertices $\{u_1, \dots, u_n\}$, V be an empty graph on vertices $\{v_1, \dots, v_n\}$ and an additional vertex z , and add edges $u_i \sim v_i$ and $v_i \sim z$ for $i \in [n]$. The graph B_n has clique number n and we show that $n+1$ robots can traverse $B_n \vee K_1$ in general position. Denote the vertex of the K_1 by x . Start with robots at $U \cup \{x\}$ and make the move $x \rightsquigarrow z$. Then for each $i \in [n]$ make the two moves $u_i \rightsquigarrow v_i \rightsquigarrow u_i$. It is easily seen that the robots are always in general position and visit all the vertices of $B_n \vee K_1$.

5. Concluding remarks and open problems

We conclude with a few open problems.

- Is there a non-trivial upper bound on $\text{Mob}_{\text{gp}}(G \square H)$, at least for the particular case $\text{Mob}_{\text{gp}}(G \square K_2)$?
- We have seen that $\text{Mob}_{\text{gp}}(P_\infty \square P_\infty) = 4 = \text{gp}(P_\infty \square P_\infty)$. In [20, Theorem 1] it was proven that $\text{gp}(P_\infty^{k,\square}) = 2^{2^{k-1}}$, where $P_\infty^{k,\square}$ is the k -tuple Cartesian product of the infinite path P_∞ . It is therefore of interest to determine whether $\text{Mob}_{\text{gp}}(P_\infty^{k,\square}) = 2^{2^{k-1}}$ holds for larger values of k .
- Are there graphs G with arbitrarily large clique number such that $\text{Mob}_{\text{gp}}(G \vee K_1) = 2$?
- In view of Proposition 3.7 and the preceding remarks, we ask what is the mobile general position number of cylinder graphs $C_7 \square P_s$ and $C_{10} \square P_s$ for $s \geq 5$?
- By Corollary 3.8, $\text{Mob}_{\text{gp}}(C_r \square C_s) \geq 5$ if $r = 9$ or $r > 10$ and $s \geq 10$. It would be interesting to classify the mobile general position numbers of all torus graphs.
- What is the mobile general position number of strong and direct products?
- What is the mobile general position number of the hypercube?

Declaration of competing interest

Sandi Klavžar is an Associate Editor of the Discrete Applied Mathematics journal and was not involved in the review and decision-making process of this article. The authors do not have any additional conflicts of interest to declare.

Acknowledgements

Sandi Klavžar was supported by the Slovenian Research and Innovation Agency (ARIS) under the grants P1-0297, N1-0285, N1-0355, N1-0431. D. Kuziak and I. G. Yero have been partially supported by the Spanish Ministry of Science and Innovation through the grant PID2023-146643NB-I00. D. Kuziak acknowledges the support from “Ministerio de Educación, Cultura y Deporte”, Spain, under the “José Castillejo” program for young researchers (reference number: CAS22/00081) to make a temporary visit to the University of Ljubljana, where this investigation has been partially developed. The research of James Tuite was partly carried out during a visit to University of Ljubljana supported by a grant from the Crowther Fund. Ethan Shallcross conducted this research with funding from an Open University research bursary. The authors thank the anonymous referees for their suggestions.

Data availability

No data was used for the research described in the article.

References

- [1] 2024, Website <https://www.starship.xyz/>. (Accessed on 09 July 2024).
- [2] A. Aljohani, P. Poudel, G. Sharma, Complete visitability for autonomous robots on graphs, in: IEEE International Parallel and Distributed Processing Symposium, IPDPS, IEEE Computer Society, 2018, pp. 733–742.
- [3] A. Aljohani, G. Sharma, Complete visibility for mobile robots with lights tolerating faults, *Int. J. Netw. Comput.* 8 (2018) 32–52.
- [4] S. Boshar, A. Evans, A. Krishnakumar, E. Shallcross, J. Tuite, Solution to some conjectures on mobile position problems, 2024, [arXiv:2507.16622](https://arxiv.org/abs/2507.16622).
- [5] B. Brešar, I.G. Yero, Lower (total) mutual-visibility number in graphs, *Appl. Math. Comput.* 465 (2024) 128411.
- [6] U. Chandran S.V., G. Di Stefano, H. Sreelatha, E.J. Thomas, J. Tuite, Colouring a graph with position sets, 2024, [arXiv:2408.13494](https://arxiv.org/abs/2408.13494).
- [7] U. Chandran S.V., S. Klavžar, J. Tuite, The general position problem: a survey, 2025, [arXiv:2501.19385](https://arxiv.org/abs/2501.19385).
- [8] U. Chandran S.V., P.K. Neethu, S. Klavžar, J. Tuite, On monophonic position sets of cartesian and lexicographic products, 2025, <https://arxiv.org/abs/2412.09837v2>.
- [9] U. Chandran S.V., G.J. Parthasarathy, The geodesic irredundant sets in graphs, *Int. J. Math. Comb.* 4 (2016) 135–143.
- [10] S. Cicerone, G. Di Stefano, S. Klavžar, On the mutual visibility in Cartesian products and triangle-free graphs, *Appl. Math. Comput.* 438 (2023) 127619.
- [11] M. Dettlaff, M. Lemańska, J.A. Rodríguez-Velázquez, I.G. Yero, Mobile mutual-visibility sets in graphs, *Ars Math. Contemp.* (2025) [http://dx.doi.org/10.26493/1855-3974.3410.9bc](https://dx.doi.org/10.26493/1855-3974.3410.9bc).
- [12] G.A. Di Luna, P. Flocchini, S.G. Chaudhuri, F. Poloni, N. Santoro, G. Viglietta, Mutual visibility by luminous robots without collisions, *Inform. and Comput.* 254 (2017) 392–418.
- [13] G. Di Stefano, Mutual visibility in graphs, *Appl. Math. Comput.* 419 (2022) 126850.
- [14] H.E. Dudeney, Amusements in Mathematics, Nelson, Edinburgh, 1917.
- [15] M. Ghorbani, S. Klavžar, H.R. Maimani, M. Momeni, F. Rahimi-Mahid, G. Rus, The general position problem on Kneser graphs and on some graph operations, *Discuss. Math. Graph Theory* 41 (2021) 1199–1213.
- [16] V. Iršič, S. Klavžar, G. Rus, J. Tuite, General position polynomials, *Results Math.* 79 (3) (2024) 110.
- [17] S. Klavžar, A. Krishnakumar, J. Tuite, I.G. Yero, Traversing a graph in general position, *Bull. Aust. Math. Soc.* 108 (2023) 353–365.
- [18] S. Klavžar, D. Kuziak, J.C. Valenzuela Tripodoro, I.G. Yero, Coloring the vertices of a graph with mutual-visibility property, *Open Math.* 23 (2025) 20250193.
- [19] S. Klavžar, B. Patkós, G. Rus, I.G. Yero, On general position sets in Cartesian products, *Results Math.* 76 (2021) 123.
- [20] S. Klavžar, G. Rus, The general position number of integer lattices, *Appl. Math. Comput.* 390 (2021) 125664.

- [21] D. Korže, A. Vesel, General position sets in two families of Cartesian product graphs, *Mediterr. J. Math.* 20 (2023) 203.
- [22] E. Krufft Welton, S. Khudairi, J. Tuite, Lower general position in Cartesian products, *Commun. Comb. Optim.* 10 (2025) 110–125.
- [23] D. Kuziak, M.L. Puertas, J.A. Rodríguez-Velázquez, I.G. Yero, Strong resolving graphs: the realization and the characterization problems, *Discrete Appl. Math.* 236 (2018) 270–287.
- [24] D. Kuziak, J.A. Rodríguez-Velázquez, Total mutual-visibility in graphs with emphasis on lexicographic and Cartesian products, *Bull. Malays. Math. Sci. Soc.* 46 (2023) 197.
- [25] P. Manuel, S. Klavžar, Graph theory general position problem, 2017, arXiv:1708.09130.
- [26] P. Manuel, S. Klavžar, A general position problem in graph theory, *Bull. Aust. Math. Soc.* 98 (2018) 177–187.
- [27] P. Manuel, S. Klavžar, The graph theory general position problem on some interconnection networks, *Fund. Inform.* 163 (2018) 339–350.
- [28] P. Manuel, R. Prabha, S. Klavžar, The edge general position problem, *Bull. Malays. Math. Sci. Soc.* 45 (2022) 2997–3009.
- [29] O.R. Oellermann, J. Peters-Fransen, The strong metric dimension of graphs and digraphs, *Discrete Appl. Math.* 155 (2007) 356–364.
- [30] J. Tian, S. Klavžar, Variety of general position problems in graphs, *Bull. Malays. Math. Sci. Soc.* 48 (2025) 5.
- [31] J. Tian, K. Xu, The general position number of Cartesian products involving a factor with small diameter, *Appl. Math. Comp.* 403 (2021) 126206.
- [32] J. Tian, K. Xu, S. Klavžar, The general position number of Cartesian product of two trees, *Bull. Aust. Math. Soc.* 104 (2021) 1–10.