

The vertex visibility number of graphs

Dhanya Roy^a, Gabriele Di Stefano^b, Sandi Klavžar^{c,d,e,*}, Aparna S. Lakshmanan^a

^a Department of Mathematics, Cochin University of Science and Technology, Cochin - 22, Kerala, India

^b Department of Information Engineering, Computer Science and Mathematics, University of L'Aquila, Italy

^c Faculty of Mathematics and Physics, University of Ljubljana, Slovenia

^d Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia

^e Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia

ARTICLE INFO

Section Editor: Vladimiro Sassone
Handling Editor: katie harris

AMS subject classifications (2020):

05C12

68Q17

05C76

Keywords:

Vertex visibility

Mutual-visibility

Shortest-path tree

Computational complexity

Cartesian product

ABSTRACT

If $x \in V(G)$, then $S \subseteq V(G) \setminus \{x\}$ is an x -visibility set if for each $y \in S$ there exists a shortest x, y -path avoiding S . The x -visibility number $v_x(G)$ is the maximum cardinality of an x -visibility set, and the maximum value of $v_x(G)$ between all vertices x of G is the vertex visibility number $vv(G)$ of G . It is proved that $vv(G)$ is equal to the largest possible number of leaves of a shortest-path tree of G . Deciding whether $v_x(G) \geq k$ holds for given G , a vertex $x \in V(G)$, and a positive integer k is NP-complete even for graphs of diameter 2. Several general sharp lower and upper bounds on the vertex visibility number are proved. The vertex visibility number of Cartesian products is also bounded from below and above, and the exact value of the vertex visibility number is determined for square grids, square prisms, and square toruses.

1. Introduction

General position and mutual-visibility are very active areas of metric and algorithmic graph theory. The concepts are complementary to each other, and progress in one of the areas typically has an impact on the other area. The general position problem has been recently surveyed in [1], see also [2–5]. For the mutual-visibility problem we refer to the seminal paper [6] and the following selection of studies [4,7–15].

In [16], the general position version was investigated from a vertex's point of view. Among the many motivations for this research, let us highlight the following. In [17][Chapter III] it is shown how to place a set of points with integer positive coordinates (i, j) , $j \leq i$, so that each point is in mutual-visibility with $(0, 0)$ and such that the number of points with the same abscissa is maximized. The problem turns out to be very interesting, as the solution is linked with the Farey series and Euler's totient function ϕ : the number of points with abscissa n is exactly $\phi(n)$.

Due to the motivation above (and more), the paper [16] gives the following definitions. If x is a vertex of a graph G , then $S \subseteq V(G)$ is said to be an x -position set if for any $y \in S$, no vertex of $S \setminus \{y\}$ lies on any shortest x, y -path. The *vertex position number* $vp(G)$ of G is the maximum cardinality of an x -position set among all vertices x of G . The paper [16] yields numerous results dealing with the

* Corresponding author.

E-mail addresses: dhanyaroyku@gmail.com, dhanyaroyku@cusat.ac.in (D. Roy), gabriele.distefano@univaq.it (G. Di Stefano), sandi.klavzar@fmf.uni-lj.si (S. Klavžar), aparnaren@gmail.com, aparnals@cusat.ac.in (A. S. Lakshmanan).

<https://doi.org/10.1016/j.tcs.2026.116193>

Received 21 October 2025; Received in revised form 1 July 2026; Accepted 3 August 2026

Available online 6 August 2026

0304-3975/© 2026 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

largest and smallest orders of maximum x -position sets, in particular giving bounds in terms of the girth, vertex degrees, diameter and radius.

In this paper, we complement the investigation from [16] by considering the mutual-visibility problem from a vertex's point of view. If x is a vertex of a graph G , then $S \subseteq V(G) \setminus \{x\}$ is an x -visibility set if for any $y \in S$ there exists a shortest x, y -path P , such that $V(P) \cap S = \{y\}$. The x -visibility number $v_x(G)$ is the maximum cardinality of an x -visibility set, we also say that $v_x(G)$ is the visibility number of x , cf. Fig. 1. An x -visibility set of cardinality $v_x(G)$ is called a v_x -set. The maximum value of $v_x(G)$ among all vertices x of G is called the vertex visibility number $\text{vv}(G)$ of G , and a corresponding v_x -set is a vv -set, see Fig. 1 again. For instance, if $n \geq 3$, then $v_x(C_n) = 2$ for any $x \in V(C_n)$, hence $\text{vv}(C_n) = 2$. Similarly, if $n \geq 3$, then $v_x(P_n) = \deg_{P_n}(x)$, hence $\text{vv}(P_n) = 2$. It is worth mentioning the following fact, where $n(G)$ denotes the order of a graph G .

Lemma 1.1. *If x is a leaf of a connected graph G with $n(G) \geq 3$, then $v_x(G) < \text{vv}(G)$.*

Proof. If G is a path graph on at least three vertices, then $v_x(G) = 1$ and $\text{vv}(G) = 2$. Assume next that G is not a path, let y be the support vertex of x , and S be a v_x -set of G . Then $S' = S \cup \{x\}$ is a y -visibility set, hence $\text{vv}(G) \geq |S'| > |S| = v_x(G)$. $\square$

The paper is organized as follows. In the following subsection, further definitions needed are listed. In Section 2 we first prove the fundamental reason for the computational difficulty of the vertex visibility number: $\text{vv}(G)$ is equal to the largest possible number of leaves of a shortest-path tree of G . This builds a bridge between the new introduced concepts and the Dijkstra's algorithm, the result of which is precisely a tree of shortest-paths. In the second main result of the section we prove that the x -VISIBILITY problem is NP-complete even for graphs of diameter 2, where the x -VISIBILITY problem asks whether $v_x(G) \geq k$ holds for given G , a vertex $x \in V(G)$, and a positive integer k . In Section 3 we deduce some general sharp lower and upper bounds on the vertex visibility number in terms of the order, the maximum degree, and the eccentricity. The vertex visibility number of Cartesian products is investigated in Section 4. After establishing a general sharp lower bound and an upper bound, the focus is on square grids, square prisms, and square toruses. For each of these Cartesian products, the exact value of the vertex visibility number is obtained. We conclude the paper with some problems for future investigation.

1.1. Concepts and notation

Here we provide further necessary definitions.

Let $G = (V(G), E(G))$ be a graph. The maximum degree of G is denoted by $\Delta(G)$, and the (open) neighborhood of $x \in V(G)$ by $N_G(x)$. A vertex x of G is *simplicial* if the subgraph induced by $N_G(x)$ is complete. A vertex of G is *universal* if it is adjacent to all other vertices of G . A *double star* is a tree in which exactly two vertices are not leaves.

The distance function $d_G(\cdot, \cdot)$ is the standard shortest-path distance. The *eccentricity* $\text{ecc}_G(x)$ of a vertex x is the maximum distance between x and the other vertices of G . A vertex z is an *eccentric vertex* of x if $d_G(x, z) = \text{ecc}_G(x)$. The open interval $I(x, y)$ between vertices x and y is the set of all vertices that lie on shortest x, y -paths other than x and y . When every two vertices x and y of G are connected by a unique shortest x, y -path, G is called *geodetic*. A vertex y is *maximally distant* from x if $d_G(x, y) \geq d_G(x, z)$, for every $z \in N_G(y)$. (This notion was introduced in [18] as a tool to study the strong metric dimension.) The collection of all maximally distant vertices from x is denoted by $\text{MD}_G(x)$.

If $S \subseteq V(G)$, then we say that $x, y \in V(G)$ are S -visible, if there exists a shortest x, y -path P such that $V(P) \cap S \subseteq \{x, y\}$. The set S is a *mutual-visibility set* if any two vertices from S are S -visible. The cardinality of a largest mutual-visibility set of G is the *mutual-visibility number* $\mu(G)$ of G . A mutual-visibility set of cardinality $\mu(G)$ is a μ -set. We also say that a vertex y is S -visible from x if there exists a shortest y, x -path P such that $V(P) \cap S \subseteq \{x, y\}$.

The *Cartesian product* $G \square H$ of graphs G and H has the vertex set $V(G) \times V(H)$, vertices (g, h) and (g', h') of $G \square H$ are adjacent if either $gg' \in E(G)$ and $h = h'$, or $g = g'$ and $hh' \in E(H)$. A G -layer is a subgraph of $G \square H$ induced by $V(G) \times \{h\}$ for some $h \in V(H)$, denoted by G^h . Analogously, for $g \in V(G)$ we have the H -layer ${}^g H$.

Finally, for a positive integer k , the set $\{1, \dots, k\}$ is denoted by $[k]$.

2. Shortest-path trees and computational complexity

To study the computational complexity of finding the x -visibility number $v_x(G)$ for a fixed vertex x of a graph G , we introduce the following decision problem.

Definition 2.1. x -VISIBILITY problem: INSTANCE: A graph G , a vertex $x \in V(G)$, a positive integer $k \leq n(G)$. QUESTION: $v_x(G) \geq k$?

In this section, we prove that the x -VISIBILITY problem is NP-complete even for graphs of diameter 2, so finding $\text{vv}(G)$ is an NP-hard problem. This result is in sharp contrast with the computational complexity of the vertex position problem, which is solvable in $O(n^4 \log(n))$ time, as shown in [16].

The intrinsic difficulty of the x -VISIBILITY problem lies in the fact that the value $\text{vv}(G)$ is equal to the largest possible number of leaves in a shortest-path tree of G . These trees are defined as follows. Let G be a graph, and $x \in V(G)$. Then a tree rooted in x constructed by the BFS search is called a *shortest-path tree* of G . That is, it is a rooted spanning tree T , such that $d_T(x, y) = d_G(x, y)$ for every $y \in V(G)$.

Theorem 2.2. *If G is a connected graph, then $\text{vv}(G)$ is equal to the largest possible number of leaves of a shortest-path tree of G .*




Fig. 1. A graph G where $\ell(G) = 11$ but $\text{vv}(G) = v_x(G) = 10$ (also attained at y and z).

Proof. Let $S = \{x_1, \dots, x_k\}$ be a vv -set of G , and let x be a vertex with $v_x(G) = \text{vv}(G)$. Then, if $n(G) \geq 3$, x is not a leaf because $v_{x'}(G) > v_x(G)$, where x' is the (support) vertex adjacent to x .

Denoting the largest possible number of leaves in a shortest-path tree of G by t_G , it is clear that $\text{vv}(G) \geq t_G$.

To prove $t_G \geq \text{vv}(G)$, we first claim that there exist shortest x, x_i -paths P_i , $i \in [k]$, such that $V(P_i) \cap S = \{x_i\}$ and the edges of $\cup_{i \in [k]} E(P_i)$ induce a tree T in G . Since S is a v_x -set, there exist shortest x, x_i -paths P_i , $i \in [k]$, such that $V(P_i) \cap S = \{x_i\}$. Assume that the paths P_i are selected such that the union $\cup_{i \in [k]} E(P_i)$ induces the smallest number of cycles possible. Suppose that this number is not zero. Then there exist $s, t \in [k]$ such that the union of edges $E(P_s) \cup E(P_t)$ induces a cycle. Let y be a vertex from $V(P_s) \cap V(P_t)$ such that $d_G(x, y)$ is as large as possible among all vertices of $V(P_s) \cap V(P_t)$. Let P'_s and P'_t be the subpaths of P_s and P_t between x and y , respectively. Since P_s and P_t are the shortest paths, P'_s and P'_t are also the shortest paths. Then in our collection $\{P_i : i \in [k]\}$ of shortest paths, replace P_t with the concatenation of P'_s and the y, x_t -subpath of P_t . In this way, the union of the edges from the new collections of shortest x, x_i -paths induces at least one cycle less. This contradicts the selection of the paths P_i . We can conclude that the union $\cup_{i \in [k]} E(P_i)$ does not induce a cycle, thus proving the claim.

Let T be the tree induced by the edges of $\cup_{i \in [k]} E(P_i)$. If T is a spanning tree, then we are done. Hence, assume that T does not span G . Let y be an arbitrary vertex from $V(G) \setminus V(T)$. Then every shortest x, y -path contains at least one vertex from S , for otherwise $S \cup \{y\}$ would be an x -position set larger than S , which is not possible. Moreover, there exists a shortest x, y -path P such that $V(P) \cap S = \{x_j\}$ for some $j \in [k]$, for otherwise S would not be an x -position set. Then replace x_j by y in S and in the collection of our shortest paths, replace the current shortest x, x_j -path by P . Proceeding in this way, we end with an x -position set of the same cardinality as S , and with a collection of paths whose edges induce a spanning tree T of G . We can conclude that $t_G \geq \text{vv}(G)$. $\square$

In view of Theorem 2.2, we first recall that the problem of finding a spanning tree with maximum number of leaves has been extensively researched and is computationally difficult. For instance, the problem is NP-hard and APX-hard for cubic graphs, see [19]. On the other hand, a 2-approximation algorithm is known for this problem [20]. We now demonstrate that the difference between the maximum number of leaves in a spanning tree of G and $\text{vv}(G)$ can be arbitrarily large.

Let $\ell(G)$ denote the maximum number of leaves in a spanning tree of a graph G . Let G_n denote the graph obtained by taking n copies of the graph G in Fig. 1 and identifying the vertex x . Then $\ell(G_n) = 11n$. By Lemma 3.2, it is easy to check that $v_x(G_n) = v_y(G_n) = v_z(G_n) = 10n$ while $v_a(G_n) = 9$, $v_b(G_n) = 8$ and $v_c(G_n) = 8$. Thus, by the symmetry of the graph and by Lemma 1.1, we can conclude that $\text{vv}(G_n) = 10n$. Hence, the difference $\ell(G_n) - \text{vv}(G_n)$ is n , which we summarize in the following result.

Proposition 2.3. *Given a graph G , the difference between $\ell(G)$ and $\text{vv}(G)$ can be arbitrarily large.*

We are now ready to prove that the x -VISIBILITY problem is hard to solve.

Theorem 2.4. *The x -VISIBILITY problem is NP-complete even for graphs of diameter 2.*

Proof. Given a set $X \subseteq V(G) \setminus \{x\}$ of G , it is possible to test in polynomial time whether it is a x -visibility set or not. Consequently, the problem is in NP.


We will now prove that the INDEPENDENT SET problem (equivalent to the CLIQUE problem in the complement graph and shown as NP-complete in [21]), polynomially reduces to the x -VISIBILITY problem. The INDEPENDENT SET problem asks if the independence number $\alpha(G)$ of a graph G , that is, the cardinality of a largest mutually non adjacent set of vertices of G , is at least a given integer t . Here we assume that G does not contain any isolated vertices. Clearly, this assumption does not affect the hardness of the problem. We consider an arbitrary instance (G, t) of the INDEPENDENT SET problem where G has no isolated vertices and construct an instance (G', k) of the x -VISIBILITY problem as follows.

Let $V(G) = [n]$ and add a universal vertex x to $V(G)$. Now for each edge $e = ij$ of G , we add a new vertex $v_e = v_{ij}$ and the edges iv_e and jv_e . Also, let $S = \{v_e : e \in E(G)\}$ induce a clique. An example of the graph G' , for $G = P_5$, is given in Fig. 2.

Let (G, t) be an instance of the INDEPENDENT SET problem, we show that $\alpha(G) \geq t$ if and only if the x -VISIBILITY problem has x -visibility number greater than k , where $k = E(G) + t$, that is, $v_x(G') \geq k$.

Let I be an independent set of G , and consider the set $X = I \cup S$ in G' . For each $v_e \in S$, where $e = ij$, either i or j is not present in I and hence all vertices in S are x -visible. In addition, all vertices in I are adjacent to x and hence are x -visible. Therefore, $v_x(G') \geq k$.

On the contrary, let X be an x -visibility set of G' of cardinality at least k . By Lemma 3.2, we can assume that all the vertices of S are in X . Then the vertices in $X \setminus S$ must form an independent set of G , as, otherwise, there would be an edge $e = ij$ such that both

Fig. 2. The graph G' for $G = P_5$.

vertices i and j are in X , but then v_e would not be in visibility with x . Since the number of vertices in S is $|E(G)|$, the number of vertices in the independent set of G is $|X \setminus S| \geq k - |E(G)| = t$. Hence, $\alpha(G) \geq t$.

Finally, the diameter of G' is two as x is adjacent to all vertices of G and at distance 2 from the vertices in S . All pairs of vertices u, v in G are at distance at most two as u, x, v is a path in G' . Each vertex in S is adjacent to all other vertices in S and, since G has no isolated vertices, it is at a distance of at most two from the vertices in G . $\square$

The above Theorem 2.4 clearly implies that finding $\text{vv}(G)$ is also an NP-hard problem.

3. General lower and upper bounds

In view of the hardness of the problem studied, as established in Section 2, in this section we prove general lower and upper bounds for the vertex visibility numbers. Along the way, several exact values are also determined.

If S is a μ -set of G and $x \in S$, then $S \setminus \{x\}$ is an x -visibility set of G . Also, $N_G(x)$ is an x -visibility set for every vertex x . Thus we have the following general bounds:

$$\max\{\mu(G) - 1, \Delta(G)\} \leq \text{vv}(G) \leq n(G) - 1. \quad (1)$$

The difference $\text{vv}(G) - (\mu(G) - 1)$ can be arbitrarily large. For instance, it can be deduced from [22][Theorem 3.2.(i)] that if $n \geq 6$, then $\mu(K_2 \square C_n) = 6$, while on the other hand $\text{vv}(K_2 \square C_n) \geq n$. The first assertion of the following result follows from (1), while the second is also straightforward to verify, hence we omit the proof.

Proposition 3.1. *If G is a connected graph with $n(G) \geq 2$, then the following properties hold.*

1. $\text{vv}(G) = n(G) - 1$ if and only if G has a universal vertex.
2. $\text{vv}(G) = n(G) - 2$ if and only if G contains no universal vertex and contains a spanning double star.

We now say that a vertex y is a *stress vertex* for x if there exists a maximally distant vertex z of x such that y lies on every x, z -shortest path in G . Note, that every cut vertex y , $y \neq x$, is a stress vertex for x . Let $\text{str}_G(x)$ denote the number of stress vertices for x .

Lemma 3.2. *If x is a vertex of a connected graph G , then there exists a v_x -set S with the following properties:*

1. every $y \in V(G) \setminus \{x\}$ is S -visible from x ;
2. $\text{MD}_G(x) \subseteq S$;
3. S contains no stress vertex for x .

In addition, $|\text{MD}_G(x)| \leq v_x(G) \leq n(G) - \text{str}_G(x) - 1$.

Proof. Let R be a v_x -set of G such that it contains as many vertices of $\text{MD}_G(x)$ as possible. Suppose that there exists a vertex $y \in \text{MD}_G(x) \setminus (R \cup \{x\})$. Then every shortest x, y -path intersects R . Also, there exists a shortest x, y -path that contains exactly one vertex from R , say z , for otherwise R would not be an x -visibility set. But then $(R \setminus \{z\}) \cup \{y\}$ is an x -visibility set of the same cardinality as R that contains more vertices from $\text{MD}_G(x)$, a contradiction.

By the above we have $\text{MD}_G(x) \subseteq R$. It follows from here by the definition of the stress vertex that R contains no stress vertex for x and that every $y \in V(G) \setminus \{x\}$ is R -visible from x . The set R is thus a v_x -set with the three required properties. Finally, the second property yields the claimed lower bound, while the third property gives the upper bound. $\square$

Note that Lemma 3.2 implies that if G is a geodetic graph, then $v_x(G) = |\text{MD}_G(x)|$ for every $x \in V(G)$. For a graph G , where $|\text{MD}_G(x)| < v_x(G) < n(G) - \text{str}_G(x) - 1$, see Fig. 3.

For block graphs, which form a special case of geodetic graphs, a more specific result can be stated. In fact, this is an example where the lower and upper bounds in Lemma 3.2 coincide. Denoting by $s(G)$ the number of simplicial vertices of a graph G , the result for block graphs reads as follows.




Fig. 3. A graph G , where $v_x(G) = 4$ while $|\text{MD}_G(x)| = \text{str}_G(x) = 2$.

Proposition 3.3. *If G is a block graph different from a complete graph, then*

$$\text{vv}(G) = s(G).$$

Proof. Since G is not complete, it contains a cut vertex x . Then $\text{MD}_G(x)$ consists of the simplicial vertices of G . By the lower bound of Lemma 3.2 we thus get $\text{vv}(G) \geq s(G)$. On the other hand, as already mentioned, cut vertices are stress vertices, therefore by the upper bound of Lemma 3.2 we also get $\text{vv}(G) \leq s(G)$. $\square$

We continue with the next upper bound depending on the order and the maximum degree.

Theorem 3.4. *If G has no universal vertex, then*

$$\text{vv}(G) \leq \left\lfloor \frac{n(G)\Delta(G) - 1}{\Delta(G) + 1} \right\rfloor,$$

and the bound is sharp.

Proof. Let S be a vv-set of G and set $n = n(G)$ and $\Delta = \Delta(G)$ for the rest of the proof.

Assume first that $\text{vv}(G) = \Delta$. Then we have

$$\text{vv}(G) = \Delta \leq \frac{n\Delta - 1}{\Delta + 1},$$

where the above inequality holds since $\Delta \leq n - 2$ by the assumption of the theorem.

Next, assume that $\text{vv}(G) > \Delta$. Let x be a vertex such that $v_x(G) = \text{vv}(G)$ and let S be a v_x -set. Then $|S| = \text{vv}(G)$. Let $S' = S \setminus N_G(x)$ and let $s' = |S'|$. Since $\text{vv}(G) > \Delta$ we have $s' \geq 1$. This in turn implies that at least one neighbor of x does not belong to S . Consequently,

$$s' \geq \text{vv}(G) - (\deg_G(x) - 1) \geq \text{vv}(G) - \Delta + 1. \quad (2)$$

Since S is an x -visibility set, each vertex of S' has a neighbor in $V(G) \setminus (S \cup \{x\})$. It follows that

$$s' \leq (n - \text{vv}(G) - 1)\Delta. \quad (3)$$

Combining (2) with (3) we get

$$\text{vv}(G) - \Delta + 1 \leq (n - \text{vv}(G) - 1)\Delta.$$

Rearranging this inequality and using the fact that $\text{vv}(G)$ is a positive integer, the claimed inequality follows.

To show the sharpness of the bound, consider the cocktail party graph $K_{k \times 2}$, (the complete k -partite graph where each partite set is of cardinality 2) of order $2k$, where $k \geq 2$. Then $\Delta(K_{k \times 2}) = 2k - 2 = \text{vv}(K_{k \times 2}) = 2k - 2$. Since

$$\left\lfloor \frac{2k(2k - 2) - 1}{(2k - 2) + 1} \right\rfloor = \left\lfloor \frac{2k - 2 - \frac{1}{2k}}{1 - \frac{1}{2k}} \right\rfloor = 2k - 2,$$

the equality holds for $K_{k \times 2}$, $k \geq 2$. $\square$

Next, we bound the x -visibility number using the eccentricity of x .

Proposition 3.5. *If G is a graph and $x \in V(G)$, then*

$$\frac{n(G) - 1}{\text{ecc}_G(x)} \leq v_x(G) \leq n(G) - \text{ecc}_G(x),$$

and the bounds are sharp.

Proof. Let S be a v_x -set, and let y be an eccentric vertex of x in G . Let $P : x = x_0, x_1, \dots, x_k = y$ be the shortest x, y -path selected such that $|V(P) \cap S|$ is as small as possible. Suppose that S contains more than one vertex from P , say x_i and x_j , where $i < j$. Since S is an x -visibility set, there exists a shortest x, x_j -path Q such that $V(Q) \cap S = \{x_j\}$. This in turn implies that the path Q followed by the x_j, y -subpath of P is a shortest x, y -path that contains fewer vertices from S than P , a contradiction to the choice of P . Hence $|V(P) \cap S| \leq 1$ which in turn yields $v_x(G) \leq n(G) - \text{ecc}_G(x)$. For the lower bound, let $N_i(x)$, $i \in [\text{ecc}_G(x)]$, be the set of all vertices at distance exactly i from x . Then each of these is an x -visibility set, and one of them must have order at least $(n(G) - 1)/\text{ecc}_G(x)$.

If x is the universal vertex in a star graph, then the lower and upper bounds coincide. $\square$

4. Cartesian products

In this section, we consider the Cartesian product graphs and first prove the general lower and upper bounds on their vertex visibility number. Next, we focus on square grids $(P_n \square P_n)$, square prisms $(P_n \square C_n)$, and square toruses $(C_n \square C_n)$, and determine exact values for the vertex visibility number in all three cases. The general bounds read as follows.

Proposition 4.1. *If G and H are graphs with $n(G) \geq n(H)$, then*

$$\max\{\Delta(G)n(H), \Delta(H)n(G)\} \leq \text{vv}(G \square H) \leq (n(G) - 1)n(H).$$

Both bounds are sharp, in particular, $\text{vv}(K_m \square K_n) = mn - m$ for $m \geq n \geq 2$.

Proof. Let $g \in V(G)$ be such that $\deg_G(g) = \Delta(G)$ and let h be an arbitrary vertex of H . Then $N_G(x) \times V(H)$ is a (g, h) -visibility set of $G \square H$ with cardinality $\Delta(G)n(H)$. By the commutativity of the Cartesian product operation, there also exists a (g', h') -visibility set of $G \square H$ with cardinality $\Delta(H)n(G)$. Thus, $\text{vv}(G \square H) \geq \max\{\Delta(G)n(H), \Delta(H)n(G)\}$.

To prove that $\text{vv}(G \square H) \leq (n(G) - 1)n(H)$ suppose, in contrast, that $G \square H$ contains a (g, h) -visibility set S with $|S| > (n(G) - 1)n(H)$. Then there exist $g' \in V(G)$ and $h' \in V(H)$ such that $V(G^{h'}) \subseteq S$ and $V(g' H) \subseteq S$. But then the vertex $(g', h') \in S$ is not S -visible from (g, h) , a contradiction.

If G has a universal vertex, then the lower and upper bounds coincide. A particular case of this situation is the product $K_n \square K_m$. $\square$

Next, we will cover each of the square grids, square prisms, and square toruses in their own subsections.

4.1. Square grids

In this subsection, we determine the vertex visibility number of square grids.

Theorem 4.2. *If $n \geq 4$, then*

$$\text{vv}(P_n \square P_n) = \frac{n^2 + n - 2}{2}.$$

Proof. Let $V(P_n \square P_n) = \{(u_k, v_l) : k, l \in [n]\}$. Consider an arbitrary vertex $x = (u_{i+1}, v_{j+1}) \in V(P_n \square P_n)$, where we may assume without loss of generality that $0 \leq i \leq j \leq \lfloor \frac{n-1}{2} \rfloor$. Let $X = V^{(u_{i+1})} P_n$ and $Y = V^{(v_{j+1})} P_n$. Set further

$$\begin{aligned} Q_1 &= \{(u_k, v_l) : k < i+1, l < j+1\}, \\ Q_2 &= \{(u_k, v_l) : k < i+1, l > j+1\}, \\ Q_3 &= \{(u_k, v_l) : k > i+1, l > j+1\}, \\ Q_4 &= \{(u_k, v_l) : k > i+1, l < j+1\}. \end{aligned}$$

Let S be a v_x -set in $P_n \square P_n$ that satisfies the properties given in Lemma 3.2 and has the maximum number of vertices from $\bigcup_{t=1}^4 Q_t$ among its choices.

For $(u_k, v_l) \in Q_1$, a shortest $(u_{i+1}, v_{j+1}), (u_k, v_l)$ -path passes through either (u_k, v_{l+1}) or (u_{k+1}, v_l) , and hence these two vertices cannot be simultaneously in S . A similar argument is valid for $(u_k, v_l) \in Q_t, t \in [4]$. Now, partition each Q_t into diagonals D_{pq} , where $p, q \in Q_t$ are its end vertices, as illustrated in Fig. 4. The rectangles indicate the sets $Q_i, i \in [4]$, and the dotted lines indicate the diagonals D_{pq} .

By the above argument, we can have at most $\lceil \frac{|D_{pq}|}{2} \rceil$ vertices in $S \cap D_{pq}$. The diagonals D_{pq} in Q_1 can be explicitly represented as $D_a = \{(u_k, v_l) : k + l = a\}, a \in [i+j] \setminus \{1\}$. Thus if i is even, then

$$|S \cap Q_1| = \sum_{a \in [i+j] \setminus \{1\}} |S \cap D_a| \leq 4 \left(1 + 2 + \dots + \left\lfloor \frac{i-1}{2} \right\rfloor \right) + (j-i+3) \left\lceil \frac{i}{2} \right\rceil,$$

and if i is odd,


$$|S \cap Q_1| = \sum_{a \in [i+j] \setminus \{1\}} |S \cap D_a| \leq 4 \left(1 + 2 + \dots + \left\lfloor \frac{i-1}{2} \right\rfloor \right) + (j-i+1) \left\lceil \frac{i}{2} \right\rceil.$$

On simplification we get,

$$|S \cap Q_1| \leq \begin{cases} \frac{i(j+1)}{2}; & i \text{ even}, \\ \frac{(i+1)j}{2}; & i \text{ odd}. \end{cases}$$

A similar computation can be done for each $Q_t, t \in [4]$. It may be noted that by our choice of i and j we have, $i \leq j \leq n-j-1 \leq n-i-1$. Consequently we obtain the following inequalities:

$$\begin{aligned} |S \cap Q_2| &\leq \begin{cases} \frac{i(n-j)}{2}; & i \text{ even}, \\ \frac{(i+1)(n-j-1)}{2}; & i \text{ odd}, \end{cases} \\ |S \cap Q_3| &\leq \begin{cases} \frac{(n-j-1)(n-i)}{2}; & n-j-1 \text{ even}, \\ \frac{((n-j)(n-i-1))}{2}; & n-j-1 \text{ odd}, \end{cases} \end{aligned}$$

Fig. 4. vv-sets of $P_6 \square P_6$ and $P_7 \square P_7$.

$$|S \cap Q_4| \leq \begin{cases} \frac{j(n-i)}{2}; & j \text{ even,} \\ \frac{(j+1)(n-i-1)}{2}; & j \text{ odd.} \end{cases}$$

By the choice of S , every vertex must be S -visible from (u_{i+1}, v_{j+1}) and hence $S \cap (X \cup Y) \subseteq \{(u_{i+1}, v_1), (u_{i+1}, v_n), (u_1, v_{j+1}), (u_n, v_{j+1})\}$. But, (u_{i+1}, v_1) can be in S only if both (u_i, v_2) and (u_{i+2}, v_2) are not in S since (u_i, v_1) and (u_{i+2}, v_1) must be S -visible from (u_{i+1}, v_{j+1}) . A similar argument holds for (u_{i+1}, v_n) , (u_1, v_{j+1}) , and (u_n, v_{j+1}) . Also, note that in Q_i , at least one end vertex of D_{pq} must be in S if $|D_{pq}|$ is even, and both end vertices of D_{pq} must be in S if $|D_{pq}|$ is odd for $S \cap Q_i$ to attain the maximum cardinality. Since $i \leq j$ we get that

$$\text{in } Q_1 : |D_{p(u_2, v_j)}| = i - 1 \text{ and } |D_{(u_i, v_2)q}| = \begin{cases} i - 1; & i = j, \\ i; & i < j. \end{cases}$$

A similar computation can be done in Q_2 , Q_3 , and Q_4 , to obtain the following respective cardinalities:

$$\begin{aligned} |D_{p(u_2, v_{j+2})}| = i - 1 \text{ and } |D_{(u_i, v_{n-1})q}| &= \begin{cases} i - 1; & i = n - j - 1, \\ i; & i < n - j - 1. \end{cases} \\ |D_{p(u_i, v_{n+1})}| = n - j - 2 \text{ and } |D_{(u_{n-1}, v_{j+2})q}| &= \begin{cases} n - j - 2; & i = j, \\ n - j - 1; & i < j. \end{cases} \\ |D_{p(u_{i+2}, v_2)}| = j - 1 \text{ and } |D_{(u_{n-1}, v_j)q}| &= \begin{cases} j - 1; & j = n - i - 1, \\ j; & j < n - i - 1. \end{cases} \end{aligned}$$

Then, since we have assumed that S has the maximum number of vertices from $\bigcup_{t=1}^4 Q_t$, we infer that $|S \cap (X \cup Y)|$ equals 0, 1, or 2, according to n , i , and j being even or odd. Considering all possible combinations, we get the maximum possible cardinality as $\frac{n^2+n-2}{2}$.

Therefore, $vv(P_n \square P_n) \leq \frac{n^2+n-2}{2}$.

Now, choose $x = (u_2, v_2)$ and let

$$S = \{(u_k, v_1) : k \in [n]\} \cup \{(u_1, v_l) : l \in [n] \setminus \{2\}\} \cup S',$$

where $S' \subset Q_3$ is chosen by taking $\lceil \frac{|D_a|}{2} \rceil$ alternate vertices from each diagonal D_a there; see Fig. 4 which illustrates this construction on $P_6 \square P_6$ and on $P_7 \square P_7$. Then S is an x -visibility set of cardinality $\frac{n^2+n-2}{2}$. We can conclude that $vv(P_n \square P_n) = \frac{n^2+n-2}{2}$. $\square$

4.2. Square prisms

The proof technique used in Theorem 4.2 can be used to determine the vertex visibility number of square prisms.




Fig. 5. vv-sets of $P_5 \square C_5$ and $P_7 \square C_7$.

Theorem 4.3. If $n \geq 4$, then

$$v\mathcal{W}(P_n \square C_n) = \begin{cases} \frac{n^2+3}{2}; & n \equiv 1 \pmod{4}, \\ \frac{n^2+n-2}{2}; & n \equiv 3 \pmod{4}, \\ \frac{2n^2+n}{4}; & n \equiv 0 \pmod{4}, \\ \frac{2n^2+n-2}{4}; & n \equiv 2 \pmod{4}. \end{cases}$$

Proof. Let $V(P_n \square C_n) = \{(u_k, v_l) : k, l \in [n]\}$. By the symmetry, a vertex $x \in V(P_n \square C_n)$ has the same x -visibility number as some vertex from $\{(u_k, v_{\lceil n/2 \rceil}) : k \in [\lceil n/2 \rceil]\}$, hence we may without loss of generality assume that $x = (u_{i+1}, v_{\lceil n/2 \rceil})$, where $0 \leq i \leq \lfloor \frac{n-1}{2} \rfloor$. Let $X = V(P_{i+1} \square C_n)$ and $Y = V(P_n^{\lceil n/2 \rceil})$. Set further

$$\begin{aligned} Q_1 &= \{(u_k, v_l) : k < i+1, l < \lceil n/2 \rceil\}, \\ Q_2 &= \{(u_k, v_l) : k < i+1, l > \lceil n/2 \rceil\}, \\ Q_3 &= \{(u_k, v_l) : k > i+1, l > \lceil n/2 \rceil\}, \\ Q_4 &= \{(u_k, v_l) : k > i+1, l < \lceil n/2 \rceil\}. \end{aligned}$$

Let S be a v_x -set in $P_n \square C_n$ that satisfies the properties given in Lemma 3.2 and has the maximum number of vertices from $\bigcup_{i=1}^4 Q_i$ among its choices.

First assume n is odd and $x = (u_{i+1}, v_{\frac{n+1}{2}})$, where $0 \leq i \leq \frac{n-1}{2}$. For $(u_k, v_l) \in Q_1$, a shortest $x, (u_k, v_l)$ -path passes through either (u_k, v_{l+1}) or (u_{k+1}, v_l) , hence these two vertices cannot be simultaneously in S . A similar argument is valid for each $(u_k, v_l) \in Q_i$, $i \in [4]$. Now, partition each Q_i into diagonals D_{pq} , where $p, q \in Q_i$ are its end vertices as in Theorem 4.2. Then we can have at most $\lceil \frac{|D_{pq}|}{2} \rceil$ vertices in $S \cap D_{pq}$. Hence we get that

$$\text{for } t \in \{1, 2\} : |S \cap Q_t| \leq \begin{cases} \frac{i(n+1)}{4}; & i \text{ even}, \\ \frac{(i+1)(n-1)}{4}; & i \text{ odd}, \end{cases}$$

and

$$\text{for } t \in \{3, 4\} : |S \cap Q_t| \leq \begin{cases} \frac{(n-1)(n-i)}{4}; & n \equiv 1 \pmod{4}, \\ \frac{(n+1)(n-i-1)}{4}; & n \equiv 3 \pmod{4}. \end{cases}$$

Thus we obtain,

$$|S \cup \bigcup_{i=1}^4 Q_i| \leq \begin{cases} \frac{n^2-1}{2}; & n \equiv 1 \pmod{4}, \\ \frac{n^2+n-4}{2}; & n \equiv 3 \pmod{4}. \end{cases}$$

By the choice of S , every vertex must be S -visible from x and hence $S \cap (X \cup Y) \subseteq \{(u_{i+1}, v_1), (u_{i+1}, v_n), (u_1, v_{\frac{n+1}{2}}), (u_n, v_{\frac{n+1}{2}})\}$. But, (u_{i+1}, v_1) can be in S only if both (u_i, v_2) and (u_{i+2}, v_2) are not in S since (u_i, v_1) and (u_{i+2}, v_1) must be S -visible from x . Similar




Fig. 6. vv-sets of $P_6 \square C_6$ and $P_8 \square C_8$.

argument holds for (u_{i+1}, v_n) , $(u_1, v_{\frac{n+1}{2}})$ and $(u_n, v_{\frac{n+1}{2}})$. Also, since $i \leq \frac{n-1}{2} \leq n-i-1$ we have,

$$\begin{aligned} \text{in } Q_1 : \quad |D_{p(u_2, v_{\frac{n-1}{2}})}| &= i-1 \text{ and } |D_{(u_i, v_2)q}| = \begin{cases} i-1; & i = \frac{n-1}{2}, \\ i; & i < \frac{n-1}{2}, \end{cases} \\ \text{in } Q_2 : \quad |D_{p(u_2, v_{\frac{n+3}{2}})}| &= i-1 \text{ and } |D_{(u_i, v_{n-1})q}| = \begin{cases} i-1; & i = \frac{n-1}{2}, \\ i; & i < \frac{n-1}{2}, \end{cases} \\ \text{in } Q_3 : \quad |D_{p(u_i, v_{n+1})}| &= \frac{n-3}{2} \text{ and } |D_{(u_{n-1}, v_{\frac{n+3}{2}})q}| = \begin{cases} \frac{n-3}{2}; & i = \frac{n-1}{2}, \\ \frac{n-1}{2}; & i < \frac{n-1}{2}, \end{cases} \\ \text{in } Q_4 : \quad |D_{p(u_{i+2}, v_2)}| &= \frac{n-3}{2} \text{ and } |D_{(u_{n-1}, v_{\frac{n-1}{2}})q}| = \begin{cases} \frac{n-3}{2}; & i = \frac{n-1}{2}, \\ \frac{n-1}{2}; & i < \frac{n-1}{2}. \end{cases} \end{aligned}$$

Then, since we have assumed that S has the maximum number of vertices from $\bigcup_{t=1}^4 Q_t$ we get,

$$|S \cap (X \cup Y)| = \begin{cases} 2; & n \equiv 1 \pmod{4}, \\ 1; & n \equiv 3 \pmod{4}. \end{cases}$$

Thus,

$$|S| \leq \begin{cases} \frac{n^2+3}{2}; & n \equiv 1 \pmod{4}, \\ \frac{n^2+n-2}{2}; & n \equiv 3 \pmod{4}. \end{cases}$$

Now, assume n is even and $x = (u_{i+1}, v_{\frac{n}{2}})$, where $0 \leq i \leq \frac{n-2}{2}$. For each $(u_k, v_l) \in Q_t$, we know that the set $N((u_k, v_l)) \cap I(x, (u_k, v_l))$ has cardinality at most three and has a vertex which is not in S . Particularly for $(u_k, v_l) \in Q_1$, this vertex is either (u_k, v_{l+1}) or (u_{k+1}, v_l) . Now, partition each Q_t into diagonals D_{pq} , where $p, q \in Q_t$ are its end vertices. Then we can have at most $\sum_{pq} \lceil |D_{pq}|/2 \rceil$ vertices in $S \cap \bigcup_{t=1}^4 Q_t$. Since we have

$$|S \cap Q_1| + |S \cap Q_2| \leq \frac{(i+1)(n-2)}{4} + \frac{(i+1)n}{4},$$

we obtain that for $n \equiv 0 \pmod{4}$,

$$|S \cup \bigcup_{t=1}^4 Q_t| = |S \cap Q_1| + |S \cap Q_2| + |S \cap Q_3| + |S \cap Q_4|$$

$$\begin{aligned} &\leq \frac{(i+1)(n-2)}{4} + \frac{(i+1)n}{4} + \frac{n(n-i)}{4} + \frac{n(n-i-1)}{4} \\ &= \frac{2n^2 + n - 4}{4}, \end{aligned}$$

and for $n \equiv 2 \pmod{4}$,

$$\begin{aligned} |S \cup \bigcup_{i=1}^4 Q_i| &= |S \cap Q_1| + |S \cap Q_2| + |S \cap Q_3| + |S \cap Q_4| \\ &\leq \frac{(i+1)(n-2)}{4} + \frac{(i+1)n}{4} + \frac{(n+2)(n-i-1)}{4} + \frac{(n-2)(n-i)}{4} \\ &= \frac{2n^2 + n - 6}{4}. \end{aligned}$$

Again, by the choice of S , every vertex must be S -visible from x and hence, $S \cap (X \cup Y) \subseteq \{(u_{i+1}, v_1), (u_{i+1}, v_{n-1}), (u_{i+1}, v_n), (u_1, v_{\frac{n}{2}}), (u_n, v_{\frac{n}{2}})\}$. Since $i \leq \frac{n-2}{2} \leq n-i-1$ we have

$$|D_{p(u_2, v_{\frac{n-2}{2}})}| = |D_{p(u_2, v_{\frac{n+2}{2}})}| = i-1$$

and

$$|D_{(u_i, v_2)q}| = |D_{(u_i, v_{n-2})q}| = \begin{cases} i-1; & i = \frac{n-2}{2}, \\ i; & i < \frac{n-2}{2}, \end{cases}$$

$$|D_{(u_i, v_{n-1})q}| = \begin{cases} i-1; & i = \frac{n}{2}, \\ i; & i < \frac{n}{2} \end{cases}$$

$$|D_{p(u_i, v_{n+1})}| = \frac{n-2}{2} \text{ and } |D_{(u_{n-1}, v_{\frac{n+2}{2}})q}| = \begin{cases} \frac{n-2}{2}; & i = \frac{n-2}{2}, \\ \frac{n}{2}; & i < \frac{n-2}{2}, \end{cases}$$

$$|D_{p(u_{i+2}, v_{n-2})}| = |D_{p(u_{i+2}, v_2)}| = \frac{n-4}{2} \text{ and } |D_{(u_{n-1}, v_{\frac{n-2}{2}})q}| = \begin{cases} \frac{n-4}{2}; & i = \frac{n}{2}, \\ \frac{n-2}{2}; & i < \frac{n}{2}. \end{cases}$$

Then, since we have assumed that S has the maximum number of vertices from $\bigcup_{i=1}^4 Q_i$, we get,

$$|S \cap (X \cup Y)| = 1.$$

Thus,

$$|S| \leq \begin{cases} \frac{2n^2+n}{4}; & n \equiv 0 \pmod{4}, \\ \frac{2n^2+n-2}{4}; & n \equiv 2 \pmod{4}. \end{cases}$$

Now, choose $x = (u_2, v_{\lceil \frac{n}{2} \rceil})$ and let


$$R = \{(u_1, v_l) : l \in [n] \setminus \{\lceil \frac{n}{2} \rceil\}\} \cup R' \cup R'',$$

where $R' \subset Q_3$ and $R'' \subset Q_4$ are chosen by taking $\lceil \frac{|D_{pq}|}{2} \rceil$ alternate vertices from each diagonal D_{pq} there; see Figs. 5 and 6 which illustrates this construction on $P_5 \square C_5$, $P_6 \square C_6$, $P_7 \square C_7$ and $P_8 \square P_8$. Then $R \cup \{(u_1, v_{\frac{n+1}{2}}), (u_n, v_{\frac{n+1}{2}})\}$ is an x -visibility set of cardinality $\frac{n^2+3}{2}$ for $n \equiv 1 \pmod{4}$ and $R \cup \{(u_1, v_{\frac{n+1}{2}})\}$ is an x -visibility set of cardinality $\frac{n^2+n-2}{2}$ for $n \equiv 3 \pmod{4}$. Also, $R \cup \{(u_1, v_{\frac{n+1}{2}})\}$ is an x -visibility set of cardinality $\frac{2n^2+n}{4}$ for $n \equiv 0 \pmod{4}$ and of cardinality $\frac{2n^2+n-2}{4}$ for $n \equiv 2 \pmod{4}$. Since we have covered all possible cases, the proof is complete. $\square$

4.3. Square toruses

Theorem 4.4. *If $n \geq 4$, then*

$$vv(C_n \square C_n) = \begin{cases} \frac{n^2-1}{2}; & n \equiv 1 \pmod{4}, \\ \frac{n^2+3}{2}; & n \equiv 3 \pmod{4}, \\ \frac{n^2}{2}; & n \text{ even.} \end{cases}$$

Fig. 7. vv-sets of $C_5 \square C_5$ and $C_7 \square C_7$.

Proof. Let $V(C_n \square C_n) = \{(u_k, v_l) : k, l \in [n]\}$. By the symmetry, all vertices of $C_n \square C_n$ have the same visibility number, and hence we may without loss of generality assume that $x = (u_{\lceil n/2 \rceil}, v_{\lceil n/2 \rceil})$. Let $X = V(C_n^{\lceil n/2 \rceil})$ and $Y = V(C_n^{\lfloor n/2 \rfloor})$. Set further

$$\begin{aligned} Q_1 &= \{(u_k, v_l) : k, l < \lceil n/2 \rceil\}, \\ Q_2 &= \{(u_k, v_l) : k < \lceil n/2 \rceil, l > \lceil n/2 \rceil\}, \\ Q_3 &= \{(u_k, v_l) : k, l > \lceil n/2 \rceil\}, \\ Q_4 &= \{(u_k, v_l) : k > \lceil n/2 \rceil, l < \lceil n/2 \rceil\}. \end{aligned}$$

Let S be a v_x -set in $C_n \square C_n$ that satisfies the properties given in Lemma 3.2 and has the maximum number of vertices from $\bigcup_{t=1}^4 Q_t$ among its choices.

First assume n is odd and $x = (u_{\frac{n+1}{2}}, v_{\frac{n+1}{2}})$. For $(u_k, v_l) \in Q_1$, a shortest $x, (u_k, v_l)$ -path contains either (u_k, v_{l+1}) or (u_{k+1}, v_l) , hence these two vertices cannot simultaneously belong to S . A similar argument is valid for each $(u_k, v_l) \in Q_t$, $t \in [4]$. Now, partition each Q_t into diagonals D_{pq} , where $p, q \in Q_t$ are its end vertices as in Theorem 4.2. Then we can have at most $\lceil \frac{|D_{pq}|}{2} \rceil$ vertices in $S \cap D_{pq}$. Thus we get,

$$|S \cap Q_t| \leq \frac{n^2 - 1}{8}$$

for each $t \in [4]$ and hence,

$$|S \cap \bigcup_{t=1}^4 Q_t| \leq \frac{n^2 - 1}{2}.$$


By the choice of S , every vertex must be S -visible from x and hence $S \cap (X \cup Y) \subseteq \{(u_{\frac{n+1}{2}}, v_1), (u_{\frac{n+1}{2}}, v_n), (u_1, v_{\frac{n+1}{2}}), (u_n, v_{\frac{n+1}{2}})\}$. But, $(u_{\frac{n+1}{2}}, v_1)$ can be in S only if both $(u_{\frac{n-1}{2}}, v_2)$ and $(u_{\frac{n+3}{2}}, v_2)$ are not in S since $(u_{\frac{n-1}{2}}, v_1)$ and $(u_{\frac{n+3}{2}}, v_1)$ must be S -visible from x . Similar argument holds for $(u_{\frac{n+1}{2}}, v_n)$, $(u_1, v_{\frac{n+1}{2}})$, and $(u_n, v_{\frac{n+1}{2}})$. Also, each of the diagonals $D_{(u_{\frac{n-1}{2}}, v_2)(u_2, v_{\frac{n-1}{2}})}$, $D_{(u_{\frac{n-1}{2}}, v_{n-1})(u_2, v_{\frac{n+3}{2}})}$, $D_{(u_{n-1}, v_{\frac{n+3}{2}})(u_{\frac{n+3}{2}}, v_{n-1})}$, and $D_{(u_{n-1}, v_{\frac{n-1}{2}})(u_{\frac{n+3}{2}}, v_2)}$ is of cardinality $\frac{n-3}{2}$. Then, since we have assumed that S has the maximum number of vertices from $\bigcup_{t=1}^4 Q_t$, we get,

$$|S \cap (X \cup Y)| = \begin{cases} 0; & n \equiv 1 \pmod{4}, \\ 2; & n \equiv 3 \pmod{4}. \end{cases}$$

Thus,

$$|S| \leq \begin{cases} \frac{n^2-1}{2}; & n \equiv 1 \pmod{4}, \\ \frac{n^2+3}{2}; & n \equiv 3 \pmod{4}. \end{cases}$$

Next assume that n is even and $x = (u_{\frac{n}{2}}, v_{\frac{n}{2}})$. For each $(u_k, v_l) \in Q_t$, we know that the set $N((u_k, v_l)) \cap I(x, (u_k, v_l))$ has cardinality at most four and has a vertex which is not in S . Particularly for $(u_k, v_l) \in Q_1$, this vertex is either (u_k, v_{l+1}) or (u_{k+1}, v_l) . As in the previous

Fig. 8. vv-sets of $C_6 \square C_6$ and $C_8 \square C_8$.

two proofs, partition each Q_t into diagonals D_{pq} , where $p, q \in Q_t$ are its end vertices. Then we can have at most $\sum_{pq} \lfloor |D_{pq}|/2 \rfloor$ vertices in $S \cap \bigcup_{t=1}^4 Q_t$. Thus, for $n \equiv 0 \pmod 4$ we get

$$\begin{aligned} |S \cap \bigcup_{t=1}^4 Q_t| &= |S \cap Q_1| + |S \cap Q_2| + |S \cap Q_3| + |S \cap Q_4| \\ &\leq \frac{(n-2)n}{8} + \frac{n^2}{8} + \frac{n(n+2)}{8} + \frac{n^2}{8} \\ &= \frac{n^2}{2}, \end{aligned}$$

and for $n \equiv 2 \pmod 4$ we get

$$\begin{aligned} |S \cap \bigcup_{t=1}^4 Q_t| &= |S \cap Q_1| + |S \cap Q_2| + |S \cap Q_3| + |S \cap Q_4| \\ &\leq \frac{(n-2)n}{8} + \frac{n^2-4}{8} + \frac{n(n+2)}{8} + \frac{n^2-4}{8} \\ &= \frac{n^2-2}{2}. \end{aligned}$$

Again, by the choice of S , every vertex must be S -visible from x and hence, $S \cap (X \cup Y) \subseteq \{(u_{\frac{n}{2}}, v_1), (u_{\frac{n}{2}}, v_{n-1}), (u_{\frac{n}{2}}, v_n), (u_1, v_{\frac{n}{2}}), (u_{n-1}, v_{\frac{n}{2}}), (u_n, v_{\frac{n}{2}})\}$. Also,

$$\text{in } Q_1 : |D_{(u_{\frac{n-2}{2}}, v_2)(u_2, v_{\frac{n-2}{2}})}| = \frac{n-4}{2},$$

$$\text{in } Q_2 : |D_{(u_{\frac{n-2}{2}}, v_{n-2})(u_2, v_{\frac{n+2}{2}})}| = \frac{n-4}{2} \text{ and } |D_{(u_{\frac{n-2}{2}}, v_{n-1})(u_1, v_{\frac{n+2}{2}})}| = \frac{n-2}{2},$$

$$\text{in } Q_3 : |D_{(u_{n-2}, v_{\frac{n+2}{2}})(u_{\frac{n+2}{2}}, v_{n-2})}| = \frac{n-4}{2} \text{ and } |D_{(u_{n-1}, v_{\frac{n+2}{2}})(u_{\frac{n+2}{2}}, v_{n-1})}| = \frac{n-2}{2},$$

$$\text{in } Q_4 : |D_{(u_{n-2}, v_{\frac{n-2}{2}})(u_{\frac{n+2}{2}}, v_3)}| = \frac{n-4}{2} \text{ and } |D_{(u_{n-1}, v_{\frac{n-2}{2}})(u_{\frac{n+2}{2}}, v_2)}| = \frac{n-2}{2}.$$

Then, since we have assumed that S has the maximum number of vertices from $\bigcup_{t=1}^4 Q_t$, we get

$$|S \cap (X \cup Y)| = \begin{cases} 1; & n \equiv 0 \pmod 4, \\ 2; & n \equiv 2 \pmod 4. \end{cases}$$

Thus for $n \equiv 0 \pmod 4$,

$$|S| \leq \frac{n^2+2}{2}.$$

Table 1
The vertex visibility number of square grids, prisms, and toruses.

	$n \equiv 1 \pmod 4$	$n \equiv 3 \pmod 4$	$n \equiv 0 \pmod 4$	$n \equiv 2 \pmod 4$
$P_n \square P_n$	$\frac{n^2+n-2}{2}$	$\frac{n^2+n-2}{2}$	$\frac{n^2+n-2}{2}$	$\frac{n^2+n-2}{2}$
$P_n \square C_n$	$\frac{n^2+3}{2}$	$\frac{n^2+n-2}{2}$	$\frac{2n^2+n}{4}$	$\frac{2n^2+n-2}{4}$
$C_n \square C_n$	$\frac{n^2-1}{2}$	$\frac{n^2+3}{2}$	$\frac{n^2+2}{2}$	$\frac{n^2+2}{2}$

To prove the required lower bound, let R be the set of $\lfloor |D_{pq}|/2 \rfloor$ alternate vertices from each D_{pq} ; see Figs. 7 and 8 which illustrate the construction on $P_5 \square C_5$, $P_6 \square C_6$, $P_7 \square C_7$, and $P_8 \square C_8$. Then R is an x -visibility set of cardinality $\frac{n^2-1}{2}$ for $n \equiv 1 \pmod 4$ and $R \cup \{(u_{\frac{n+1}{2}}, v_1), (u_{\frac{n+1}{2}}, v_n)\}$ is an x -visibility set of cardinality $\frac{n^2+3}{2}$ for $n \equiv 3 \pmod 4$. Also, $R \cup \{(u_{\frac{n+1}{2}}, v_1)\}$ is an x -visibility set of cardinality $\frac{n^2+2}{2}$ for $n \equiv 0 \pmod 4$ and $R \cup \{(u_{\frac{n+1}{2}}, v_n), (u_n, v_{\frac{n+1}{2}})\}$ is an x -visibility set of cardinality $\frac{n^2+2}{2}$ for $n \equiv 2 \pmod 4$. $\square$

The results of Theorems 4.2–4.4 are summarized in Table 1.

5. Concluding remarks

We conclude the paper by outlining some interesting directions for future research.

From a graph theoretical point of view, in connection with the sharp example presented in Proposition 4.1, a natural next step is to determine the vertex visibility number of the Cartesian product of an arbitrary product of finitely many complete graphs. Following our investigation of the vertex visibility property within Cartesian products, it is also of interest to extend this study to other graph products. In particular, since mutual-visibility has already been explored for strong products [23], it would be valuable to analyze the vertex visibility in the strong product of two graphs.

From a computer science point of view, given that mutual-visibility and general position problems have been successfully analyzed on many graph classes (e.g., [12,22,24,25]), it would be interesting to extend our study to other classes, in particular those suitable to describe interconnection networks, like hypercubes, butterflies, cube-connected cycles, as done for the mutual-visibility in [26].

Finally, since the mutual visibility problem is NP-complete, in [27] the authors provide approximation algorithms for this problem. We have shown, as a consequence of Theorem 2.4, that computing the vertex visibility number of a graph is also hard. Then it would be interesting to devise approximation algorithms with performance guarantees even for the new posed problem.

CRedit authorship contribution statement

Dhanya Roy: Writing – review & editing, Writing – original draft, Validation, Investigation; **Gabriele Di Stefano:** Writing – review & editing, Writing – original draft, Validation, Methodology, Investigation, Conceptualization; **Sandi Klavžar:** Writing – review & editing, Writing – original draft, Validation, Methodology, Investigation, Conceptualization; **Aparna S. Lakshmanan:** Writing – review & editing, Writing – original draft, Validation, Methodology, Investigation, Conceptualization.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors thank Ullas Chandran S V for the initial idea for this research, which greatly influenced the direction of this project. We also thank Manoj Changat for some initial discussions. Dhanya Roy thank Cochin University of Science and Technology for providing financial support under University JRF Scheme. Sandi Klavžar was supported by the Slovenian Research and Innovation Agency (ARIS) under the grants P1-0297, N1-0285, N1-0355, N1-0431, J1-70045. Gabriele Di Stefano was supported by the Italian National Group for Scientific Computation (GNCS-INdAM) and by the academic project “Monet” of the University of L’Aquila.

Data availability

No data was used for the research described in the article.

References

- [1] U.S.V. Chandran, S. Klavžar, J. Tuite, The general position problem: a survey (2025). [arXiv:2501.19385](https://arxiv.org/abs/2501.19385)
- [2] J. Araújo, M.C. Dourado, F. Protti, R. Sampaio, The iteration time and the general position number in graph convexities, *Appl. Math. Comput.* 487 (2025) 129084.

- [3] D. Korze, A. Vesel, Mutual-visibility and general position sets in Sierpiński triangle graphs, *Bull. Malays. Math. Sci. Soc.* 48 (2025) 106.
- [4] D. Roy, S. Klavžar, A.S. Lakshmanan, Mutual-visibility and general position in double graphs and in Mycielskians, *Appl. Math. Comput.* 488 (2025) 129131.
- [5] E.J. Thomas, U.S.V. Chandran, J. Tuite, G. Di Stefano, On the general position number of Mycielskian graphs, *Discrete Appl. Math.* 353 (2024) 29–43.
- [6] G. Di Stefano, Mutual visibility in graphs, *Appl. Math. Comput.* 419 (2022) 126850.
- [7] B. Bresar, I.G. Yero, Lower (total) mutual-visibility number in graphs, *Appl. Math. Comput.* 465 (2024) 128411.
- [8] C. Bujtás, S. Klavžar, J. Tian, Visibility polynomials, dual visibility spectrum, and characterization of total mutual-visibility sets, *Aequat. Math.* 99 (2025) 1883–1901.
- [9] S. Cicerone, G.D. Stefano, L. Drožek, J. Hedžet, S. Klavžar, I.G. Yero, Variety of mutual-visibility problems in graphs, *Theoret. Comput. Sci.* 974 (2023) 114096.
- [10] S. Cicerone, G. Di Stefano, S. Klavžar, I.G. Yero, Mutual-visibility problems on graphs of diameter two, *Eur. J. Combin.* 120 (2024) 103995.
- [11] S. Klavžar, D. Kuziak, J.C. Valenzuela Tripodoro, I.G. Yero, Coloring the vertices of a graph with mutual-visibility property, *Open Math.* 23 (2025) 20250193.
- [12] S. Klavžar, A.S. Lakshmanan, D. Roy, Counting largest mutual-visibility and general position sets of glued t -ary trees, *Results Math.* 80 (2025) 207.
- [13] D. Korze, A. Vesel, Mutual-visibility sets in Cartesian products of paths and cycles, *Results Math.* 79 (2024) 116.
- [14] D. Kuziak, J.A. Rodríguez-Velázquez, Total mutual-visibility in graphs with emphasis on lexicographic and Cartesian products, *Bull. Malays. Math. Sci. Soc.* 46 (2023) Paper 197.
- [15] J. Tian, S. Klavžar, Graphs with total mutual-visibility number zero and total mutual-visibility in Cartesian products, *Discuss Math. Graph Theory* 44 (2024) 1277–1291.
- [16] M.G.S. Thankachy, S.V. U. Chandran, J. Tuite, E.J. Thomas, G.D. Stefano, G. Erskine, On the vertex position number of graphs, *Discuss. Math. Graph Theory* 44 (2024) 1169–1188.
- [17] G.H. Hardy, E.M. Wright, *An Introduction to the Theory of Numbers*, Oxford University Press, 2008.
- [18] O.R. Oellermann, J. Peters-Fransen, The strong metric dimension of graphs and digraphs, *Discrete Appl. Math.* 155 (2007) 356–364.
- [19] P. Bonsma, Max-leaves spanning tree is APX-hard for cubic graphs, *J. Discrete Algorithms* 12 (2012) 14–23.
- [20] R. Solis-Oba, P. Bonsma, S. Lowski, A 2-approximation algorithm for finding a spanning tree with maximum number of leaves, *Algorithmica* 77 (2017) 374–388.
- [21] R.M. Karp, Reducibility among combinatorial problems, in: *Complexity of Computer Computations*, 1972, pp. 85–103. *Proc. Sympos.*, IBM Thomas J. Watson Res. Center, Yorktown Heights, N. Y.
- [22] S. Cicerone, G. Di Stefano, S. Klavžar, On the mutual visibility in Cartesian products and triangle-free graphs, *Appl. Math. Comput.* 438 (2023) 127619.
- [23] S. Cicerone, G.D. Stefano, S. Klavžar, I.G. Yero, Mutual-visibility in strong products of graphs via total mutual-visibility, *Discrete Appl. Math.* 358 (2024) 136–146.
- [24] S. Cicerone, G. Di Stefano, Characterizing and computing in linear time mutual-visibility parameters in distance-hereditary graphs, *Discrete Appl. Math.* 376 (2025) 359–373.
- [25] D. Roy, S. Klavžar, A.S. Lakshmanan, J. Tian, Varieties of mutual-visibility and general position on Sierpiński graphs, *Discuss. Math. Graph Theory* (2026). <https://doi.org/10.7151/dmgt.2625>
- [26] S. Cicerone, A.D. Fonso, G. Di Stefano, A. Navarra, F. Piselli, Mutual and total mutual visibility in hypercube-like graphs, *Appl. Math. Comput.* 491 (2025) 129216.
- [27] D. Bilò, A. Di Fonso, G. Di Stefano, S. Leucci, On the approximability of graph visibility problems, *Theoret. Comput. Sci.* 1067 (2026) 115766.