

On the general position number of complementary prisms

Neethu P. K.^a Ullas Chandran S. V.^a Manoj Changat^b
 Sandi Klavžar^{c,d,e}

January 8, 2020

^a Department of Mathematics, Mahatma Gandhi College, Kesavadasapuram,
 Thiruvananthapuram-695004, Kerala, India
 p.kneethu.pk@gmail.com, svuc.math@gmail.com

^b Department of Futures Studies, University of Kerala
 Thiruvananthapuram-695034, Kerala, India
 mchangat@gmail.com

^c Faculty of Mathematics and Physics, University of Ljubljana, Slovenia
 sandi.klavzar@fmf.uni-lj.si

^d Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia

^e Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia

Abstract

The general position number $\text{gp}(G)$ of a graph G is the cardinality of a largest set of vertices S such that no element of S lies on a geodesic between two other elements of S . The complementary prism $G\overline{G}$ of G is the graph formed from the disjoint union of G and its complement \overline{G} by adding the edges of a perfect matching between them. It is proved that $\text{gp}(G\overline{G}) \leq n(G) + 1$ if G is connected and $\text{gp}(G\overline{G}) \leq n(G)$ if G is disconnected. Graphs G for which $\text{gp}(G\overline{G}) = n(G) + 1$ holds, provided that both G and \overline{G} are connected, are characterized. A sharp lower bound on $\text{gp}(G\overline{G})$ is proved. If G is a connected bipartite graph or a split graph then $\text{gp}(G\overline{G}) \in \{n(G), n(G) + 1\}$. Connected bipartite graphs and block graphs for which $\text{gp}(G\overline{G}) = n(G) + 1$ holds are characterized. A family of block graphs is constructed in which the gp -number of their complementary prisms is arbitrary smaller than their order.

Key words: general position set; complementary prism; bipartite graph; split graph; block graph

AMS Subj. Class: 05C12; 05C69

1 Introduction

The general position problem in graphs was introduced in [14] as a graph theory variant of the classical, century old Dudeney's no-three-in-line problem [6] and the general position subset selection problem from discrete geometry [7, 13, 17, 19, 20]. A set S of vertices in a graph G is a *general position set* if no element of S lies on a geodesic between any two other elements of S . A largest general position set is called a *gp-set* and its size is the *general position number* (*gp-number* for short) $gp(G)$ of G . The same concept was in use two years earlier in [21] under the name *geodetic irredundant sets*.

Let us briefly recall the progress on the general position problem so far. In [14], general upper and lower bounds on the gp-number were proved as well as NP-completeness of the problem for arbitrary graphs. The gp-number of a large class of subgraphs of the infinite grid graph and of some other classes were obtained in [15]. The paper [1] gives a characterization of general position sets which is then applied in determining the gp-number of graphs of diameter 2, cographs, graphs with at least one universal vertex, bipartite graphs and their complements. Subsequently, the gp-number of complements of trees, of grids, and of hypercubes were deduced in [1]. In [8], a sharp lower bound on the gp-number of Cartesian products is proved, and the gp-number for different graph operations determined. The gp-number of Cartesian products has been further studied in [12]. In [11] the general position number has been connected with strong resolving graphs, and in [18] the general position number of Kneser graphs was investigated.

If G is a graph and \overline{G} its complement, then the *complementary prism* $G\overline{G}$ of G is the graph formed from the disjoint union of G and \overline{G} by adding the edges of a perfect matching between the corresponding vertices of G and \overline{G} [10]. For example, $C_5\overline{C}_5$ is the Petersen graph. Solely from this particular reason, but also from many additional ones, complementary prisms were studied from different perspectives. Since the Petersen graph is a key example in the theory of edge colorings, it is no surprise that the chromatic index of complementary prisms was studied in [22]. Other topics studied on complementary prisms include domination [9], cycle structure [16], complexity properties [5], spectral properties [3], convexity number [4], and b-chromatic number [2]. In this paper, we add to this list the general position problem. We proceed as follows.

The next section contains definitions, observations, and known results needed

in the rest of the paper. In Section 3 we prove that $\text{gp}(G\overline{G})$ is bounded from above by $n(G) + 1$ if G is connected and by $n(G)$ if G is disconnected, where $n(G)$ is the order of G . We also introduce the concept of 3-general position sets and apply it to derive a characterization of graphs G for which $\text{gp}(G\overline{G}) = n(G) + 1$ holds provided that both G and \overline{G} are connected. Then, in Section 4, we prove a sharp lower bound on $\text{gp}(G\overline{G})$. We follow with two sections on complementary prisms of bipartite graphs and split graphs, respectively. In both cases, provided that a bipartite graph in question is connected, $\text{gp}(G\overline{G})$ lies between $n(G)$ and $n(G) + 1$. For connected bipartite graphs we characterize the graphs G for which $\text{gp}(G\overline{G}) = n(G) + 1$ holds, while for split graphs we give two partial results about the split graphs G for which $\text{gp}(G\overline{G}) = n(G)$ holds. We conclude with Section 7 in which we give a characterization of block graphs G for which $\text{gp}(G\overline{G}) = n(G) + 1$ holds and provide a family of block graphs in which the gp-number of their complementary prisms is arbitrary smaller than their order.

2 Preliminaries

Graphs in this paper are finite and simple. Let $G = (V(G), E(G))$ be a graph. The maximum order of its complete subgraph is denoted by $\omega(G)$. Let further $\eta(G)$ denote the maximum order of an induced complete multipartite subgraph of the complement of G . The *distance* $d_G(u, v)$ between vertices u and v is the length of a shortest u, v -path. An u, v -path of minimum length is also called an u, v -*geodesic*. The *interval* $I_G[u, v]$ between u and v is the set of vertices that lie on some u, v -geodesic of G . For $S \subseteq V(G)$ we set $I_G[S] = \bigcup_{u, v \in S} I_G[u, v]$. The *eccentricity* of u is $\text{ecc}_G(u) = \max\{d_G(u, v) : v \in V(G)\}$. The *radius* and the *diameter* of G are $\text{rad}(G) = \min\{\text{ecc}_G(v) : v \in V(G)\}$ and $\text{diam}(G) = \max\{\text{ecc}_G(v) : v \in V(G)\}$, respectively. A vertex v is a *central vertex* of G if $\text{ecc}_G(v) = \text{rad}(G)$. The set of all central vertices is denoted by $C(G)$. We may simplify the above notation by omitting the index G whenever G is clear from the context. On the other hand, when we will want to emphasize that a vertex is central in a graph G , we will say that it is G -*central*.

For a characterization of general position sets we need some additional definitions. Let G be a connected graph, $S \subseteq V(G)$, and $\mathcal{P} = \{S_1, \dots, S_p\}$ a partition of S . Then \mathcal{P} is *distance-constant* if for any $i, j \in [p]$, $i \neq j$, the distance $d(u, v)$, where $u \in S_i$ and $v \in S_j$ is independent of the selection of u and v . If \mathcal{P} is a distance-constant partition, and $i, j \in [p]$, $i \neq j$, then let $d(S_i, S_j)$ be the distance between a vertex from S_i and a vertex from S_j . A distance-constant partition \mathcal{P} is *in-transitive* if $d(S_i, S_k) \neq d(S_i, S_j) + d(S_j, S_k)$ holds for arbitrary pairwise different $i, j, k \in [p]$. Now all is ready to recall the announced characterization.

Theorem 2.1 [1, Theorem 3.1] *Let G be a connected graph. Then $S \subseteq V(G)$ is a general position set if and only if the components of $G[S]$ are complete subgraphs, the vertices of which form an in-transitive, distance-constant partition of S .*

We will also make use of the following two results.

Theorem 2.2 [1, Theorem 4.1] *If $\text{diam}(G) = 2$, then $\text{gp}(G) = \max\{\omega(G), \eta(G)\}$.*

Theorem 2.3 [1, Theorem 5.1] *If G is a connected, bipartite graph with $n(G) \geq 3$, then $\text{gp}(G) \leq \alpha(G)$. Moreover, if $\text{diam}(G) \in \{2, 3\}$, then $\text{gp}(G) = \alpha(G)$.*

Let G be a graph and $G\overline{G}$ its complementary prism. Then we will consider $V(G\overline{G})$ as the disjoint union of $V(G)$ and $V(\overline{G})$. We will use the convention that if $u \in V(G) \cap V(G\overline{G})$, then its unique neighbour in $V(G\overline{G}) \cap V(\overline{G})$ will be denoted with \overline{u} and called the *partner* of u in \overline{G} . We will extend this notation to sets of vertices, that is, if $X \subseteq V(G)$, then the set of the partners of the vertices from X will be denoted with \overline{X} . Since the complementation is an idempotent operation, $\overline{G\overline{G}}$ is isomorphic to $G\overline{G}$. Note further that if $\text{diam}(G) = 2$, then $\text{diam}(G\overline{G}) = 2$, while if G is an arbitrary connected graph, then $\text{diam}(G\overline{G}) \leq 3$. Note finally that $G\overline{G}$ is always connected, no matter whether G is connected or not.

3 Upper bounds

In this section we bound $\text{gp}(G\overline{G})$ from the above by $n(G) + 1$ for connected graphs G and by $n(G)$ for disconnected graphs G , both bounds being sharp. We also characterize the graphs G for which $\text{gp}(G\overline{G}) = n(G) + 1$ holds provided that both G and \overline{G} are connected. For this purpose, the concept of 3-general position sets is introduced along the way.

Theorem 3.1 *Let G be a graph.*

- (i) *If G is connected, then $\text{gp}(G\overline{G}) \leq n(G) + 1$.*
- (ii) *If G is disconnected, then $\text{gp}(G\overline{G}) \leq n(G)$.*

Proof. (i) Let $S \subseteq V(G\overline{G})$, where $|S| \geq n(G) + 2$. By the pigeonhole principle there exist vertices $u, v \in V(G)$ such that $\{u, v, \overline{u}, \overline{v}\} \subseteq S$. Since either $uv \in E(G)$ or $\overline{u}\overline{v} \in E(\overline{G})$, we see that either $v \in I[u, \overline{v}]$ or $\overline{v} \in I[v, \overline{u}]$. It follows that S is not a general position set and we can conclude that $\text{gp}(G\overline{G}) \leq n(G) + 1$.

(ii) Let G_1, \dots, G_r , $r \geq 2$, be the components of G . Assume that $S \subseteq V(G\overline{G})$, where $|S| \geq n(G) + 1$, is a general position set of $G\overline{G}$. Using the pigeonhole principle again, there exists $v \in V(G)$ such that $\{v, \overline{v}\} \subseteq S$. We may without loss of generality

assume that $v \in V(G_1)$. Let $x \in S \cap V(G)$, $x \neq v$. Then $xv \notin E(G)$, for otherwise v would lie on a x, \bar{v} -geodesic. Moreover, $d_{G\bar{G}}(x, v) \leq 2$, for otherwise, having in mind that $\bar{x}\bar{v} \in E(G\bar{G})$, we would have that \bar{v} would lie on a x, v -geodesic. It follows that $d_{G\bar{G}}(x, v) = 2$ holds. This in particular implies that $S \cap V(G) \subseteq V(G_1)$. Using a parallel argument we infer that if $\bar{y} \in S \cap V(\bar{G})$, then $d_{G\bar{G}}(\bar{y}, \bar{v}) = 2$. This in turn implies that $\bar{y}\bar{v} \notin E(\bar{G})$ so that $yv \in E(G)$. It follows that $\{y : \bar{y} \in S \cap V(\bar{G})\} \subseteq N_G[v]$.

We have thus proved that $S \subseteq V(G_1) \cup V(\bar{G}_1)$. Since G_1 is connected, the proof of (i) restricted to $G_1\bar{G}_1$ implies that v is the unique vertex of $S \cap V(G)$ such that $\bar{v} \in S$. Hence $|S| \leq n(G_1) + 1 \leq n(G)$, a contradiction. Hence $\text{gp}(G\bar{G}) \leq n(G)$. \square

To quickly demonstrate that the bounds of Theorem 3.1 are sharp, consider the following sporadic examples. First, as observed in [14] for the Petersen graph, $\text{gp}(C_5\bar{C}_5) = 6$, which demonstrates sharpness of (i). Second, $\text{gp}(\bar{P}_2\bar{P}_2) = 2$ demonstrates sharpness of (ii).

Our next goal is to characterize the graphs G such that both G and \bar{G} are connected and $\text{gp}(G\bar{G}) = n(G) + 1$. By the above, the Petersen graph belongs to this family.

Lemma 3.2 *Let G be a graph with $n(G) \geq 2$ and such that both G and \bar{G} are connected. If $\text{gp}(G\bar{G}) = n(G) + 1$, then the following properties hold.*

(i) $\text{rad}(G) = 2$.

(ii) *If S is a gp-set of $G\bar{G}$, then there exists a G -central vertex $v \in S$ such that $S \cap V(G) = \{u \in V(G) : d_G(u, v) = 2\} \cup \{v\}$ and $S \cap V(\bar{G}) = \bar{N}_G[v]$.*

Proof. Let S be a gp-set of $G\bar{G}$. Then by lemma's assumption, $|S| = n(G) + 1$, and hence there exists a vertex $v \in V(G)$ such that $\{v, \bar{v}\} \subseteq S$.

Suppose that there exists a vertex $x \in S \cap V(G)$ for which $d_G(x, v) \geq 3$ holds. Then the path on the vertices x, \bar{x}, \bar{v}, v is a x, v -geodesic in $G\bar{G}$ containing the vertex \bar{v} , a contradiction. Hence $d_G(x, v) \leq 2$ for all $x \in S \cap V(G)$, $x \neq v$. Since \bar{G} is connected, $V(G) \setminus N_G(v) \neq \emptyset$, for otherwise \bar{v} would be an isolated vertex in \bar{G} . We have shown that $\text{ecc}_G(v) = 2$ which in turn implies (i).

Theorem 2.1 implies that $N_G(v) \cap S = \emptyset$ and $S \cap V(\bar{G}) \subseteq \bar{N}_G[v]$. Since $|S| = n(G) + 1$ and because by the proof of Theorem 3.1(i), the vertex v is the unique vertex of $S \cap V(G)$ such that its partner in \bar{G} also belongs to S , we must have that $S \cap V(G) = V(G) \setminus N_G(v)$ and $S \cap V(\bar{G}) = \bar{N}_G[v]$. Combining this with the already proved property (i), the assertion (ii) follows. \square

The condition (i) of Lemma 3.2 is not sufficient for G to have $\text{gp}(G\bar{G}) = n(G) + 1$. To see it, consider the double star G shown in Fig. 1.




Figure 1: The double star G

Clearly, $\text{rad}(G) = 2$ and both G and \overline{G} are connected. If $\text{gp}(G\overline{G}) = 7$, and S is a gp-set of $G\overline{G}$, then by Lemma 3.2 either $v \in S$ or $u \in S$. Assume without loss of generality that $v \in S$. Using Lemma 3.2 again we get that $S = \{v, \overline{v}, u_1, u_2, \overline{u}, \overline{v}_1, \overline{v}_2\}$. But then the path on the vertices $\overline{u}, u, v, \overline{v}$ is a geodesic containing three vertices of S , a contradiction.

To state the announced characterization, we introduce the following concept. A set S of vertices in a graph G is a *3-general position set* if no three vertices from S lie on a common geodesic of length at most 3.

Lemma 3.3 *Let G be a graph. A set $S \subseteq V(G)$ is a 3-general position set if and only if the components of $G[S]$ are complete subgraphs, and if $d(u, v) = 2$, where u and v lie in respective components Q and Q' of $G[S]$, then $d(x, y) = 2$ for all $x \in Q$ and all $y \in Q'$.*

Proof. Let $S \subseteq V(G)$ be a 3-general position set. Then each component of $G[S]$ must be complete, for otherwise a non-complete component would contain an induced P_3 which is a geodesic. Suppose now that $d(u, v) = 2$, where u and v lie in components Q and Q' of $G[S]$. Let $x \in Q$ and $y \in Q'$. Clearly, $1 \leq d(x, y) \leq 3$. Suppose $d(x, y) = 1$. Assuming without loss of generality that $x \neq u$ we see that the vertices y, x, u induce a geodesic, a contradiction. Suppose $d(x, y) = 3$. Again assuming without loss of generality that $x \neq u$ we see that the vertices x, u, v lie on a common geodesic of length 3, another contradiction. Hence $d(x, y) = 2$.

Conversely, suppose that the components of $G[S]$ are complete subgraphs and that the distance condition is fulfilled. Suppose on the contrary that S is not a 3-general position set, that is, there are three vertices $u, v, w \in S$ that lie on a common geodesic P of length at most 3. As the components of $G[S]$ are complete, P cannot be of length 2, so we may without loss of generality assume that P is the path on vertices u, v, x, w for some vertex $x \in V(G)$. Then u and w lie in different cliques of $G[S]$, say $u \in Q$ and $w \in Q'$. Clearly, then also $v \in Q$. Since $d(v, w) = 2$, the distance condition implies that $d(u, w) = 2$ as well. But this contradicts the fact that P is a geodesic. \square

We are now ready for the announced characterization.

Theorem 3.4 *Let G be a graph with $n(G) \geq 2$ and such that both G and \overline{G} are connected. Then $\text{gp}(G\overline{G}) = n(G) + 1$ if and only if $\text{rad}(G) = 2$ and there exists $v \in C(G)$ such that*

(i) $\overline{N_G(v)}$ is a 3-general position set in \overline{G} and $N_G^2(v)$ is a 3-general position set in G , and

(ii) for each $x \in N_G(v)$ there exists $y \in N_G^2(v)$ such that $xy \notin E(G)$.

Proof. First suppose that $\text{gp}(G\overline{G}) = n(G) + 1$. Let S be a gp-set of G , so that $|S| = n(G) + 1$. Then by Lemma 3.2, $\text{rad}(G) = 2$ and there exists a vertex $v \in C(G)$ such that $S \cap V(G) = N_G^2(v)$ and $S \cap V(\overline{G}) = \overline{N_G(v)}$ are both general position sets in $G\overline{G}$. Hence $\overline{N_G(v)}$ is a 3-general position set in \overline{G} and $N_G^2(v)$ is a 3-general position set in G . Thus (i) holds for the vertex v . Let now $x \in N_G(v)$ and assume that $xy \in E(G)$ for every $y \in N_G^2(v)$. This implies that $d_{G\overline{G}}(\overline{x}, \overline{v}) = 3$. Since $d_{G\overline{G}}(\overline{x}, v) = 2$ and v, \overline{v} are in the same clique of $G\overline{G}[S]$, Lemma 3.3 implies that also $d_{G\overline{G}}(\overline{x}, \overline{v}) = 2$ must hold. This contradiction proves property (ii) for the vertex v .

Conversely, suppose that $\text{rad}(G) = 2$ and that there exists a vertex v in $C(G)$ satisfying properties (i) and (ii). We claim that the set $S = N_G^2(v) \cup \overline{N_G(v)} \cup \{v, \overline{v}\}$ induces an in-transitive, distance-constant partition into cliques. By Lemma 3.3, each of $N_G^2(v)$ and $\overline{N_G(v)}$ induces an in-transitive, distance-constant partition into cliques. Now, for any $u \in V(G)$ and $\overline{w} \in V(\overline{G})$ with $u \neq w$, we have $d_{G\overline{G}}(u, \overline{w}) = 2$. Which implies that $N_G^2(v) \cup \overline{N_G(v)} \cup \{v\}$ has an intransitive-distance constant partition into cliques. Next, for any $y \in N_G(v)$ we have $d_{G\overline{G}}(\overline{y}, v) = 2$ and by condition (ii) of our assumption, $d_{G\overline{G}}(\overline{y}, \overline{v}) = 2$. Hence S induces an in-transitive, distance-constant partition into cliques. Since $\text{ecc}_G(v) = 2$, we have that $|S| = n(G) + 1$. Hence by Theorem 2.1, S is a general position set of size $n(G) + 1$ and so $\text{gp}(G\overline{G}) = n(G) + 1$ by Theorem 3.1. \square

As an example consider again the Petersen graph $P = C_5\overline{C_5}$. Let the vertices of the C_5 be v_1, v_2, v_3, v_4, v_5 . Observe that $\text{rad}(C_5) = 2$ and $C(C_5) = V(C_5)$. Select v_1 as a vertex from $C(C_5)$. Then $N_{C_5}^2(v_1) = \{v_3, v_4\}$ and $\overline{N_{C_5}(v_1)} = \{\overline{v_2}, \overline{v_5}\}$. It is now straightforward to check that the conditions of Theorem 3.4 are fulfilled, hence $\text{gp}(P) = 6$ with $\{v_1, \overline{v_1}, v_3, v_4, \overline{v_2}, \overline{v_5}\}$ being its gp-set.

Next, we give an infinite family of graphs that satisfies Theorem 3.4. Let G be the graph obtained from the path on vertices u, v, w and disjoint cliques K_{n_1}, \dots, K_{n_r} and K_{m_1}, \dots, K_{m_s} by joining u to all the vertices of K_{n_i} , $i \in [r]$, and w to all the vertices of K_{m_j} , $j \in [s]$. Then $\text{rad}(G) = 2$ and $C(G) = \{v\}$. Since the sets $N_G(v) = \{u, w\}$ and $N_G^2(v) = V(G) \setminus \{u, v, w\}$ satisfy the conditions of Theorem 3.4, we conclude that $\text{gp}(G\overline{G}) = n(G) + 1$.

4 A lower bound

In this section we prove a sharp lower bound on $\text{gp}(G\overline{G})$. For this sake we define a new invariant $\overline{\text{gp}}_3(G)$ of a graph G as follows:

$$\overline{\text{gp}}_3(G) = \max\{\text{gp}_3(\overline{G}[V(\overline{G}) \setminus \overline{S}]) : S \text{ gp}_3\text{-set of } G\}.$$

Theorem 4.1 *If G is a graph, then*

$$\text{gp}(G\overline{G}) \geq \max\{\text{gp}_3(G) + \overline{\text{gp}}_3(G), \text{gp}_3(\overline{G}) + \overline{\text{gp}}_3(\overline{G})\}.$$

Moreover, the bound is sharp.

Proof. Let S be a gp_3 -set of G and let Q_1, \dots, Q_k be the complete subgraphs of $G[S]$ according to Lemma 3.3. For $x \in Q_i$ and $y \in Q_j$, where $i \neq j$, $d_G(x, y) \in \{2, 3\}$ and hence $d_G(x, y) = d_{G\overline{G}}(x, y)$. This already implies that the cliques Q_1, \dots, Q_k form an in-transitive, distant-constant partition of S in $G\overline{G}$. Consider now a gp_3 -set \overline{T} of $\overline{G}[V(\overline{G}) \setminus \overline{S}]$ and let R_1, \dots, R_t be the complete subgraphs of $\overline{G}[\overline{T}]$ according to Lemma 3.3. Then, by the same argument as we used for Q_1, \dots, Q_k , we infer that the cliques R_1, \dots, R_t form an in-transitive, distant-constant partition of \overline{T} in $G\overline{G}$. Let now $x \in Q_i$, $i \in [k]$, and $y \in R_j$, $j \in [t]$. Then by the structure of $G\overline{G}$ it follows that $d_{G\overline{G}}(x, y) = 2$. It follows that the cliques $Q_1, \dots, Q_k, R_1, \dots, R_t$ form an in-transitive, distant-constant partition in $G\overline{G}$. With Theorem 2.1 in hands we have thus proved that $\text{gp}(G\overline{G}) \geq \text{gp}_3(G) + \overline{\text{gp}}_3(G)$. Starting with a gp_3 -set of \overline{G} and repeating the argument from the paragraph above, we also get that $\text{gp}(G\overline{G}) \geq \text{gp}_3(\overline{G}) + \overline{\text{gp}}_3(\overline{G})$.

By a simple argument we can see that if $n \geq 2$, then $\text{gp}(K_n \overline{K}_n) = n$. Since $\text{gp}_3(K_n) = n$ (as well as $\text{gp}_3(\overline{K}_n) = n$), the bound is sharp. \square

From Theorem 4.1 it is clear that $\text{gp}(G\overline{G}) \geq \max\{\text{gp}_3(G), \text{gp}_3(\overline{G})\}$. We next characterize the connected graphs for which the equality holds.

Theorem 4.2 *Let G be a connected graph. Then $\text{gp}(G\overline{G}) = \max\{\text{gp}_3(G), \text{gp}_3(\overline{G})\}$ if and only if G is a complete multipartite graph.*

Proof. Let $\text{gp}(G\overline{G}) = \max\{\text{gp}_3(G), \text{gp}_3(\overline{G})\}$.

Suppose first that $\text{gp}(G\overline{G}) = \max\{\text{gp}_3(G), \text{gp}_3(\overline{G})\} = \text{gp}_3(G)$. Let S be a maximum 3-general position set of G . Then S is also a maximum general position set of $G\overline{G}$. If $S \neq V(G)$, then $S \cup \{\overline{v}\}$ is a general position set of $G\overline{G}$ for all $v \in V(G) \setminus S$. This is impossible. Hence $\text{gp}_3(G) = |S| = n(G)$. Then $G[S]$ must be connected and from Lemma 3.3 we conclude that $G = K_n$.

Suppose second that $\text{gp}(G\overline{G}) = \max\{\text{gp}_3(G), \text{gp}_3(\overline{G})\} = \text{gp}_3(\overline{G})$. If \overline{G} is connected, then as above we have that $G = \overline{K}_n$. This is a contradiction to the fact that

G is connected. Hence \overline{G} must be disconnected. Let S be a maximum 3-general position set of \overline{G} . Then S is also a maximum general position set of $G\overline{G}$. If $S \neq V(\overline{G})$, then $S \cup \{v\}$ is a general position set of $G\overline{G}$ for all $\overline{v} \in V(\overline{G}) \setminus S$. This is impossible and so $\text{gp}_3(\overline{G}) = |S| = n(\overline{G})$. By Lemma 3.3, each component of \overline{G} must be a clique. This shows that G is a complete multipartite graph.

Conversely, Let G be a complete multipartite graph. Then \overline{G} is a disjoint union of cliques. This shows that $V(G)$ is a general position set of $G\overline{G}$. Hence it follows from Theorem 3.1(ii) that $\text{gp}(G\overline{G}) = n(G) = \max\{\text{gp}_3(G), \text{gp}_3(\overline{G})\}$. \square

Note that if $\text{gp}(G) \geq n(G) - 2$, then Theorem 4.1 readily implies that $\text{gp}(G\overline{G}) \geq n(G)$. On the other hand, the bound of Theorem 4.1 is never larger than $n(G)$, hence the existence of graphs G for which the equality $\text{gp}(G\overline{G}) = n(G) + 1$ holds (cf. Theorem 3.4), implies that the bound is not sharp in general. Additional sharpness cases for the bound of Theorem 4.1 will be presented in the subsequent sections.

5 Bipartite graphs

In this section we give our attention to the complementary prisms of bipartite graphs. If $G = (V(G), E(G))$ is a bipartite graph with bipartition $V(G) = A \cup B$, then we write G as a triple $(A, B, E(G))$. In $G = (A, B, E(G))$, set

$$U_G = \{u \in A : \deg(u) = |B|\} \cup \{v \in B : \deg(v) = |A|\}.$$

Theorem 5.1 *If $G = (A, B, E(G))$ is a connected, bipartite graph, then $n(G) \leq \text{gp}(G\overline{G}) \leq n(G) + 1$. Moreover, $\text{gp}(G\overline{G}) = n(G) + 1$ if and only if $\text{rad}(G) = 2$ and $C(G)$ is an independent set.*

Proof. By Theorem 3.1, $\text{gp}(G\overline{G}) \leq n(G) + 1$. Since independent sets and cliques are 3-general position sets in any graph, it follows from Theorem 4.1 that the set $S = A \cup \overline{B}$ is a general position set in $G\overline{G}$. Thus $\text{gp}(G\overline{G}) \geq n(G)$. It thus remains to characterize the graphs G for which $\text{gp}(G\overline{G}) = n(G) + 1$ holds.

Suppose that $\text{gp}(G\overline{G}) = n(G) + 1$. Then, in view of Theorem 3.1, both G and \overline{G} are connected. Let S be a general position set in $G\overline{G}$ of size $n(G) + 1$. Then by Lemma 3.2, we have that $\text{rad}(G) = 2$ and there exists a G -central vertex $v \in S$ such that $S \cap V(G) = \{u \in V(G) : d_G(u, v) = 2\}$ and $S \cap V(\overline{G}) = \overline{N_G[v]}$. Now, without loss of generality we may assume that $v \in A$. Since $\text{ecc}_G(v) = 2$ and G is bipartite, it follows that $v \in U_G$ and so $S \cap V(\overline{G}) = \overline{N_G[v]} = \overline{B} \cup \{\overline{v}\}$. Now, we claim that $U_G \subseteq A$. Assume on the contrary that there exists a vertex $u \in B \cap U_G$. Then $d_{G\overline{G}}(\overline{u}, \overline{v}) = 3$ and $\overline{u}, \overline{v} \in S$. Moreover, $\overline{u}, u, v, \overline{v}$ is a $\overline{u}, \overline{v}$ -geodesic in G containing the vertex v . This leads to a contradiction to the fact that S is a general position set in

$G\overline{G}$. Thus $U_G \subseteq A$. This shows that $\text{rad}(G) = 2$ and that $C(G)$ is an independent set in G .

Conversely, suppose that $\text{rad}(G) = 2$ and $C(G)$ is independent in G . Since $\text{rad}(G) = 2$ and G is bipartite, $C(G) = U_G$. Because $C(G)$ is independent, and $C(G) = U_G$, it follows that either $C(G) \subseteq A$ or $C(G) \subseteq B$, say $C(G) \subseteq A$. Let v be a vertex in $C(G)$. We claim that the set $S = A \cup \overline{B} \cup \{\overline{v}\}$ is a general position set in $G\overline{G}$. Now, the set $N_G^2(v) = A \setminus \{v\}$ is an independent set in G and so it is a 3-general position set in $G\overline{G}$. Also, $N_G(v) = \overline{B}$ is a clique in \overline{G} and so it is a 3-general position set in $G\overline{G}$. Moreover, since $v \in U_G = C(G) \subseteq A$, we have that $d_{G\overline{G}}(\overline{y}, \overline{v}) = 2$ for all $y \in N_G(v)$. Hence it follows from Theorem 3.4 that S is a general position set in $G\overline{G}$ and hence $\text{gp}(G\overline{G}) = n(G) + 1$. \square

In the rest of the section we present the general position number of complementary prism of some standard families of bipartite graphs. Let T be a tree. Then $\text{gp}(T)$ is the number of its leaves [1, 21], and $\text{gp}(\overline{T}) = \max\{\alpha(T), \Delta(T) + 1\}$ [1]. Since $1 \leq |C(T)| \leq 2$, Theorem 5.1 implies that $\text{gp}(T\overline{T}) = n(G) + 1$ if and only if $|C(T)| = 1$ and $\text{rad}(T) = 2$. This is possible only when T has diameter 4. We have thus deduced:

Corollary 5.2 *If T is a tree, then*

$$\text{gp}(T\overline{T}) = \begin{cases} n(G) + 1; & \text{diam}(T) = 4, \\ n(G); & \text{otherwise.} \end{cases}$$

The *Cartesian product* $G \square H$ of graphs (factors) G and H has $V(G \square H) = V(G) \times V(H)$ and vertices (g, h) and (g', h') are adjacent if either $g = g'$ and $hh' \in E(G)$, or $h = h'$ and $gg' \in E(G)$. For $n, m \geq 2$ set $P_{nm} = P_n \square P_m$. In [15] it was proved that $\text{gp}(P_{nm}) = 4$ for $n, m \geq 3$, while in [1] the following result was deduced:

$$\text{gp}(\overline{P}_{nm}) = \begin{cases} 4; & n = m = 2, \\ \lceil \frac{n}{2} \rceil \lceil \frac{m}{2} \rceil + \lfloor \frac{n}{2} \rfloor \lfloor \frac{m}{2} \rfloor; & \text{otherwise.} \end{cases}$$

With the help of Theorem 5.1, we can add to these results the following.

Corollary 5.3 *If $n, m \geq 2$, then*

$$\text{gp}(P_{nm}\overline{P}_{nm}) = \begin{cases} 10; & n = m = 3, \\ nm; & \text{otherwise.} \end{cases}$$

As the last subclass of bipartite graphs consider hypercubes. Recall that the n -cube Q_n is the n -fold Cartesian product of K_2 . Once more applying Theorem 5.1 we get:

Corollary 5.4 *If $n \geq 2$, then $\text{gp}(Q_n\overline{Q}_n) = 2^n$.*

6 Split graphs

A graph $G = (V(G), E(G))$ is a *split graph* if $V(G)$ can be partitioned into a clique C and an independent set I . If so, the pair (C, I) is a *split partition* of G and we write $G = (C, I, E(G))$.

Theorem 6.1 *If $G = (C, I, E(G))$ is a split graph, then $n(G) \leq gp(\overline{G\overline{G}}) \leq n(G) + 1$. Moreover, the following hold.*

- (i) *If $\deg_G(x) \geq |C| + 1$ for all $x \in C$ and $\deg_G(y) \leq |C| - 2$ for all $y \in I$, then $gp(\overline{G\overline{G}}) = n(G)$.*
- (ii) *If $gp(\overline{G\overline{G}}) = n(G)$, then \overline{G} is disconnected or $\deg_G(x) \geq |C|$ for all $x \in C$.*

Proof. By Theorem 4.1, the set $H = C \cup \overline{I}$ is a general position set of $\overline{G\overline{G}}$ and hence $gp(\overline{G\overline{G}}) \geq n(G)$. The upper bound again follows from Theorem 3.1.

(i) Suppose that $\deg_G(x) \geq |C| + 1$ for all $x \in C$ and $\deg_G(y) \leq |C| - 2$ for all $y \in I$. By way of contradiction suppose that $gp(\overline{G\overline{G}}) = n(G) + 1$. Then by Lemma 3.2, there exists a G -central vertex v such that $S = N_G^2(v) \cup \overline{N_G(v)} \cup \{v, \overline{v}\}$ is a general position set of $\overline{G\overline{G}}$. We consider two cases.

Case 1: $v \in C$.

Then $\deg_G(v) \geq |C| + 1$ and let z be a neighbour of v in I . Now, since $\overline{N_G[v]} \subseteq S$, we have that $\overline{z} \in S$. By our hypothesis, $\deg_G(z) \leq |C| - 2$. Thus we can choose two distinct vertices, say u_1 and u_2 in C , such that both u_1 and u_2 are non-adjacent to z in G . This shows that $\overline{z} \in I_{\overline{G\overline{G}}}[\overline{u_1}, \overline{u_2}] \subseteq I_{\overline{G\overline{G}}}[S]$. This is a contradiction to the fact that S is a general position set in $\overline{G\overline{G}}$.

Case 2: $v \in I$.

In this case $\deg_G(v) \leq |C| - 2$. Let u be a vertex in C such that u and v are non-adjacent in G . Then by Lemma 3.2, $u \in S$. Then $\deg_G(u) \geq |C| + 1$, thus we can choose two distinct vertices, say v_1 and v_2 in I such that both v_1 and v_2 are adjacent to u in G . This shows that $u \in I_{\overline{G\overline{G}}}[v_1, v_2] \subseteq I_{\overline{G\overline{G}}}[S]$. This is a contradiction to the fact that S is a general position set in $\overline{G\overline{G}}$.

Since in both cases we got a contradiction, we conclude that $gp(\overline{G\overline{G}}) = n(G)$.

(ii) Assume that $gp(\overline{G\overline{G}}) = n(G)$ and \overline{G} is connected. Then G has no universal vertex and so $\text{rad}(G) = 2$. Suppose that $\deg_G(v) = |C| - 1$ for some $v \in C$. Then v is a G -central vertex and $\overline{N_G(v)} = \overline{C}$ is a 3-general position set in \overline{G} and $N_G^2(v) = I$ is a 3-general position set of G . Moreover, since G has no universal vertices, we have that for each $x \in C$ there exists $y \in I$ such that $xy \notin E(G)$. Hence by Theorem 3.4, $gp(\overline{G\overline{G}}) = n(G) + 1$, a contradiction. \square




Figure 2: Split graphs

The converse for neither of Theorem 6.1(i) and (ii) is true. For this sake consider the split graphs G_1 and G_2 shown in Fig. 2.

Consider first the split graph G_1 with $C = \{u_1, u_2, u_3\}$ and $I = \{v_1, v_2, v_3\}$. The center of G_1 is $C(G_1) = \{u_1, u_2, u_3\}$. Since no central vertex satisfies conditions of Theorem 3.4, $\text{gp}(G_1\overline{G}_1) = n(G_1) = 6$. Also, since $\deg_{G_1}(x) = 3 < 4 = |C| + 1$ for all $x \in C(G_1)$, we infer that the converse of Theorem 6.1(i) does not hold. Next, in G_2 , the vertex u_1 is a G_2 -central vertex which satisfies the conditions of Theorem 3.4. Hence $\text{gp}(G_2\overline{G}_2) = 7 = n(G) + 1$, whereas both G_2 and \overline{G}_2 are connected and have no simplicial vertices in the corresponding cliques. Hence the converse of Theorem 6.1(ii) also does not hold.

Again, consider a split graph $G = (C, I, E(G))$ such that both G and \overline{G} are connected. Then it is clear from Theorem 6.1 that $\text{gp}(G\overline{G}) = n(G) + 1$ if either C or \overline{I} contains a simplicial vertex.

We conclude the section with the natural problem arising from Theorem 6.1.

Problem 6.2 *Characterize the split graphs G for which $\text{gp}(G\overline{G}) = n(G) + 1$ holds.*

7 Block graphs

A graph is a *block graph* if every maximal 2-connected component is a clique. In a block graph each vertex is either a simplicial vertex or a cut vertex. The following theorem gives a characterization of block graphs for which the bound in Theorem 3.1(i) is attained.

Theorem 7.1 *Let G be a block graph. Then $\text{gp}(G\overline{G}) = n(G) + 1$ if and only if $\text{rad}(G) = 2$ and there exists a G -central vertex v such that $N_G(v)$ is either a clique or an independent set, and $N_G(v)$ contains at least two cut vertices of G .*

Proof. First suppose that $\text{gp}(G\overline{G}) = n(G) + 1$. Then by Lemma 3.2, we have $\text{rad}(G) = 2$ and G contains a central vertex v such that $N_G^2(v) \cup \overline{N}_G(v) \cup \{v, \overline{v}\}$ is

a gp-set of $G\overline{G}$. We claim that $N_G(v)$ is either a clique or an independent set. Let B_1, \dots, B_k be the blocks of G containing v . Then we have $N_G(v) = \cup_{i=1}^k N_{B_i}(v)$. Since each $N_{B_i}(v)$ induces a clique, $\overline{N_G(v)}$ induces a multipartite set. By Theorem 3.4, we have that $\overline{N_G(v)}$ is a 3-general position set in \overline{G} . Thus either $k = 1$ or $\overline{N_G(v)}$ is a clique. This proves the claim. Next we prove that $N_G(v)$ contains at least two cut vertices of G . Since v is a central vertex and $\text{rad}(G) = 2$, we have that $|N_G(v)| \geq 2$. Let $y \in N_G(v)$. Since $\text{ecc}_G(v) = 2$, it follows from Theorem 3.4 that corresponding to the vertex y there exist vertices $x \in N_G^2(v)$ and $w \in N_G(v)$ such that $xw \in E(G)$ and $xy \notin E(G)$. Similarly, corresponding to the vertex w there exist vertices $x' \in N_G^2(v)$ and $z \in N_G(v)$ such that $x'z \in E(G)$ and $x'w \notin E(G)$. Since G is block graph, both w and z must be cut vertices in G .

Conversely, suppose that $\text{rad}(G) = 2$ and G contains a central vertex v such that $N_G(v)$ is either a clique or an independent set, and $N_G(v)$ contains at least two cut vertices of G . Since $\overline{N_G(v)}$ is either a clique or an independent set in G , it is a 3-general position set of \overline{G} . Let B'_1, \dots, B'_r be the blocks of G which do not contain v . Then each $B'_i, i \in [r]$, must contain a vertex from $N_G(v)$, say u_i . Then $N_G^2(v) = \cup_{i=1}^r V(B'_i) \setminus \{u_i\}$. Since $V(B'_i) \setminus \{u_i\}$ induces a clique in G , we have that $N_G^2(v)$ is a union of disjoint cliques in G . This implies that $N_G^2(v)$ is a 3-general position set in G . We now claim that for each $y \in N_G(v)$, there exists a vertex x in $N_G^2(v)$ such that $xy \notin E(G)$. Suppose on the contrary a vertex $y \in N_G(v)$ is adjacent to each vertex in $N_G^2(v)$. But this would mean that $N_G(v)$ has at most one cut vertex, a contradiction. Hence, by Theorem 3.4, we conclude that $N_G^2(v) \cup \overline{N_G(v)} \cup \{v, \overline{v}\}$ is a gp-set of $G\overline{G}$ and so $\text{gp}(G\overline{G}) = n(G) + 1$. \square

In the rest of the section we present an infinite family of block graphs such that the gp-number of their complementary prisms is arbitrary smaller than their order. Let G_0 be the complete graph K_3 with the vertex set $\{v_1, u_1, v_2\}$. For $k \geq 1$, the graph G_k is obtained from G_{k-1} by adding two new adjacent vertices u_{k+1} and v_{k+2} and joining both u_{k+1} and v_{k+2} to the vertex v_{k+1} . Note that G_k is a block graph with $k + 1$ blocks B_1, \dots, B_{k+1} , where each block B_i is a triangle with $V(B_i) = \{v_i, u_i, v_{i+1}\}$. Note further that $E_k = \{v_1, v_{k+2}\} \cup \{u_1, \dots, u_{k+1}\}$ is the set of simplicial vertices of G_k and that the set of remaining vertices, that is $A_k = \{v_2, \dots, v_{k+1}\}$, is the set of cut vertices of G_k . Clearly, A_k induces a path of length $k - 1$, let X_k and Y_k form the bipartition of A_k where $|X_k| \geq |Y_k|$.

Theorem 7.2 *If $k \geq 5$, then $\text{gp}(G_k\overline{G}_k) = n(G_k) - \lfloor \frac{k}{2} \rfloor$.*

Proof. Let $k \geq 5$ and let E_k, A_k, X_k , and Y_k be the sets of vertices as defined before the theorem. Using the proof of Theorem 4.1 we infer that $S = E_k \cup \overline{X}_k$ is a general position set of $G_k\overline{G}_k$ of size $n(G_k) - \lfloor \frac{k}{2} \rfloor$. Hence $\text{gp}(G_k\overline{G}_k) \geq n(G_k) - \lfloor \frac{k}{2} \rfloor$.

To prove the other inequality, assume on the contrary that there exists a general position set S of $G_k \overline{G}_k$ with $|S| > n(G_k) - \lfloor \frac{k}{2} \rfloor$. Let $M = S \cap V(G_k)$ and $\overline{N} = S \cap V(\overline{G}_k)$. Then both M and \overline{N} are 3-general position sets of G_k and \overline{G}_k , respectively. Since $|S| > n(G_k) - \lfloor \frac{k}{2} \rfloor$, it follows that either $|M| > k + 3$ or $|N| = |\overline{N}| > \lfloor \frac{k}{2} \rfloor$.

Claim A: $\text{gp}_3(G_k) = k + 3$ for all $k \geq 0$.

We proceed by induction on k , the cases $k = 0$ and $k = 1$ being easily verified. Assume that $\text{gp}_3(G_i) = i + 3$ for all i with $2 \leq i < k$ and consider G_k . Let H_k be a maximum 3-general position set of G_k and suppose that $|H_k| > k + 3$. Then both $u_{k+1}, v_{k+2} \in H_k$. Otherwise, if $u_{k+1} \notin H_k$ (say), then $H_k \setminus \{v_{k+2}\}$ is a 3-general position set of G_{k-1} of size at least $k + 3$. This is a contradiction to the induction hypothesis. Now, let $H_{k-1} = H_k \setminus \{u_{k+1}, v_{k+2}\}$. Then H_{k-1} is a 3-general position set of G_{k-1} of size at least $k + 2$. hence by induction H_{k-1} is a maximum 3-general position set of G_{k-1} of size $k + 2$. Again, since $\text{gp}(G_{k-2}) = k + 1$, it follows that either $u_k \in H_{k-1}$ or $v_{k+1} \in H_{k-1}$. Recall that H_k is a 3-general position set in G_k . This shows that H_{k-1} contains exactly one vertex from the set $\{u_k, v_{k+1}\}$, say u_k . But this leads to the fact that $H_{k-1} \cup \{v_{k+1}\}$ is a 3-general position set of G_{k-1} of size $k + 3$, a contradiction, and the claim is proved.

Since M is a 3-general position set in G_k , Claim A implies that $|M| \leq k + 3$. This shows that $|N| > \lfloor \frac{k}{2} \rfloor$. We consider the following three cases.

Case 1: $N \subseteq E_k$.

Recall that N contains at most one vertex from M . First suppose that $M \cap N \neq \emptyset$. In this case there exists a simplicial vertex x such that $x \in M$ and $\overline{x} \in \overline{N}$. Let y be a vertex from N distinct from x . If x and y are non-adjacent in G_k , then $x, \overline{x}, \overline{y}$ is a shortest path in $G_k \overline{G}_k$ with $x, \overline{x}, \overline{y} \in M \cup \overline{N} = S$. This is impossible. Hence x and y are adjacent in G_k . Now, since $x, y \in N \subseteq E_k$, it follows that either $x, y \in B_1$, or $x, y \in B_{k+1}$. This shows that $\overline{E_k} \setminus \{x, y\} \subseteq I_{G_k \overline{G}_k}[\overline{x}, \overline{y}] \subseteq I_{G_k \overline{G}_k}[\overline{N}]$ and so $N = \{x, y\}$. This is a contradiction to the fact that $|N| > \lfloor \frac{k}{2} \rfloor \geq 3$. So, assume that $M \cap N = \emptyset$. Because \overline{N} is a 3-general position set in \overline{G}_k , we have that the components of the induced subgraph of \overline{N} in \overline{G}_k are cliques. Hence N induces a complete multipartite graph in G_k . Since $N \subseteq E_k$, it follows that $N \subseteq E_k \setminus \{v_1, v_{k+1}\}$ and so $|N| \leq k + 1$. On the other hand, since M is a 3-general position set in G_k and A_k induces a path of order k , it follows that $|M \cap A_k| \leq \lfloor \frac{k}{2} \rfloor$.

Thus $|S| = |M| + |\overline{N}| = |M| + |N| = |M \cup N| = |(M \cup N) \cap E_k| + |(M \cup N) \cap A_k| \leq k + \lfloor \frac{k}{2} \rfloor + 3 = n(G_k) - \lfloor \frac{k}{2} \rfloor$.

Case 2: $N \subseteq A_k$.

Recall that N induces a complete multipartite graph in G_k and A_k induces a path of order k . Hence it is clear that either $N \subseteq X_k$ or $N \subseteq Y_k$ and so $|N| \leq \lfloor \frac{k}{2} \rfloor$. Hence this case cannot occur.

Case 3: $N \cap E_k \neq \emptyset$ and $N \cap A_k \neq \emptyset$.

Claim B: N is an independent set in G_k for all $k \geq 5$.

Assume on the contrary that there exists adjacent vertices u and v in N and consider the following cases.

Subcase 3.1: u is a simplicial vertex and v is a cut vertex.

We may assume that $u = u_i$ and $v = v_{i+1}$. Then $I_{G_k} \overline{G_k}[\overline{u}, \overline{v}]$ covers both the $\overline{E_k} \setminus \{u_{i+1}\}$ and $\overline{A_k} \setminus \{v_i, v_{i+2}\}$. This shows that $N \subseteq \{v_i, v_{i+1}, v_{i+2}, u_i, u_{i+1}\}$. Again, since N induces a complete multipartite graph, it follows that $N = \{v_i, v_{i+1}, u_i\}$ and so $k = 4$.

Subcase 3.2: both u and v are simplicial vertices.

In this situation either $u, v \in B_1$ or $u, v \in B_{k+1}$, say $u, v \in B_1$. Then $u = v_1$ and $v = u_1$. As in Subcase 3.1 we can then prove that $N \subseteq \{v_1, v_2, u_1\}$ and so $k \leq 4$.

Subcase 3.3: both u and v are cut vertices.

Then $u = v_i$ and $v = v_{i+1}$ for some $i \geq 2$. Similarly as in Subcase 3.1 we have that $N \subseteq \{v_{i-2}, v_{i-1}, v_i, v_{i+1}, u_i, u_{i+1}\}$. Again, since N induces a complete multipartite graph, it follows that $|N| \leq 3$ and so $k \leq 4$. Hence Claim B follows.

Since N is an independent set in G_k , we have that N contains at most one vertex from each block and so $|N| \leq k + 1$. Recall that N contains both cut vertices and simplicial vertices. This shows that $|N| \leq k$. Again, since N is an independent set, it follows that $M \cap N = \emptyset$.

Claim C: If $v_2 \in N$ or $v_{k+1} \in N$, then $|N| \leq \lceil \frac{k}{2} \rceil$ for all $k \geq 2$.

To prove this claim, we use induction on the number of cut vertices k . If $k = 2$ or 3 the result holds. Assume the result holds for all integers i with $3 \leq i < k$ and consider G_k . Suppose that $v_2 \in N$ or $v_{k+1} \in N$, say $v_{k+1} \in N$. Now, since N is independent in G_k , we have that both $u_{k+1} \notin N$ and $v_{k+2} \notin N$. Hence $N \subseteq V(G_{k-1})$ and so by induction hypothesis $|N| \leq \lceil \frac{k-1}{2} \rceil \leq \lceil \frac{k}{2} \rceil$. Hence Claim C follows.

In the following, we prove that $|N| \leq \lceil \frac{k}{2} \rceil$. Choose α with $2 \leq \alpha \leq k + 1$ such that the cut vertex $v_\alpha \in N$. Then it follows from Claim C that

$$|N \cap V(G_{\alpha-1})| \leq \left\lceil \frac{\alpha - 1}{2} \right\rceil.$$

and

$$|N \cap [(V(G_k) \setminus V(G_{\alpha-3})) \cup \{v_{\alpha-1}\}]| \leq \left\lceil \frac{k + 1 - \alpha}{2} \right\rceil.$$

Since v_α is counted twice, we have that

$$\begin{aligned} |N| &\leq |N \cap V(G_{\alpha-1})| + |N \cap [(V(G_k) \setminus V(G_{\alpha-3})) \cup \{v_{\alpha-1}\}]| \\ &\leq \left\lceil \frac{\alpha-1}{2} \right\rceil + \left\lceil \frac{k+1-\alpha}{2} \right\rceil - 1 \\ &\leq \left\lceil \frac{k}{2} \right\rceil. \end{aligned}$$

This shows that $|M \cup N| \leq k + \lceil \frac{k}{2} \rceil + 3$.

Since in all the cases we have arrived at a contradiction, we conclude that $\text{gp}(G_k \overline{G_k}) = n(G_k) - \lfloor \frac{k}{2} \rfloor$. \square

Acknowledgments

Sandi Klavžar acknowledges the financial support from the Slovenian Research Agency (research core funding P1-0297 and projects J1-9109, J1-1693, N1-0095). Neethu P K acknowledges the Council of Scientific and Industrial Research(CSIR), Govt. of India for providing financial assistance in the form of Junior Research Fellowship. Manoj Changat acknowledges the financial support from the Department of Science and Technology(research project under 'MATRICS' scheme No. MTR/2017/000238).

References

- [1] B. S. Anand, S. V. Ullas Chandran, M. Changat, S. Klavžar, E. J. Thomas, A characterization of general position sets in graphs, *Appl. Math. Comput.* 359 (2019) 84–89.
- [2] A. Bendali-Braham, N. Ikhlef-Eschouf, M. Blidia, Some results on the b-chromatic number in complementary prism graphs, *RAIRO Oper. Res.* 53 (2019) 1187–1195.
- [3] D. M. Cardoso, P. Carvalho, M. A. A. de Freitas, C. T. M. Vinagre, Spectra, signless Laplacian and Laplacian spectra of complementary prisms of graphs, *Linear Algebra Appl.* 544 (2018) 325–338.
- [4] D. Castonguay, E. M. M. Coelho, H. Coelho, J. R. A. Nascimento, A note on the convexity number of complementary prisms, *Discrete Math. Theor. Comput. Sci.* 21 (2019), Paper No. 4, 10 pp.

- [5] M. A. Duarte, L. Penso, D. Rautenbach, U. S. Souza, Complexity properties of complementary prisms, *J. Comb. Optim.* 33 (2017) 365–372.
- [6] H. E. Dudeney, *Amusements in Mathematics*, Nelson, Edinburgh, 1917.
- [7] V. Froese, I. Kanj, A. Nichterlein, R. Niedermeier, Finding points in general position, *Internat. J. Comput. Geom. Appl.* 27 (2017) 277–296.
- [8] M. Ghorbani, S. Klavžar, H. R. Maimani, M. Momeni, F. Rahimi-Mahid, G. Rus, The general position problem on Kneser graphs and on some graph operations, *Discuss. Math. Graph Theory* (2019) doi:10.7151/dmgt.2269.
- [9] T. W. Haynes, M. A. Henning, L. C. van der Merwe, Domination and total domination in complementary prisms, *J. Comb. Optim.* 18 (2009) 23–37.
- [10] T. W. Haynes, M. A. Henning, P. J. Slater, L. C. van der Merwe, The complementary product of two graphs, *Bull. Inst. Combin. Appl.* 51 (2007) 21–30.
- [11] S. Klavžar, I. G. Yero, The general position problem and strong resolving graphs, *Open Math.* 17 (2019) 1126–1135.
- [12] S. Klavžar, B. Patkós, G. Rus, I. G. Yero, On general position sets in Cartesian grids, *arXiv [math.CO]* (25 Jul 2019).
- [13] C. Y. Ku, K. B. Wong, On no-three-in-line problem on m -dimensional torus, *Graphs Combin.* 34 (2018) 355–364.
- [14] P. Manuel, S. Klavžar, A general position problem in graph theory, *Bull. Aust. Math. Soc.* 98 (2018) 177–187.
- [15] P. Manuel, S. Klavžar, The graph theory general position problem on some interconnection networks, *Fund. Inform.* 163 (2018) 339–350.
- [16] D. Meierling, F. Protti, D. Rautenbach, A. R. de Almeida, Cycles in complementary prisms, *Discrete Appl. Math.* 193 (2015) 180–186.
- [17] A. Misiak, Z. Stępień, A. Szymaszkiwicz, L. Szymaszkiwicz, M. Zwierzchowski, A note on the no-three-in-line problem on a torus, *Discrete Math.* 339 (2016) 217–221.
- [18] B. Patkós, On the general position problem on Kneser graphs, *arXiv:1903.08056v2 [math.CO]* (21 Jul 2019).
- [19] M. Payne, D. R. Wood, On the general position subset selection problem, *SIAM J. Discrete Math.* 27 (2013) 1727–1733.

- [20] M. Skotnica, No-three-in-line problem on a torus: periodicity, *Discrete Math.* 342 (2019) 111611, 13 pp.
- [21] S. V. Ullas Chandran, G. Jaya Parthasarathy, The geodesic irredundant sets in graphs, *Int. J. Math. Combin.* 4 (2016) 135–143.
- [22] L. M. Zatesko, R. Carmo, A. L. P. Guedes, A. Zorzi, R. C. S. Machado, C. M. H. Figueiredo, On the chromatic index of complementary prisms, *Acta Math. Univ. Comenian. (N.S.)* 88 (2019) 1071–1077.