

On the difference between the eccentric connectivity index and eccentric distance sum of graphs

Yaser Alizadeh ^a Sandi Klavžar ^b

^a Department of Mathematics, Hakim Sabzevari University, Sabzevar, Iran
e-mail: y.alizadeh.s@gmail.com

^b Faculty of Mathematics and Physics, University of Ljubljana, Slovenia
e-mail: sandi.klavzar@fmf.uni-lj.si

Abstract

The eccentric connectivity index of a graph G is $\xi^c(G) = \sum_{v \in V(G)} \varepsilon(v) \deg(v)$, and the eccentric distance sum is $\xi^d(G) = \sum_{v \in V(G)} \varepsilon(v) D(v)$, where $\varepsilon(v)$ is the eccentricity of v , and $D(v)$ the sum of distances between v and the other vertices. A lower and an upper bound on $\xi^d(G) - \xi^c(G)$ is given for an arbitrary graph G . Regular graphs with diameter at most 2 and joins of cocktail-party graphs with complete graphs form the graphs that attain the two equalities, respectively. Sharp lower and upper bounds on $\xi^d(T) - \xi^c(T)$ are given for arbitrary trees. Sharp lower and upper bounds on $\xi^d(G) + \xi^c(G)$ for arbitrary graphs G are also given, and a sharp lower bound on $\xi^d(G)$ for graphs G with a given radius is proved.

Key words: eccentricity; eccentric connectivity index; eccentric distance sum; tree

AMS Subj. Class (2020): 05C12, 05C09, 05C92

1 Introduction

In this paper we consider simple and connected graphs. If $G = (V(G), E(G))$ is a graph and $u, v \in V(G)$, then the *distance* $d_G(u, v)$ between u and v is the number of edges on a shortest u, v -path. The eccentricity of a vertex and its total distance are distance properties of central interest in (chemical) graph theory; they are defined as follows. The *eccentricity* $\varepsilon_G(v)$ of a vertex v is the distance between v and a farthest vertex from v , and the *total distance* $D_G(v)$ of v is the sum of distances between v and the other vertices of G . Even more fundamental property of a vertex in (chemical) graph theory is its degree (or valence in chemistry), denoted by $\deg_G(v)$. (We may skip the

index G in the above notations when G is clear.) Multiplicatively combining two out of these three basic invariants naturally leads to the *eccentric connectivity index* $\xi^c(G)$, the *eccentric distance sum* $\xi^d(G)$, and the *degree distance* $DD(G)$, defined as follows:

$$\begin{aligned}\xi^c(G) &= \sum_{v \in V(G)} \varepsilon(v) \deg(v). \\ \xi^d(G) &= \sum_{v \in V(G)} \varepsilon(v) D(v). \\ DD(G) &= \sum_{v \in V(G)} \deg(v) D(v).\end{aligned}$$

ξ^c was introduced by Sharma, Goswami, and Madan [17], ξ^d by Gupta, Singh, and Madan [7], and DD by Dobrynin and Kochetova [6] and by Gutman [8]. These three topological indices are well investigated, selected contributions to the eccentric connectivity index are [10, 13, 22], to the eccentric distance sum [1, 14, 21], and to the degree distance [15, 18, 19]. The three invariants were also compared to other invariants, cf. [2, 3, 4, 5, 23]. For information on additional topological indices based on eccentricity see [16].

In [11] the eccentric distance sum and the degree distance are compared, while in [24] the difference between the eccentric connectivity index and the (not defined here) connective eccentricity index is studied. The primary motivation for the present paper, however, are the papers [12, 25] in which $\xi^d(G) - \xi^c(G)$ was investigated. In [25], Zhang, Li, and Xu, besides other results on the two indices, determined sharp upper and lower bounds on $\xi^d(G) - \xi^c(G)$ for graphs G of given order and diameter 2. Parallel results were also derived for sub-classes of diameter 2 graphs with specified one of the minimum degree, the connectivity, the edge-connectivity, and the independence number. Hua, Wang, and Wang [12] extended the last result to general graphs. More precisely, they characterized the graphs that attain the minimum value of $\xi^d(G) - \xi^c(G)$ among all connected graphs G of given independence number. They also proved a related result for connected graphs with given matching number.

In this paper we continue the investigation along the lines of [12, 25] and proceed as follows. In the rest of this section definitions and some observations needed are listed. In Section 2, we give a lower and an upper bound on $\xi^d(G) - \xi^c(G)$ and in both cases characterize the equality case. The upper bound involves the Wiener index, the first Zagreb index, as well as the degree distance of G . In Section 3 we focus on trees and first prove that among all trees T with given order and diameter, $\xi^d(T) - \xi^c(T)$ is minimized on caterpillars. Using this result we give a lower bound on $\xi^d(T) - \xi^c(T)$ for all trees T with given order, the bound being sharp precisely on stars. We also give a sharp upper bound on $\xi^d(T) - \xi^c(T)$ for trees T with given order. In the last section we give a sharp lower bound and a sharp upper bound on $\xi^d(G) + \xi^c(G)$, compare $\xi^d(G)$ with $\xi^c(G)$ for graphs G with not too large maximum degree, and give a sharp lower bound on $\xi^d(G)$ for graphs G with a given radius.

1.1 Preliminaries

The order and the size of a graph G will be denoted by $n(G)$ and $m(G)$, respectively. The star of order $n \geq 2$ is denoted by S_n ; in other words, $S_n = K_{1,n-1}$. If $n \geq 2$, then the *cocktail party graph* CP_{2n} is the graph obtained from K_{2n} by removing a perfect matching. The *join* $G \oplus H$ of graphs G and H is the graph obtained from the disjoint union of G and H by connecting by an edge every vertex of G with every vertex of H . The maximum degree of a vertex of G is denoted by $\Delta(G)$. A graph G is *regular* if all vertices have the same degree. The *first Zagreb index* [9] $M_1(G)$ of G is the sum of the squares of the degrees of the vertices of G . The *Wiener index* [20] $W(G)$ of G is the sum of distances between all pairs of vertices in G .

The *diameter* $\text{diam}(G)$ and the *radius* $\text{rad}(G)$ of a graph G are the maximum and the minimum vertex eccentricity in G , respectively. A graph G is *self-centered* if all vertices have the same eccentricity. If this eccentricity is d , we further say that G is *d-self-centered*. The *eccentricity* $\varepsilon(G)$ of G is

$$\varepsilon(G) = \sum_{v \in V(G)} \varepsilon(v).$$

The eccentric connectivity index of G can be equivalently written as

$$\xi^c(G) = \sum_{uv \in E(G)} \varepsilon(u) + \varepsilon(v), \quad (1)$$

and the eccentric distance sum as

$$\xi^d(G) = \sum_{\{u,v\} \subseteq V(G)} (\varepsilon(u) + \varepsilon(v))d(u,v). \quad (2)$$

2 The difference on general graphs

In this section we give some sharp upper and lower bounds on $\xi^d(G) - \xi^c(G)$ for an arbitrary graph G . The bounds are in terms of the eccentricity, the Wiener index, the first Zagreb index, the degree distance, the maximum degree, the size, and the order of G .

Theorem 2.1 *If G is a connected graph, then the following hold.*

- (i) $\xi^d(G) - \xi^c(G) \geq 2(n(G) - 1 - \Delta(G))\varepsilon(G)$. Moreover, the equality holds if and only if G is a regular graph with $\text{diam}(G) \leq 2$.
- (ii) $\xi^d(G) - \xi^c(G) \leq 2n(G)(W(G) - m(G)) + M_1(G) - DD(G)$. Moreover, the equality holds if and only if $G \in \{P_4\} \cup \{CP_{2k} \oplus K_{n(G)-2k} : 0 \leq k \leq n/2\}$.

Proof. (i) Let v be a vertex of G . If w is not adjacent to v , then $d(v, w) \geq 2$ and consequently $D(v) - \deg(v) \geq 2(n(G) - 1 - \Delta(G))$. Thus:

$$\begin{aligned}\xi^d(G) - \xi^c(G) &= \sum_{v \in V(G)} \varepsilon(v)(D(v) - \deg(v)) \\ &\geq \sum_{v \in V(G)} 2\varepsilon(v)(n(G) - 1 - \Delta(G)) \\ &= 2\varepsilon(G)(n(G) - 1 - \Delta(G)).\end{aligned}$$

The equality holds if and only if $D(v) - \deg(v) = 2(n(G) - 1 - \Delta(G))$ for every vertex v . As the last equality in particular holds for a vertex of maximum degree, we infer that G must be regular. Then the condition $D(v) - \deg(v) = 2(n(G) - 1 - \Delta(G))$ simplifies to

$$D(v) + \Delta(G) = 2n(G) - 2. \quad (3)$$

Suppose that $\text{diam}(G) = d$, and let x_i , $i \in \{2, \dots, d\}$, be the number of vertices at distance i from v . Then $n(G) = 1 + \Delta(G) + x_2 + \dots + x_d$ and $D(v) = \Delta(G) + 2x_2 + \dots + dx_d$. Plugging these equalities into (3) yields

$$2\Delta(G) + 2x_2 + \dots + dx_d = 2 + 2\Delta(G) + 2x_2 + \dots + 2x_d - 2$$

which implies that $x_3 = \dots = x_d = 0$, that is, $\text{diam}(G) = 2$. Finally, if $\text{diam}(G) = 2$, then $D(v) = \Delta(G) + 2(n(G) - \Delta(G) - 1)$, so (3) is fulfilled for every regular graph of diameter 2. Clearly, (3) is also fulfilled for graphs of diameter 1, that is, complete graphs.

(ii) If $v \in V(G)$, then clearly $\varepsilon(v) \leq n(G) - \deg(v)$. Then we deduce that

$$\begin{aligned}\xi^d(G) - \xi^c(G) &= \sum_{v \in V(G)} \varepsilon(v)(D(v) - \deg(v)) \\ &\leq \sum_{v \in V(G)} (n(G) - \deg(v))(D(v) - \deg(v)) \\ &= n(G) \sum_{v \in V(G)} (D(v) - \deg(v)) + \sum_{v \in V(G)} \deg(v)^2 \\ &\quad - \sum_{v \in V(G)} \deg(v)D(v) \\ &= 2n(G)(W(G) - m(G)) + M_1(G) - DD(G).\end{aligned}$$

The equality in the above computation holds if and only if $\varepsilon(v) = n(G) - \deg(v)$ holds for all $v \in V(G)$. So suppose that G is a graph for which $\varepsilon(v) = n(G) - \deg(v)$ holds for all $v \in V(G)$ and distinguish the following two cases.

Suppose first that $\text{diam}(G) \geq 3$. Let P be a diametral path in G and let v and v' be its endpoints. Since $\varepsilon(v) = n(G) - \deg(v)$ and $|V(P) \setminus N[v]| = \varepsilon(v) - 1$, it follows

that $n(G) = 1 + \deg(v) + |V(P) \setminus N[v]|$. The latter means that $V(G) = N[v] \cup V(P)$. Since $\text{diam}(G) = \varepsilon(v) \geq 3$ it follows that $\deg(v') = 1$. Since we have also assumed that $\varepsilon(v') = n(G) - \deg(v')$ holds we see that $\varepsilon(v') = n(G) - 1$ which in turn implies that G is a path. Among the paths P_n , $n \geq 4$, the path P_4 is the unique one which fulfills the condition $\varepsilon(v) = n - \deg(v)$ for all $v \in V(P_n)$.

Suppose second that $\text{diam}(G) \leq 2$. Then $\varepsilon(v) \in \{1, 2\}$ for every $v \in (G)$. Since $\varepsilon(v) = n(G) - \deg(v)$ it follows that $\deg(v) \in \{n(G) - 1, n(G) - 2\}$. Let $V_1 = \{v : \deg(v) = n(G) - 1\}$ and $V_2 = \{v : \deg(v) = n(G) - 2\}$. Then $V(G) = V_1 \cup V_2$. Clearly, the subgraph of G induced by V_1 is complete, and there are all possible edges between V_1 and V_2 . Moreover, the complement of the subgraph of G induced by V_2 is a disjoint union of copies of K_2 , which means that V_2 induces a cocktail party graph. In summary, G must be of the form $CP_{2k} \oplus K_{n(G)-2k}$, where $0 \leq k \leq n/2$. On the other hand, the condition $\varepsilon(v) = n(G) - \deg(v)$ clearly holds for each vertex of $CP_{2k} \oplus K_{n(G)-2k}$, hence these graphs together with P_4 from the previous case are precisely the graphs that attain the equality. \square

3 The difference on trees

In this section we turn our attention to $\xi^d(T) - \xi^c(T)$ for trees T , and in particular on extremal trees regarding this difference.

Theorem 3.1 *Among all trees T with given order and diameter, $\min\{\xi^d(T) - \xi^c(T)\}$ is achieved on caterpillars.*

Proof. Fix the order and diameter of trees to be considered. Let T be an arbitrary tree that is not a caterpillar with this fixed order and diameter. Let P be a diametral path of T connecting x to y . Then the eccentricity of each vertex w of T is equal to $\max\{d(w, x), d(w, y)\}$. Let $z \neq x, y$ be a vertex of P and let T_z be a maximal subtree of T which contains z but no other vertex of P . We may assume that z can be selected such that $\varepsilon_{T_z}(z) = k \geq 2$, for otherwise T is a caterpillar. Let u be vertex of T_z with $d(u, z) = k - 1$ and let v be the neighbor of u with $d(v, z) = k - 2$. Let $S = N(u) \setminus \{v\}$ and let $s = |S|$. Note that $s > 0$. Let now T' be the tree obtained from T by replacing the edges between u and the vertices of S with the edges between v and the vertices of S .

Claim A: $\xi^d(T) - \xi^c(T) > \xi^d(T') - \xi^c(T')$.

Set $X_d = \xi^d(T) - \xi^d(T')$ and $X_c = \xi^c(T) - \xi^c(T')$. To prove the claim it is equivalent to show that $X_d - X_c > 0$.

For a vertex $w \in V(G) \setminus (S \cup \{u\})$ we have $D_{T'}(w) = D_T(w) - s$ and $\varepsilon_{T'}(w) \leq \varepsilon_T(w)$. Moreover if $w \in S$, then $\varepsilon_{T'}(w) = \varepsilon_T(w) - 1$ and $D_T(w) = D_{T'}(w) + n - s - 2$. With

these facts in hand we can compute as follows.

$$\begin{aligned}
X_d &= \sum_{w \in V(T)} \varepsilon_T(w) D_T(w) - \sum_{w \in V(T')} \varepsilon_{T'}(w) D_{T'}(w) \\
&= \varepsilon_T(u) D_T(u) - \varepsilon_{T'}(u) D_{T'}(u) + \varepsilon_T(v) D_T(v) - \varepsilon_{T'}(v) D_{T'}(v) \\
&\quad + \sum_{w \in S} \varepsilon_T(w) D_T(w) - \varepsilon_{T'}(w) D_{T'}(w) \\
&\quad + \sum_{w \in V(T) - (S \cup \{u, v\})} \varepsilon_T(w) D_T(w) - \varepsilon_{T'}(w) D_{T'}(w) \\
&\geq s(\varepsilon_T(v) - \varepsilon_T(u)) + \sum_{w \in V(T) - (S \cup \{u, v\})} \varepsilon_T(w) s \\
&\quad + \sum_{w \in S} (\varepsilon_T(w) D_T(w) - (\varepsilon_T(w) - 1)(D_T(w) - n + 2 + s)) \\
&= -s + \sum_{w \in V(T) - (S \cup \{u, v\})} \varepsilon_T(w) s \\
&\quad + \sum_{w \in S} ((D_T(w) - n + 2 + s) - \varepsilon_T(w)(-n + 2 + s)) \\
&= -s + \sum_{w \in V(T) \setminus (S \cup \{u, v\})} \varepsilon_T(w) s + (n - s - 2) \sum_{w \in S} \varepsilon_T(w) - 1 + D_T(w) \\
&= -s + \sum_{w \in V(T) \setminus (S \cup \{u, v\})} \varepsilon_T(w) s \\
&\quad + s(n - s - 2)\varepsilon_T(u) + s(D_T(u) + n - 2) \\
&= s[\varepsilon(T) - \varepsilon_T(u)(s + 2) - s + 1 + (n - s - 2)\varepsilon_T(u) + D_T(u) + n - 3].
\end{aligned}$$

Similarly, but shorter, we get that $X_c = 2s$. Thus

$$\begin{aligned}
X_d - X_c &\geq s[\varepsilon(T) - \varepsilon_T(u)(s + 2) \\
&\quad + (n - s - 2)\varepsilon_T(u) + D_T(u) + n - s - 4] \\
&> 0.
\end{aligned}$$

This proves Claim A. If T' is not a caterpillar, we can repeat the construction as many times as required to arrive at a caterpillar. Since at each step the value of $\xi^d - \xi^c$ is decreased, the minimum of this difference is indeed achieved on caterpillars. \square

Theorem 3.2 *If T is a tree of order $n \geq 3$, then*

$$\xi^d(T) - \xi^c(T) \geq 4n^2 - 12n + 8.$$

Moreover, equality holds if and only if $T = S_n$.

Proof. Let $n \geq 3$ be a fixed integer. By Theorem 3.1, it suffices to consider caterpillars. More precisely, let T be a caterpillar of order n and with $\text{diam}(T) = d \geq 3$. Then we wish to prove that $\xi^d(T) - \xi^c(T) > \xi^d(S_n) - \xi^c(S_n) = 4n^2 - 12n + 8$. The latter equality is straightforward to check, for the strict inequality we proceed as follows.

Let $w, z \in V(T)$ be two adjacent vertices of eccentricities $d-1$ and $d-2$, respectively. Let $S = N(w) \setminus \{z\}$ and set $s = |S|$. As $\varepsilon(w) = d-1$, we have $s \geq 1$. Let further $S_1 = V(G) \setminus (S \cup \{w, z\})$. Construct now a tree T' from T by replacing the edges between w and the vertices of S with the edges between z and the vertices of S . Note that $\deg_T(w) = \deg_{T'}(w) + s = 1 + s$ and $\deg_T(z) = \deg_{T'}(z) - s$, while the other vertices have the same degree in T and T' . Further, it is straightforward to verify the following relations:

$$\begin{aligned} D_T(w) &= D_{T'}(w) - s, & \varepsilon_T(w) &= \varepsilon_{T'}(w); \\ D_T(z) &= D_{T'}(z) + s, & \varepsilon_{T'}(z) &\leq \varepsilon_T(z) \leq \varepsilon_{T'}(z) + 1; \\ D_T(x) &= D_{T'}(x) + n - s - 2, & \varepsilon_T(x) &= \varepsilon_{T'}(x) + 1 \quad (x \in S); \\ D_T(y) &= D_{T'}(y) + s, & \varepsilon_{T'}(y) &\leq \varepsilon_T(y) \leq \varepsilon_{T'}(y) + 1 \quad (y \in S_1). \end{aligned}$$

Setting $X_d = \xi^d(T) - \xi^d(T')$ we have:

$$\begin{aligned} X_d &= \sum_{v \in \{w, z\}} D_T(v)\varepsilon_T(v) - D_{T'}(v)\varepsilon_{T'}(v) + \sum_{v \in S} D_T(v)\varepsilon_T(v) - D_{T'}(v)\varepsilon_{T'}(v) \\ &\quad + \sum_{v \in S_1} D_T(v)\varepsilon_T(v) - D_{T'}(v)\varepsilon_{T'}(v) \\ &\geq s(\varepsilon_{T'}(z) - \varepsilon_T(w)) + \sum_{v \in S} D_T(v)\varepsilon_T(v) - (D_T(v) - (n - s - 2))(\varepsilon_T(v) - 1) \\ &\quad + \sum_{v \in S_1} D_T(v)\varepsilon_T(v) - (D_T(v) - s)\varepsilon_T(v) \\ &\geq -s + (n - s - 2) \sum_{v \in S} \varepsilon_T(v) + \sum_{v \in S} D_T(v) - s(n - s - 2) + s \sum_{v \in S_1} \varepsilon_T(v) \\ &\geq -s + 3s(n - s - 2) + s(2(n - s - 2) + 2s + 1) - s(n - s - 2) \\ &\quad + 3s(n - s - 2) \\ &= 5s(n - s - 3) + 2s(n - 1). \end{aligned}$$

Similarly, setting $X_c = \xi^c(T) - \xi^c(T')$, we have

$$\begin{aligned}
X_c &= \sum_{v \in \{w, z\}} (\deg_T(v)\varepsilon_T(v) - \deg_{T'}(v)\varepsilon_{T'}(v)) \\
&\quad + \sum_{v \in S} (\deg_T(v)\varepsilon_T(v) - \deg_{T'}(v)\varepsilon_{T'}(v)) \\
&\quad + \sum_{v \in S_1} (\deg_T(v)\varepsilon_T(v) - \deg_{T'}(v)\varepsilon_{T'}(v)) \\
&\leq s\varepsilon_T(w) + \deg_T(z)\varepsilon_T(z) - (\deg_T(z) + s)(\varepsilon_T(z) - 1) \\
&\quad + s + \sum_{v \in S_1} \deg_T(v)\varepsilon_T(v) - \deg_T(v)(\varepsilon_T(v) - 1) \\
&= 2s + \deg_T(z) + \sum_{v \in S_1} \deg(v) \\
&= 2n - 3.
\end{aligned}$$

Therefore,

$$X_d - X_c \geq (5s(n - s - 3) + 2s(n - 1)) - (2n - 3) > 0,$$

that is, $\xi^d(T) - \xi^c(T) > \xi^d(T') - \xi^c(T')$. Repeating the above transformation until S_n is constructed finishes the argument. \square

To bound the difference $\xi^d(T) - \xi^c(T)$ for an arbitrary tree T from above, we first recall the following result.

Lemma 3.3 [14, Theorem 2.1] *Let w be a vertex of graph G . For non-negative integers p and q , let $G(p, q)$ denotes the graph obtained from G by attaching to vertex w pendant paths $P = wv_1 \cdots v_p$ and $Q = wu_1 \cdots u_q$ of lengths p and q , respectively. Let $G(p + q, 0) = G(p, q) - wu_1 + v_p u_1$. If $r = \varepsilon_G(w)$ and $r \geq p \geq q \geq 1$, then*

$$\begin{aligned}
\xi^d(G(p + q, 0)) - \xi^d(G(p, q)) &\geq \frac{pq}{6} [6D_G(w) + p(2p - 3) + q(2q - 3) + 3pq - 12r \\
&\quad + 6n(G)(p + q + r + 1) + 6 \sum_{v \in V(G)} \varepsilon(v)].
\end{aligned}$$

Lemma 3.4 *Let G , p , q , $G(p, q)$, and $G(p + q, 0)$ be as in Lemma 3.3. Then*

$$\xi^c(G(p + q, 0)) - \xi^c(G(p, q)) \leq q(3p + 2m(G) - 1).$$

Proof. Let $\deg'(v)$ and $\varepsilon'(v)$ (resp. $\deg(v)$ and $\varepsilon(v)$) denote the degree and the eccentricity of v in $G(p+q, 0)$ (resp. $G(p, q)$). Then we have:

$$\begin{aligned}\deg'(w) &= \deg(w) - 1, \quad \varepsilon'(w) \leq \varepsilon(w) + q; \\ \deg'(v_i) &= \deg(v_i), i \in [p-1], \quad \deg'(v_p) = \deg(v_p) + 1; \\ \varepsilon'(v_i) &\leq \varepsilon(v_i) + q, i \in [p]; \\ \deg'(u_j) &= \deg(u_j), \quad \varepsilon'(u_j) = \varepsilon(u_j) + p; \\ \varepsilon'(x) &\leq \varepsilon(x) + q, x \in V(G).\end{aligned}$$

Moreover, the degrees of vertices in $G(p+q, 0)$ do not decrease. Calculating the difference of contributions of vertices in ξ^c for $G(p+q, 0)$ and $G(p, q)$, we can estimate the difference $X_c = \xi^c(G(p+q, 0)) - \xi^c(G(p, q))$ as follows:

$$\begin{aligned}X_c &\leq \sum_{w \neq x \in V(G)} \deg(x)q + \sum_{i=1}^q \deg(u_i)p + \sum_{i=1}^{\lfloor \frac{p}{2} \rfloor} \deg(v_i)q \\ &\quad + \varepsilon(v_p) + (\deg(w) - 1)(r+q) - \deg(w)r \\ &= (2m(G) - \deg(w))q + (2q-1)p + pq + p + q(\deg(w) - 1) \\ &= 2qm(G) + 3pq - q.\end{aligned}$$

□

Theorem 3.5 *If T is a tree of order n , then*

$$\xi^d(T) - \xi^c(T) \leq \begin{cases} \frac{25n^4}{96} - \frac{n^3}{6} - \frac{89n^2}{48} + \frac{19n}{6} - \frac{45}{32}; & n \text{ odd}, \\ \frac{25n^4}{96} - \frac{n^3}{6} - \frac{43n^2}{24} + \frac{19n}{6} - 2; & n \text{ even}. \end{cases}$$

Moreover, equality holds if and only if $T = P_n$.

Proof. The right side of the above inequality is equal to $\xi^d(P_n) - \xi^c(P_n)$. (The value of $\xi^d(P_n)$ has been determined in [14], while it is straightforward to deduce $\xi^c(P_n)$. Combining the two formulas, the polynomials from the right hand side of the inequality are obtained.) Suppose now that $T \neq P_n$. Then there is always a vertex w of degree at least 3 such that we can apply Lemmas 3.3 and 3.4. Setting

$$X_{dc} = (\xi^d(T(p+q, 0)) - \xi^c(p+q, 0)) - (\xi^d(T(p, q)) - \xi^c(p, q))$$

we have:

$$\begin{aligned}
X_{dc} &\geq pqD_T(w) + \frac{pq}{6}(p(2p-3) + q(2q-3)) + \frac{(pq)^2}{2} - 2pqr \\
&\quad + pqn(T)(p+q+r+1) + pq \sum_{v \in V(T)} \varepsilon(v) - (2qm(T) + 3pq - q) \\
&= pq(D_T(w) + \sum_{v \in V(T)} \varepsilon(v) - 3) + \frac{pq}{6}(2p^2 - 3p + 2q^2 - 3q + 3pq) \\
&\quad + pqr(n(G) - 2) + q(pn(T)(p+q+r) - 2m(T) + 1) > 0
\end{aligned}$$

and the result follows. \square

4 Further comparison

In this concluding section we give sharp lower and upper bounds on $\xi^d(G) + \xi^c(G)$, compare $\xi^d(G)$ with $\xi^c(G)$ for graphs G with $\Delta(G) \leq \frac{2}{3}(n-1)$, and give a sharp lower bound on $\xi^d(G)$ for graphs G with a given radius.

Theorem 4.1 *If G is a connected graph, then the following hold.*

- (i) $\xi^d(G) + \xi^c(G) \leq 2(n(G) - 1)\varepsilon(G) + 2\text{diam}(G)(W(G) + m(G) - 2\binom{n(G)}{2})$.
- (ii) $\xi^d(G) + \xi^c(G) \geq 2(n(G) - 1)\varepsilon(G) + 2\text{rad}(G)(W(G) + m(G) - 2\binom{n(G)}{2})$.

Moreover, each of the equalities holds if and only if G is a self-centered graph.

Proof. (i) Partition the pairs of vertices of G into neighbors and non-neighbors, and using (1), we can compute as follows:

$$\begin{aligned}
\xi^d(G) &= \sum_{\{u,v\} \subseteq V(G)} (\varepsilon(u) + \varepsilon(v))d(u,v) \\
&= \sum_{uv \in E(G)} (\varepsilon(u) + \varepsilon(v)) + 2 \sum_{\substack{\{u,v\} \subseteq V(G) \\ d(u,v) \geq 2}} (\varepsilon(u) + \varepsilon(v)) \\
&\quad + \sum_{\substack{\{u,v\} \subseteq V(G) \\ d(u,v) \geq 2}} (\varepsilon(u) + \varepsilon(v))(d(u,v) - 2) \\
&= \xi^c(G) + \sum_{\{u,v\} \subseteq V(G)} (\varepsilon(u) + \varepsilon(v)) - 2\xi^c(G) \\
&\quad + \sum_{\substack{\{u,v\} \subseteq V(G) \\ d(u,v) \geq 2}} (\varepsilon(u) + \varepsilon(v))(d(u,v) - 2) \\
&\leq -\xi^c(G) + 2(n(G) - 1)\varepsilon(G) \\
&\quad + 2\text{diam}(G)(W(G) + m(G) - 2\binom{n(G)}{2}).
\end{aligned}$$

The inequality above becomes equality if and only if $\varepsilon(v) = \text{diam}(G)$ for every $v \in V(G)$. That is, the equality holds if and only if G is a self-centered graph.

(ii) This inequality as well as its equality case are proved along the same lines as (i). The only difference is that the inequality $\varepsilon(u) + \varepsilon(v) \leq 2\text{diam}(G)$ is replaced by $\varepsilon(u) + \varepsilon(v) \geq 2\text{rad}(G)$. \square

In our next result we give a relation between $\xi^d(G)$ and $\xi^c(G)$ for graph G with maximum degree at most $\frac{2}{3}(n(G) - 1)$.

Theorem 4.2 *If G is a graph with $\Delta(G) \leq \frac{2}{3}(n - 1)$, then $\xi^d(G) \geq 2\xi^c(G)$. Moreover, the equality holds if and only if G is 2-self-centered, $\frac{2}{3}(n(G) - 1)$ -regular graph.*

Proof. Set $n = n(G)$ and let v be a vertex of G . Since $\deg(v) < n - 1$ we have $\varepsilon(v) \geq 2$. Therefore $D(v) \geq 2(n - 1) - \deg(v)$ with equality holding if and only if $\varepsilon(v) = 2$. Using the assumption that $\deg(v) \leq \frac{2}{3}(n - 1)$, equivalently, $2n - 2 \geq 3\deg(v)$, we infer that $\varepsilon(v)D(v) \geq 2\varepsilon(v)\deg(v)$. Summing over all vertices of G the inequality is proved. Its derivation also reveals that the equality holds if and only if $\deg(v) = \frac{2}{3}(n - 1)$ and $\varepsilon(v) = 2$ for each vertex $v \in V(G)$. \square

To conclude the paper we give a lower bound on the eccentric distance sum in terms of the radius of a given graph. Interestingly, the cocktail-party graphs are again among the extreme graphs.

Theorem 4.3 *If G is a graph with $\text{rad}(G) = r$, then*

$$\xi^d(G) \geq (n(G) - 1 + \binom{r}{2})\varepsilon(G).$$

Equality holds if and only if G is a complete graph or a cocktail-party graph.

Proof. Set $n = n(G)$ and let $v \in V(G)$. Let P be a longest path starting in v . Separately considering the neighbors of v , the last $\varepsilon(v) - 2$ vertices of P , and all the other vertices, we can estimate that

$$\begin{aligned} D(v) &\geq \deg(v) + (3 + \cdots + \varepsilon(v)) + 2(n - 1 - \deg(v) - (\varepsilon(v) - 2)) \\ &= 2n - \deg(v) + \frac{\varepsilon(v)^2 - 3\varepsilon(v)}{2} - 1. \end{aligned}$$

Since $n - \deg(v) \geq \varepsilon(v)$ for every vertex $v \in V(G)$, we have $D(v) \geq n + \varepsilon(v) + \frac{\varepsilon(v)^2 - 3\varepsilon(v)}{2} - 1$. Consequently, having the fact $\varepsilon(v) \geq r$ in mind, we get $D(v) \geq n - 1 + \binom{r}{2}$. Multiplying this inequality by $\varepsilon(v)$ and summing over all vertices of G the claimed inequality is proved.

From the above derivation we see that the equality can hold only if $\varepsilon(v) = r = n - \deg(v)$ holds for every $v \in V(G)$. From the equality part of the proof of Theorem 2.1(ii) we know that this implies $\text{diam}(G) \leq 2$. For the equality we must also have $D(v) = n - 1 + \binom{r}{2}$ for every v . If $r = 2$ this means that $D(v) = n$ and hence $\deg(v) = n - 2$. It follows that G is a cocktail-party graph. And if $r = 2$, then we get a complete graph. \square

Acknowledgements

Sandi Klavžar acknowledges the financial support from the Slovenian Research Agency (research core funding P1-0297 and projects J1-9109, J1-1693, N1-0095).

References

- [1] S. Chen, S. Li, Y. Wu, L. Sun, Connectivity, diameter, minimal degree, independence number and the eccentric distance sum of graphs, *Discrete Appl. Math.* **247** (2018) 135–146.
- [2] P. Dankelmann, M.J. Morgan, S. Mukwembi, H.C. Swart, On the eccentric connectivity index and Wiener index of a graph, *Quaest. Math.* **37** (2014) 39–47.
- [3] K.Ch. Das, N. Trinajstić, Relationship between the eccentric connectivity index and Zagreb indices, *Comput. Math. Appl.* **62** (2011) 1758–1764.
- [4] K. Ch. Das, M.J. Nadjafi-Arani, Comparison between the Szeged index and the eccentric connectivity index, *Discrete Appl. Math.* **186** (2015) 74–86.
- [5] K.Ch. Das, G. Su, L. Xiong, Relation between degree distance and Gutman index of graphs, *MATCH Commun. Math. Comput. Chem.* **76** (2016) 221–232.
- [6] A.A. Dobrynin, A.A. Kochetova, Degree distance of a graph: a degree analogue of the Wiener index, *J. Chem. Inf. Comput. Sci.* **34** (1994) 1082–1086.
- [7] S. Gupta, M. Singh, A.K. Madan, Eccentric distance sum: A novel graph invariant for predicting biological and physical properties, *J. Math. Anal. Appl.* **275** (2002) 386–401.
- [8] I. Gutman, Selected properties of the Schultz molecular topological index, *J. Chem. Inf. Comput. Sci.* **34** (1994) 1087–1089.
- [9] I. Gutman, N. Trinajstić, Graph theory and molecular orbitals. Total π -electron energy of alternant hydrocarbons, *Chem. Phys. Lett.* **17** (1972) 535–538.
- [10] P. Hauweele, A. Hertz, H. Mélot, B. Ries, G. Devillez, Maximum eccentric connectivity index for graphs with given diameter, *Discrete Appl. Math.* **268** (2019) 102–111.
- [11] H. Hua, H. Wang, X. Hu, On eccentric distance sum and degree distance of graphs, *Discrete Appl. Math.* **250** (2018) 262–275.
- [12] H. Hua, H. Wang, M. Wang, The difference between the eccentric distance sum and eccentric connectivity index, *Ars Combin.* **144** (2019) 3–12.

- [13] A. Ilić, I. Gutman, Eccentric connectivity index of chemical trees, *MATCH Commun. Math. Comput. Chem.* **65** (2011) 731–744.
- [14] A. Ilić, G. Yu, L. Feng, On the eccentric distance sum of graph, *J. Math. Anal. Appl.* **381** (2011) 590–600.
- [15] S. Li, Y. Song, H. Zhang, On the degree distance of unicyclic graphs with given matching number, *Graphs Combin.* **31** (2015) 2261–2274.
- [16] A.K. Madan, H. Dureja, Eccentricity based descriptors for QSAR/QSPR. In: Novel Molecular Structure Descriptors - Theory and Applications II, I. Gutman, B. Furtula (Eds.), *Univ. Kragujevac, Kragujevac* (2010) 91–138.
- [17] V. Sharma, R. Goswami, A.K. Madan, Eccentric connectivity index: A novel highly discriminating topological descriptor for structure - property and structure - activity studies, *J. Chem. Inf. Comput. Sci.* **37** (1997) 273–282.
- [18] A.I. Tomescu, Unicyclic and bicyclic graphs having minimum degree distance, *Discrete Appl. Math.* **156** (2008) 125–130.
- [19] H. Wang, L. Kang, Further properties on the degree distance of graphs, *J. Comb. Optim.* **31** (2016) 427–446.
- [20] H. Wiener, Structural determination of paraffin boiling points, *J. Amer. Chem. Soc.* **69** (1947) 17–20.
- [21] Y.-T. Xie, S.-J. Xu, On the maximum value of the eccentric distance sums of cubic transitive graphs, *Appl. Math. Comput.* **359** (2019) 194–201.
- [22] K. Xu, K.C. Das, H. Liu, Some extremal results on the connective eccentricity index of graphs, *J. Math. Anal. Appl.* **433** (2016) 803–817.
- [23] K. Xu, X. Li, Comparison between two eccentricity-based topological indices of graphs, *Croat. Chem. Acta* **89** (2016) 499–504.
- [24] K. Xu, Y. Alizadeh, K.Ch. Das, On two eccentricity-based topological indices of graphs, *Discrete Appl. Math.* **233** (2017) 240–251.
- [25] H. Zhang, S. Li, B. Xu, Extremal graphs of given parameters with respect to the eccentricity distance sum and the eccentric connectivity index, *Discrete Appl. Math.* **254** (2019) 204–221.