

Maker-Breaker resolving game

Cong X. Kang¹ Sandi Klavžar^{2,3,4} Ismael G. Yero⁵
 Eunjeong Yi¹

Texas A&M University at Galveston, Galveston, TX 77553, USA¹
 Faculty of Mathematics and Physics, University of Ljubljana, Slovenia²
 Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia³
 Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia⁴
 Universidad de Cádiz, Av. Ramón Puyol s/n, 11202 Algeciras, Spain⁵
 kangc@tamug.edu; sandi.klavzar@fmf.uni-lj.si
 ismael.gonzalez@uca.es; yie@tamug.edu

May 28, 2020

Abstract

A set of vertices W of a graph G is a resolving set if every vertex of G is uniquely determined by its vector of distances to W . In this paper, the Maker-Breaker resolving game is introduced. The game is played on a graph G by Resolver and Spoiler who alternately select a vertex of G not yet chosen. Resolver wins if at some point the vertices chosen by him form a resolving set of G , whereas Spoiler wins if the Resolver cannot form a resolving set of G . The outcome of the game is denoted by $o(G)$ and $R_{\text{MB}}(G)$ (resp. $S_{\text{MB}}(G)$) denotes the minimum number of moves of Resolver (resp. Spoiler) to win when Resolver has the first move. The corresponding invariants for the game when Spoiler has the first move are denoted by $R'_{\text{MB}}(G)$ and $S'_{\text{MB}}(G)$. Invariants $R_{\text{MB}}(G)$, $R'_{\text{MB}}(G)$, $S_{\text{MB}}(G)$, and $S'_{\text{MB}}(G)$ are compared among themselves and with the metric dimension $\dim(G)$. A large class of graphs G is constructed for which $R_{\text{MB}}(G) > \dim(G)$ holds. The effect of twin equivalence classes and pairing resolving sets on the Maker-Breaker resolving game is described. As an application $o(G)$, as well as $R_{\text{MB}}(G)$ and $R'_{\text{MB}}(G)$ (or $S_{\text{MB}}(G)$ and $S'_{\text{MB}}(G)$), are determined for several graph classes, including trees, complete multi-partite graphs, grid graphs, and torus grid graphs.

Keywords: resolving set; metric dimension; Maker-Breaker game; Maker-Breaker resolving game; twin equivalence class, pairing resolving set

2020 Mathematics Subject Classification: 05C12, 05C57, 05C69

1 Introduction

Let $G = (V(G), E(G))$ be a finite, simple, undirected, connected graph of order at least 2. A set $W \subseteq V(G)$ is a *resolving set* of G if, for every pair of distinct vertices x and y of G , there exists $z \in W$ such that $d(x, z) \neq d(y, z)$, where $d(u, v)$ denotes the shortest-path distance between u and v . The *metric dimension* $\dim(G)$ of G is the minimum of the cardinalities over all resolving sets of G . A resolving set of cardinality $\dim(G)$ is called a *metric basis* for G . These concepts were independently introduced by Slater [27] and by Harary and Melter [16]. Soon after it was noted in [11] that determining the metric dimension of a graph is an NP-hard problem. Metric dimension has found applications in fields as diverse as robot navigation, network discovery and verification,

chemistry, combinatorial optimization, and strategies for the mastermind game. See [2, 7] for history and surveys and [1, 22, 28] for some of the more recent results on metric dimension.

The Maker-Breaker game, introduced in 1973 by Erdős and Selfridge [9], is played on an arbitrary hypergraph $H = (V, E)$. Two players, named Maker and Breaker, alternately select a vertex from V not yet chosen in the course of the game. Maker wins the game if he is able to select all the vertices of one of the hyperedges from E , while Breaker wins if she is able to prevent Maker from doing so. We refer to the books of Beck [3] and of Hefetz et al. [17] for more information on this game as well as to papers [15, 24] for recent related developments.

Motivated by the Maker-Breaker game and the domination game [4], Duchêne, Gledel, Parreau, and Renault introduced the Maker-Breaker domination game [8]. This game is played on a graph G and can be described as the Maker-Breaker game on the hypergraph with the same vertex set as G and with hyperedges corresponding to the dominating sets of G . The game was further investigated in [13], while in [12] its total version was introduced. Inspired by these developments, we introduce in this paper the *Maker-Breaker resolving game* (MBRG for short) as follows.

The MBRG is played on a graph G by two players, Resolver and Spoiler, which will be denoted throughout the paper by R^* and S^* , respectively. R^* and S^* alternately select (without missing their turn) a vertex of G that was not yet chosen in the course of the game. If R^* is the first to play, we speak of an *R-game*, otherwise we have an *S-game*. R^* wins if at some point the vertices R^* has chosen form a resolving set of G , whereas S^* wins if R^* cannot form a resolving set of G . The *outcome* of the MBRG on a graph G is denoted by $o(G)$, and there are four possible outcomes as follows: (1) $o(G) = \mathcal{R}$, if R^* has a winning strategy in the R-game and the S-game; (2) $o(G) = \mathcal{S}$, if S^* has a winning strategy in the R-game and the S-game; (3) $o(G) = \mathcal{N}$, if the first player has a winning strategy; (4) $o(G) = \tilde{\mathcal{N}}$, if the second player has a winning strategy.

Now, suppose a company X tries to secure its network by installing transmitters at certain locations within the company, so that the robot is aware of its security status at all times and thus identifying the exact location (or a specific computer with virus infection) in the network, whereas a rival company Y tries to prevent X from forming a secure network by occupying or controlling strategic locations or computers within the network of X . With this application in mind and considering the time constraint (the longer it takes for a player to win a game, the more it costs for the player), we introduce the following terminology and notation.

- The *Maker-Breaker resolving number* $R_{\text{MB}}(G)$ of G is the minimum number of moves of R^* to win the R-game provided he has a winning strategy. Otherwise, we set $R_{\text{MB}}(G) = \infty$.
- $R'_{\text{MB}}(G)$ is the minimum number of moves of R^* to win the S-game provided he has a winning strategy. Otherwise, we set $R'_{\text{MB}}(G) = \infty$.
- The *Maker-Breaker spoiling number* $S_{\text{MB}}(G)$ of G is the minimum number of moves of S^* to win the R-game provided she has a winning strategy. Otherwise, we set $S_{\text{MB}}(G) = \infty$.
- $S'_{\text{MB}}(G)$ is the minimum number of moves of S^* to win the S-game provided she has a winning strategy. Otherwise, we set $S'_{\text{MB}}(G) = \infty$.

This paper is organized as follows. In the next section we obtain some general results on the outcome of the MBRG. In Section 3 the effect of twin equivalence classes and pairing resolving sets on the MBRG is described and as an application a large class of graphs G is constructed for which $R_{\text{MB}}(G) > \dim(G)$ holds. In Section 4 we determine $o(G)$, as well as $R_{\text{MB}}(G)$ and $R'_{\text{MB}}(G)$ or $S_{\text{MB}}(G)$ and $S'_{\text{MB}}(G)$, when G is a tree, the Petersen graph, a bouquet of cycles, a complete multi-partite graph, a grid graph, or a torus grid graph.

2 Some general properties of the MBRG

In this section we compare parameters of the MBRG with the metric dimension, $R_{\text{MB}}(G)$ with $R'_{\text{MB}}(G)$, and $S_{\text{MB}}(G)$ with $S'_{\text{MB}}(G)$. Along the way we prove the so-called No-Skip Lemma for the

Maker-Breaker game played on a hypergraph. But first we comment on the possible outcomes of the MBRG.

Among the four possible outcomes listed in the introduction, the outcome $o(G) = \tilde{\mathcal{N}}$ never occurs as follows from a general result on the Maker-Breaker game, cf. [3, 17]. In the case of the Maker-Breaker domination game, this statement and its proof are given in [8, Proposition 2]. The same argument applies also to the Maker-Breaker resolving game. The other three possible outcomes in the latter game are realized, as the reader can verify on the examples given in Fig. 1.




Figure 1: Three examples realizing the three outcomes.

For further examples, note that if $n \geq 2$, then $o(P_n) = \mathcal{R}$ and $R_{\text{MB}}(P_n) = R'_{\text{MB}}(P_n) = \dim(P_n) = 1$. Moreover, $o(C_3) = \mathcal{N}$, and if $n \geq 4$, then $o(C_n) = \mathcal{R}$ and $R_{\text{MB}}(C_n) = R'_{\text{MB}}(C_n) = \dim(C_n) = 2$.

The order of a graph G will be denoted by $n(G)$. We have the following simple relations between the outcome of the MBRG and the metric dimension.

Proposition 2.1. *If G is a connected graph, then the following properties hold.*

- (i) *If $o(G) = \mathcal{R}$, then $\dim(G) \leq \lfloor \frac{n(G)}{2} \rfloor$.*
- (ii) *If $\dim(G) \geq \lceil \frac{n(G)}{2} \rceil + 1$, then $o(G) = \mathcal{S}$.*

Proof. (i) Suppose that $o(G) = \mathcal{R}$ and consider the S-game. After the game is finished, R^* has clearly selected at most $\lfloor \frac{n(G)}{2} \rfloor$ vertices. As the set of vertices selected by R^* forms a resolving set of G , we conclude that $\dim(G) \leq \lfloor \frac{n(G)}{2} \rfloor$.

(ii) No matter whether the R-game or the S-game is played, R^* selects at most $\lceil \frac{n(G)}{2} \rceil$ vertices by the end of the game. As $\dim(G) \geq \lceil \frac{n(G)}{2} \rceil + 1$, these vertices do not form a resolving set of G , hence S^* wins the R-game as well as the S-game. \square

If $n \geq 4$, then $\dim(K_n) = n - 1 \geq \lceil \frac{n}{2} \rceil + 1$, thus $o(K_n) = \mathcal{S}$ by Proposition 2.1(ii). Similarly, let B_m be the graph obtained from $m \geq 5$ disjoint copies of C_4 by identifying a vertex from each C_4 at a common vertex. Then $n(B_m) = 3m + 1$ and $\dim(B_m) = 2m - 1 \geq \lceil \frac{n(B_m)}{2} \rceil + 1$; thus $o(B_m) = \mathcal{S}$ by Proposition 2.1(ii).

Next, we compare $R_{\text{MB}}(G)$ with $R'_{\text{MB}}(G)$ and $S_{\text{MB}}(G)$ with $S'_{\text{MB}}(G)$. To this end, we consider the possibility that a player is allowed to skip a move; equivalently, a player allows the other player to select two vertices in one move. The observation that skipping offers no advantage to a player in the Maker-Breaker domination game was proved in [13]. We next show that a parallel argument works for the Maker-Breaker game played on an arbitrary hypergraph.

Lemma 2.2. (No-Skip Lemma) *If the Maker-Breaker game is played on a hypergraph H , then in an optimal strategy of R^* to win in the minimum number of moves it is never an advantage for him to skip a move. Moreover, it never disadvantages R^* for S^* to skip a move.*

Proof. Suppose the R-game or the S-game is played. Let R^* and S^* play optimally until S^* decides to skip a move. Then R^* imagines that S^* played an arbitrary legal move w , and replies optimally. R^* continues to use this strategy until the end of the game. It may happen that in the course of the

game S^* selects a vertex which was already selected in the imagined game of R^* . In that case, R^* imagines that some other legal move has been played by S^* . In this way, the game on G will finish in no more than the minimum number of moves played in the usual Maker-Breaker game. With a strategy of S^* parallel to the above strategy of R^* , it also follows that it is never an advantage for R^* to skip a move. \square

No-Skip Lemma quickly implies the announced comparison of $R_{\text{MB}}(G)$ with $R'_{\text{MB}}(G)$ and $S_{\text{MB}}(G)$ with $S'_{\text{MB}}(G)$.

Proposition 2.3. *If G is a connected graph, then the following properties hold.*

- (i) *If $o(G) = \mathcal{R}$, then $R'_{\text{MB}}(G) \geq R_{\text{MB}}(G) \geq \dim(G)$.*
- (ii) *If $o(G) = \mathcal{S}$, then $S_{\text{MB}}(G) \geq S'_{\text{MB}}(G)$.*

Proof. (i) The R-game can be viewed as the S-game in which S^* has skipped her first move. Hence the first inequality follows from Lemma 2.2 specialized to the MBRG. The second inequality follows from the fact that when the MBRG is finished, the set of vertices selected by R^* forms a resolving set of G .

(ii) The S-game can be viewed as the R-game in which R^* has skipped his first move, and thus the inequality follows. \square

For the lexicographic product graph $G = C_m[K_2]$, where $m \geq 4$, we have $\dim(G) = R_{\text{MB}}(G) = R'_{\text{MB}}(G) = m = \lfloor \frac{n(G)}{2} \rfloor$ (see [14] for the definition of the lexicographic product). This shows the sharpness of the bounds of Proposition 2.3(i). On the other hand, at the end of Section 3 we will construct a large family of graphs G for which $R_{\text{MB}}(G) > \dim(G)$ holds.

3 Twin equivalence classes and pairing resolving sets

In this section we consider two concepts that are very useful when dealing with the MBRG; this fact will be demonstrated in the rest of the paper.

The *open neighborhood* of a vertex $v \in V(G)$ is $N(v) = \{u \in V(G) \mid uv \in E(G)\}$. Vertices u and v are *twins* if $N(u) \setminus \{v\} = N(v) \setminus \{u\}$; notice that a vertex is its own twin. Hernando et al. [18, Lemma 2.7] observed that the twin relation is an equivalence relation and that an equivalence class under it, hereafter called a *twin equivalence class*, induces either a clique or an independent set. We recall the following well-known fact.

Observation 3.1. [18, Corollary 2.4] *If W is a resolving set of G and u and v are distinct members of the same twin equivalence class of G , then $W \cap \{u, v\} \neq \emptyset$.*

Here is now a relation of twin equivalence classes with the MBRG.

Proposition 3.2. *Let G be a connected graph with $n(G) \geq 4$.*

- (a) *If G has a twin equivalence class of cardinality at least 4, then $o(G) = \mathcal{S}$ and $S_{\text{MB}}(G) = S'_{\text{MB}}(G) = 2$.*
- (b) *If G has two distinct twin equivalence classes of cardinality at least 3, then $o(G) = \mathcal{S}$ and $S_{\text{MB}}(G) = S'_{\text{MB}}(G) = 2$.*

Proof. Let W be a resolving set of G .

(a) Let $Q = \{u_1, \dots, u_k\} \subseteq V(G)$ be a twin equivalence class of G , where $k \geq 4$. Then $|W \cap Q| \geq k - 1$ by Observation 3.1. Since $k \geq 4$, we infer that S^* can occupy two vertices of Q after her second move, regardless of whether S^* plays first or second. So, R^* can occupy at most $k - 2$ vertices of Q , and thus R^* fails to occupy vertices that form a resolving set of G . Thus $o(G) = \mathcal{S}$ and $S_{\text{MB}}(G) = S'_{\text{MB}}(G) = 2$.

(b) Let Q'_1 and Q'_2 be different twin equivalence classes of G , each of cardinality at least 3. Clearly, $Q'_1 \cap Q'_2 = \emptyset$. Let $Q_1 = \{u_1, u_2, u_3\} \subseteq Q'_1$ and $Q_2 = \{u'_1, u'_2, u'_3\} \subseteq Q'_2$. Then $|W \cap Q_1| \geq 2$ and $|W \cap Q_2| \geq 2$ by Observation 3.1. Note that S^* can occupy three vertices of $Q_1 \cup Q_2$ after her third move, regardless of whether S^* plays first or second. So, after the third move by S^* , there are the following four possibilities: (i) S^* occupies all vertices of Q_1 ; (ii) S^* occupies two vertices of Q_1 and one vertex of Q_2 ; (iii) S^* occupies one vertex of Q_1 and two vertices of Q_2 ; (iv) S^* occupies all three vertices of Q_2 . In each case, R^* fails to occupy vertices that form a resolving set of G . Thus $o(G) = \mathcal{S}$.

Next, we determine $S_{\text{MB}}(G)$ and $S'_{\text{MB}}(G)$. If S^* plays first, then after her second move, S^* can occupy two vertices of Q_1 . If S^* plays second, then after her second move, S^* can occupy two vertices of Q_1 (if R^* occupies a vertex in Q_2 in his first move) or S^* can occupy two vertices of Q_2 (if R^* occupies a vertex in Q_1 in his first move). So, $S_{\text{MB}}(G) = S'_{\text{MB}}(G) = 2$. \square

Let $[k]$ denote the set $\{1, \dots, k\}$. Let $A = \{\{u_1, w_1\}, \dots, \{u_k, w_k\}\}$ be a set of 2-subsets of $V(G)$ such that $|\cup_{i=1}^k \{u_i, w_i\}| = 2k$. We say that A is a *pairing resolving set* of G if every set $\{x_1, \dots, x_k\}$, where $x_i \in \{u_i, w_i\}$ and $i \in [k]$, is a resolving set of G .

Proposition 3.3. *If a graph G admits a pairing resolving set, then $o(G) = \mathcal{R}$.*

Proof. Let A be a pairing resolving set of G with $|A| = k$. Regardless of whether the R-game or the S-game is played, R^* is guaranteed to select a vertex of each pair from A after his k^{th} move. Thus, the vertices chosen by R^* , after his k^{th} move, form a resolving set of G . So, $o(G) = \mathcal{R}$. \square

Despite its simplicity, Proposition 3.3 has fine applications. (We note in passing that for the Maker-Breaker domination game a parallel concept of *pairing dominating sets* was introduced in [13].) If $\dim(G) = k$ and A is a pairing resolving set of G with $|A| = k$, then we say that A is a *dim-pairing resolving set* of G ; this, together with Proposition 3.3, immediately yields the following

Corollary 3.4. *If G admits a dim-pairing resolving set, then $R_{\text{MB}}(G) = R'_{\text{MB}}(G) = \dim(G)$.*

For an example, consider the graph G of Fig. 2. The graph G has the following dim-pairing resolving sets:

- $\{\{u_1, w_1\}, \{u_2, w_2\}, \{u_3, w_3\}, \{u_4, v_4\}\}$,
- $\{\{u_1, w_1\}, \{u_2, w_2\}, \{u_3, w_3\}, \{u_4, w_4\}\}$, and
- $\{\{u_1, w_1\}, \{u_2, w_2\}, \{u_3, w_3\}, \{v_4, w_4\}\}$,

hence Corollary 3.4 implies that $R_{\text{MB}}(G) = R'_{\text{MB}}(G) = 4$.




Figure 2: A graph which admits three dim-pairing resolving sets.

To conclude the section we are going to show how pairing resolving sets can be applied to construct a large family of graphs G for which $R_{\text{MB}}(G) > \dim(G)$ holds.

Definition 3.5. If $k \geq 3$, then let G_k be a graph of order $k + 2(2^k - 1)$ with $V(G_k) = A \cup B \cup C$, where A , B , and C are pairwise disjoint sets with $|A| = k$ and $|B| = 2^k - 1 = |C|$. The edge set of G_k is specified as follows: (i) each of the sets A , B , and C induces a clique in G_k ; (ii) indexing

the elements of B (and separately of C) by nonempty subsets of A , let $b_S \in B$ be adjacent to each vertex in $S \subseteq A$; (iii) let $b_S \in B$ be adjacent to $c_S \in C$ for each nonempty subset S of A ; (iv) there are no other edges.

We will at times subscript a vertex $b \in B$ (and a vertex $c \in C$) by an element of \mathbb{Z}_2^k , where 1 (resp., 0) in the j -th coordinate indicates that b is adjacent (resp., not adjacent) to the j -th vertex in A . See Fig. 3 for G_3 and the labeling of its vertices. From now on, for a given vertex $x \in V(G)$ and an ordered set of vertices $S \subseteq V(G)$, by $\text{code}_S(x)$ we denote the vector of distances between x and all the vertices in S .




Figure 3: G_3 satisfying $o(G_3) = \mathcal{R}$ and $R_{\text{MB}}(G_3) > \dim(G_3) = 3$.

Theorem 3.6. *If $k \geq 3$ and G_k is as in Definition 3.5, then the following holds.*

- (i) $\dim(G_k) = k$.
- (ii) *The set A is the unique metric basis of G_k .*
- (iii) *The set $\bigcup_{\emptyset \neq S \subseteq A} \{b_S, c_S\}$ is a pairing resolving set of G_k .*

Proof. (i) First, we show that A is a metric basis of G_k . Clearly, A forms a resolving set of G_k ; thus, $\dim(G_k) \leq |A| = k$. To show $\dim(G_k) \geq k$, suppose S is a metric basis of G_k with $S \neq A$. If $S \cap A = \emptyset$, then, for any distinct α and β , $S \cap \{b_\alpha, b_\beta, c_\alpha, c_\beta\} \neq \emptyset$; otherwise, $\text{code}_S(b_\alpha) = \text{code}_S(b_\beta)$ and $\text{code}_S(c_\alpha) = \text{code}_S(c_\beta)$; then, $|S| \geq 2^k - 2 \geq k + 1$ for $k \geq 3$. Thus, $S \cap A \neq \emptyset$. By relabeling the vertices of A if necessary, we can assume that $S \cap A = \cup_{i=1}^t \{a_i\}$, where $1 \leq t \leq k - 1$. Then, there are 2^{k-t} vertices of B that are not resolved by $S \cap A$, and there are 2^{k-t} vertices of C that are not resolved by $S \cap A$; thus, $|S \cap (V(G_k) - A)| \geq 2^{k-t} - 1 \geq k - t$, where the last inequality holds since $2^x - x - 1 \geq 0$ for $x \geq 1$. So, $\dim(G_k) = |S| = |S \cap A| + |S \cap (V(G_k) - A)| \geq t + (k - t) = k$. Thus, $\dim(G_k) = k$.

(ii) Suppose $S \neq A$ is a metric basis of G_k ; then, we have $S \cap A \neq \emptyset$ from above argument. If $|S \cap A| = t \leq k - 2$, then $|S \cap (V(G_k) - A)| \geq 2^{k-t} - 1 \geq k - t + 1$ for $k - t \geq 2$, where the last inequality holds since $2^x - x - 2 \geq 0$ for $x \geq 2$. If $|S \cap A| = k - 1$, then there exist $(2^{k-1} - 1)$ pairs in B not resolved by $S \cap A$. So, $|S \cap (V(G_k) - A)| \geq 2^{k-1} - 2 \geq 2$ for $k \geq 3$, and thus $|S| \geq k + 1$. In both cases, we find $|S| > k$, contradicting the assumption of S being a metric basis.

(iii) First, note that B and C are resolving sets of G_k . To see that B resolves G_k , set $B = \{b_1, \dots, b_{2^k-1}\}$. Notice that $\text{code}_B(c_i)$ has 1 in the i th entry and 2 in the rest of its entries, while

$\text{code}_B(a_j)$ has 1 in exactly 2^{k-1} of its entries and 2 in the rest of its entries. And $\text{code}_B(a_i) \neq \text{code}_B(a_j)$ for $i \neq j$, since there is $b \in B$ such that $N(b) \cap A = \{a_i\}$. The set C is seen to be a resolving set by a very similar argument.

Now, let $R = \cup_{\alpha=1}^{2^k-1} \{x_\alpha\}$, where $x_\alpha \in \{b_\alpha, c_\alpha\}$, and assume that $R \neq B$ and $R \neq C$. We show that R resolves any two vertices of G_k by considering memberships of the two vertices with respect to the sets A , B , and C ; there are altogether six cases to consider.

For distinct vertices $a_i, a_j \in A$, there exists a vertex $b_\gamma \in B$ such that $d(b_\gamma, a_i) = 2 = 1 + d(b_\gamma, a_j)$ and $d(c_\gamma, a_i) = 3 = 1 + d(c_\gamma, a_j)$. Since $R \cap \{b_\gamma, c_\gamma\} \neq \emptyset$, $\text{code}_R(a_i) \neq \text{code}_R(a_j)$.

Let distinct vertices $b_\alpha, b_\beta \in B$ be given. If $R \cap \{b_\alpha, b_\beta\} \neq \emptyset$, then $\text{code}_R(b_\alpha) \neq \text{code}_R(b_\beta)$. If $R \cap \{b_\alpha, b_\beta\} = \emptyset$, then $\{c_\alpha, c_\beta\} \subset R$. Then $d(c_\alpha, b_\beta) = 2 = 1 + d(c_\alpha, b_\alpha)$, and thus $\text{code}_R(b_\alpha) \neq \text{code}_R(b_\beta)$. The case of distinct vertices $c_\alpha, c_\beta \in C$ is handled in the same manner.

If $a_i \in A$ and $b_\alpha \in B$, then $\text{code}_R(a_i) \neq \text{code}_R(b_\alpha)$ since $R \cap \{b_\alpha, c_\alpha\} \neq \emptyset$.

Let $a_i \in A$ and $c_\beta \in C$ be given. There exists a vertex $c_\gamma \in R$ such that $d(c_\gamma, c_\beta) \leq 1 < 2 \leq d(c_\gamma, a_i)$, and thus $\text{code}_R(a_i) \neq \text{code}_R(c_\beta)$.

Let $b_\alpha \in B$ and $c_\beta \in C$ be given. If $\alpha = \beta$, then $|R \cap \{b_\alpha, c_\alpha\}| = 1$, and thus $\text{code}_R(b_\alpha) \neq \text{code}_R(c_\beta)$. If $\alpha \neq \beta$ and $R \cap \{b_\alpha, c_\beta\} \neq \emptyset$, then $\text{code}_R(b_\alpha) \neq \text{code}_R(c_\beta)$. If $\alpha \neq \beta$ and $R \cap \{b_\alpha, c_\beta\} = \emptyset$, then there exists $\gamma \notin \{\alpha, \beta\}$ such that $|\{b_\gamma, c_\gamma\} \cap R| = 1$, and this yields $\text{code}_R(b_\alpha) \neq \text{code}_R(c_\beta)$: taking the case $c_\gamma \in R$ for example, we have $d(c_\gamma, b_\alpha) = 2 = 1 + d(c_\gamma, c_\beta)$. \square

From Theorem 3.6, we immediately conclude the following

Corollary 3.7. *For each $k \geq 3$, we have $o(G_k) = \mathcal{R}$ and $R_{\text{MB}}(G_k) > \dim(G_k)$.*

4 Some applications

With the help of the results from the previous section, we now determine $o(G)$ for some classes of graphs G . We also determine $R_{\text{MB}}(G)$ and $R'_{\text{MB}}(G)$ when R^* has a winning strategy, and we determine $S_{\text{MB}}(G)$ and $S'_{\text{MB}}(G)$ when S^* has a winning strategy.

Trees

Fix a tree T . A *support vertex* is a vertex that is adjacent to a vertex of degree one, a *major vertex* is a vertex of degree at least three. A vertex ℓ of degree 1 is called a *terminal vertex* of a major vertex v if $d(\ell, v) < d(\ell, w)$ for every other major vertex w in T . The *terminal degree*, $\text{ter}(v)$, of a major vertex v is the number of terminal vertices of v in T , and an *exterior major vertex* is a major vertex that has positive terminal degree. We denote by $\text{ex}(T)$ the number of exterior major vertices of T , and $\sigma(T)$ the number of leaves of T . Let $M(T)$ be the set of exterior major vertices of T . Let $M_1(T) = \{w \in M(T) : \text{ter}(w) = 1\}$ and let $M_2(T) = \{w \in M(T) : \text{ter}(w) \geq 2\}$; note that $M(T) = M_1(T) \cup M_2(T)$. For each $v \in M(T)$, let T_v be the subtree of T induced by v and all vertices belonging to the paths joining v with its terminal vertices, and let L_v be the set of terminal vertices of v in T .

Theorem 4.1. [6, 23, 25] *If T is a tree that is not a path, then $\dim(T) = \sigma(T) - \text{ex}(T)$.*

Theorem 4.2. [25] *Let T be a tree with $\text{ex}(T) = k \geq 1$, and let v_1, \dots, v_k be the exterior major vertices of T . For each $i \in [k]$, let $\ell_{i,1}, \dots, \ell_{i,\sigma_i}$ be the terminal vertices of v_i with $\text{ter}(v_i) = \sigma_i \geq 1$, and let $P_{i,j}$ be the $v_i - \ell_{i,j}$ path, where $j \in [\sigma_i]$. Let $W \subseteq V(T)$. Then W is a metric basis of T if and only if W contains exactly one vertex from each of the paths $P_{i,j} - v_i$, where $j \in [\sigma_i]$ and $i \in [k]$, with exactly one exception for each $i \in [k]$ and W contains no other vertices of T .*

Theorem 4.3. *If T is a tree that is not a path, then,*

$$o(T) = \begin{cases} \mathcal{S}; & |N(v) \cap L_v| \geq 4 \text{ for some } v \in M_2(T), \\ & \text{or } |N(u) \cap L_u| = 3 = |N(w) \cap L_w| \text{ for distinct } u, w \in M_2(T), \\ \mathcal{N}; & |N(w) \cap L_w| = 3 \text{ for exactly one } w \in M_2(T) \\ & \text{and } |N(v) \cap L_v| \leq 2 \text{ for each } v \in M_2(T) - \{w\}, \\ \mathcal{R}; & |N(v) \cap L_v| \leq 2 \text{ for each } v \in M_2(T). \end{cases}$$

Moreover, if $o(T) = \mathcal{S}$, then $S_{\text{MB}}(T) = S'_{\text{MB}}(T) = 2$; if $o(T) = \mathcal{R}$, then $R_{\text{MB}}(T) = R'_{\text{MB}}(T) = \dim(T) - \sigma(T) - \text{ex}(T)$.

Proof. Let T be a tree that is not a path. Hence $\text{ex}(T) \geq 1$.

First, suppose that there exists an exterior major vertex $x \in M_2(T)$ such that $|N(x) \cap L_x| \geq 4$. Since $N(x) \cap L_x$ is a twin equivalence class of cardinality at least 4, by Proposition 3.2(a), $o(T) = \mathcal{S}$ and $S_{\text{MB}}(T) = S'_{\text{MB}}(T) = 2$.

Second, suppose that $|N(v) \cap L_v| \leq 3$ for each $v \in M_2(T)$. If there exist distinct $x, y \in M_2(T)$ such that $|N(x) \cap L_x| = |N(y) \cap L_y| = 3$, then $N(x) \cap L_x$ and $N(y) \cap L_y$ are distinct twin equivalence classes of cardinality 3; thus, by Proposition 3.2(b), $o(T) = \mathcal{S}$ and $S_{\text{MB}}(T) = S'_{\text{MB}}(T) = 2$.

Now, suppose there exists exactly one $z \in M_2(T)$ with $|N(z) \cap L_z| = 3$, and $|N(v) \cap L_v| \leq 2$ for each $v \in M_2(T) - \{z\}$. Let $L_z = \{\ell'_1, \dots, \ell'_a\}$, where $a \geq 3$ and $d(z, \ell'_i) = 1$ for $i \in [3]$; if $a \geq 4$, let s'_j be the support vertex that lies on the $z - \ell'_j$ path for each $j \in [a] - [3]$. Note that, for any resolving set W of T , Observation 3.1 yields $|W \cap \{\ell'_1, \ell'_2, \ell'_3\}| \geq 2$. If there exists a vertex $v \in M_2(T) - \{z\}$ with $|N(v) \cap L_v| \leq 2$, then, for a fixed $w \in M_2(T) - \{z\}$, let $L_w = \{\ell_1, \dots, \ell_b\}$ such that $d(w, \ell_1) \leq \dots \leq d(w, \ell_b)$; if $b \geq 3$, let s_i be the support vertex that lies on the $w - \ell_i$ path for each $i \in [b] - [2]$. Then $R_w = \{\{\ell_1, \ell_2\}\} \cup (\cup_{i=3}^b \{\{s_i, \ell_i\}\})$ is a dim-pairing resolving set of T_w . In the S-game, S^* can occupy two vertices of $\{\ell'_1, \ell'_2, \ell'_3\}$ after her second move; thus R^* fails to occupy vertices that form a resolving set of G , and hence S^* wins. In the R-game, R^* can occupy two vertices of $\{\ell'_1, \ell'_2, \ell'_3\}$ after his second move, and occupy exactly one vertex of each pair in $(\cup_{i=4}^a \{\{s'_i, \ell'_i\}\}) \cup (\cup_{w \in M_2(T) - \{z\}} R_w)$ thereafter until he completes his $\dim(T)$ th move; thus, the set of vertices selected by R^* , after his $\dim(T)$ th move, forms a resolving set of T , and hence R^* wins. Therefore, $o(G) = \mathcal{N}$.

Third, suppose that $|N(v) \cap L_v| \leq 2$ for each $v \in M_2(T)$. For a fixed $v \in M_2(T)$ with $\text{ter}(v) = k \geq 2$, let ℓ_1, \dots, ℓ_k be the terminal vertices of v such that $d(v, \ell_1) \leq \dots \leq d(v, \ell_k)$; if $d(v, \ell_i) \geq 2$, let s_i be the support vertex that lies on the $v - \ell_i$ path. Then $W_v = \{\{\ell_1, \ell_2\}\} \cup (\cup_{i=3}^k \{\{s_i, \ell_i\}\})$ is a dim-pairing resolving set of T_v , and $\cup_{v \in M_2(T)} W_v$ is a dim-pairing resolving set of T . So, $o(T) = \mathcal{R}$ by Proposition 3.3, and $R_{\text{MB}}(T) = R'_{\text{MB}}(T) = \dim(T)$ by Corollary 3.4. \square

The Petersen graph

For the Petersen graph \mathcal{P} (see Fig. 4) we first recall the following results.

Theorem 4.4. [21] *For the Petersen graph \mathcal{P} , $\dim(\mathcal{P}) = 3$.*

Porism 4.5. [10] *If W is a metric basis of the Petersen graph \mathcal{P} , then the subgraph of \mathcal{P} induced by W is an edge-less graph.*

Lemma 4.6. *Let the vertices of the Petersen graph \mathcal{P} be labeled as in Fig. 4. Let $W_1 = \{u_1, w_2, w_3\}$, $W_2 = \{u_1, u_4, w_2\}$, $W_3 = \{u_1, w_4, w_5\}$, $W_4 = \{u_1, u_3, w_5\}$, $W_5 = \{u_1, u_4, w_3\}$ and $W_6 = \{u_1, u_3, w_4\}$. Then W is a metric basis of \mathcal{P} with $u_1 \in W$ if and only if $W = W_i$ for some $i \in [6]$.*

Proof. (\Leftarrow) Let $W = W_1 = \{u_1, w_2, w_3\}$. Then $\text{code}_W(u_2) = (1, 1, 2)$, $\text{code}_W(u_3) = (2, 2, 1)$, $\text{code}_W(u_4) = (2, 2, 2)$, $\text{code}_W(u_5) = (1, 2, 2)$, $\text{code}_W(w_1) = (1, 2, 1)$, $\text{code}_W(w_4) = (2, 1, 2)$ and $\text{code}_W(w_5) = (2, 1, 1)$. So, W_1 is a metric basis of \mathcal{P} by Theorem 4.4. For $i \in [6] - [1]$, one can easily check that W_i is a metric basis of \mathcal{P} .




Figure 4: The Petersen graph.

(\Rightarrow) Let W be a metric basis of \mathcal{P} with $u_1 \in W$. By Porism 4.5, $W \cap \{u_2, w_1, u_5\} = \emptyset$; thus, $W \cap \{w_2, w_5\} \neq \emptyset$ or $W \cap \{w_3, w_4\} \neq \emptyset$ or $W \cap \{u_3, u_4\} \neq \emptyset$.

First, let $W \cap \{w_2, w_5\} \neq \emptyset$. If $w_2 \in W$, say $X_1 = \{u_1, w_2\} \subset W$, then $W \cap \{u_2, u_5, w_1, w_4, w_5\} = \emptyset$ by Porism 4.5 and $\text{code}_{X_1}(u_3) = \text{code}_{X_1}(u_4) = \text{code}_{X_1}(w_3) = (2, 2)$. Since $d(u_3, w_3) = d(u_3, u_4)$, either $w_3 \in W$ (i.e., $W = W_1$) or $u_4 \in W$ (i.e., $W = W_2$). Similarly, if $w_5 \in W$, then $w_4 \in W$ (i.e., $W = W_3$) or $u_3 \in W$ (i.e., $W = W_4$).

Second, let $W \cap \{w_3, w_4\} \neq \emptyset$. If $w_3 \in W$, say $X_2 = \{u_1, w_3\} \subset W$, then $W \cap \{u_2, u_3, u_5, w_1, w_5\} = \emptyset$ by Porism 4.5 and $\text{code}_{X_2}(u_4) = \text{code}_{X_2}(w_2) = \text{code}_{X_2}(w_4) = (2, 2)$. Since $d(w_4, w_2) = d(w_4, u_4)$, either $w_2 \in W$ (i.e., $W = W_1$) or $u_4 \in W$ (i.e., $W = W_5$). Similarly, if $w_4 \in W$, then $w_5 \in W$ (i.e., $W = W_3$) or $u_3 \in W$ (i.e., $W = W_6$).

Third, let $W \cap \{u_3, u_4\} \neq \emptyset$. If $u_3 \in W$, say $X_3 = \{u_1, u_3\} \subset W$, then $W \cap \{u_2, u_4, u_5, w_1, w_3\} = \emptyset$ by Porism 4.5 and $\text{code}_{X_3}(w_2) = \text{code}_{X_3}(w_4) = \text{code}_{X_3}(w_5) = (2, 2)$. Since $d(w_2, w_4) = d(w_2, w_5)$, either $w_4 \in W$ (i.e., $W = W_6$) or $w_5 \in W$ (i.e., $W = W_4$). Similarly, if $u_4 \in W$, then $w_2 \in W$ (i.e., $W = W_2$) or $w_3 \in W$ (i.e., $W = W_5$). \square

Theorem 4.7. $o(\mathcal{P}) = \mathcal{R}$ and $R_{\text{MB}}(\mathcal{P}) = R'_{\text{MB}}(\mathcal{P}) = 3$.

Proof. Let the vertices of \mathcal{P} be labeled as in Fig. 4, and let $A_1 = \{w_2, w_5\}$, $A_2 = \{w_3, w_4\}$, and $A_3 = \{u_3, u_4\}$. First, we consider the R-game. Since \mathcal{P} is vertex-transitive (see [19]), we may assume that R^* occupies u_1 after his first move. If S^* selects a vertex in $N(u_1)$ on her first move, R^* can select a vertex of an A_i , $i \in [3]$, on his second move; if S^* selects a vertex of an A_j , $j \in [3]$, on her first move, then R^* can select the other vertex of A_j on his second move. If R^* selects a vertex of $A_1 = \{w_2, w_5\}$, say w_2 , on his second move, he can select a vertex in $\{u_4, w_3\}$ on his third move; if R^* selects a vertex of $A_2 = \{w_3, w_4\}$, say w_3 , on his second move, he can select a vertex of $\{u_4, w_2\}$ on his third move; if R^* selects a vertex of $A_3 = \{u_3, u_4\}$, say u_3 , on his second move, he can select a vertex in $\{w_4, w_5\}$ on his third move. In each case, the set of vertices occupied by R^* , after his third move, forms a resolving set of \mathcal{P} by Lemma 4.6.

Second, we consider the S-game. Since \mathcal{P} is edge-transitive (see [19]), we may assume that S^* selects u_5 on her first move and R^* selects u_1 on his first move. If S^* selects a vertex of an A_i , $i \in [3]$, on her second move, then R^* can select the other vertex of A_i on his second move; if S^* selects a vertex of $N(u_1) - \{u_5\} = \{u_2, w_1\}$ on her second move, R^* can select a vertex of an A_j , $j \in [3]$, on his second move. By applying the above argument for the R-game, it is easy to see that R^* can occupy a resolving set of \mathcal{P} after his third move.

Thus, $o(\mathcal{P}) = \mathcal{R}$ and $R_{\text{MB}}(\mathcal{P}) = R'_{\text{MB}}(\mathcal{P}) = 3$. \square

Bouquet of cycles

Let B_m , $m \geq 2$, be a bouquet of m cycles (i.e., the vertex sum of m cycles at one common vertex), and let w be the cut-vertex of B_m (see Fig. 5). Let C^1, \dots, C^m be the m cycles of B_m . For each $i \in [m]$, let $P^i = C^i - w$.




Figure 5: A bouquet of four cycles B_4 .

Theorem 4.8. [20] *If B_m is a bouquet of $m \geq 2$ cycles of which x cycles are even, then*

$$\dim(B_m) = \begin{cases} m; & x = 0, \\ m + x - 1; & x \geq 1. \end{cases}$$

Lemma 4.9. [20] *If W is a resolving set of a bouquet of cycles B_m , $m \geq 2$, then*

- (a) *for each $i \in [m]$, $|W \cap V(P^i)| \geq 1$; and*
- (b) *for any two distinct even cycles C^i and C^j of B_m , $|W \cap (V(P^i) \cup V(P^j))| \geq 3$.*

Theorem 4.10. *If B_m is a bouquet of $m \geq 2$ cycles of which z are 4-cycles, then*

$$o(B_m) = \begin{cases} \mathcal{R}; & z \leq 2, \\ \mathcal{N}; & z = 3, \\ \mathcal{S}; & z \geq 4. \end{cases}$$

Moreover, if $o(B_m) = \mathcal{R}$, then $R_{\text{MB}}(B_m) = R'_{\text{MB}}(B_m) = \dim(B_m)$; if $o(B_m) = \mathcal{S}$, then $S_{\text{MB}}(B_m) = S'_{\text{MB}}(B_m) = 4$.

Proof. Let w be the cut-vertex of B_m , where $m \geq 2$. Let C^1, \dots, C^z be cycles isomorphic to C_4 , let C^{z+1}, \dots, C^x be even cycles that are not isomorphic to C_4 , and let C^{x+1}, \dots, C^m be odd cycles of B_m ; notice $x \geq z$. If C^i is an odd cycle of length $2k_i + 1$, let C^i be given by $w, u_{i,1}, u_{i,2}, \dots, u_{i,k_i}, u_{i,k_i+1}, \dots, u_{i,2k_i}, w$; if C^j is an even cycle of length $2k_j$, let C^j be given by $w, u_{j,1}, u_{j,2}, \dots, u_{j,k_j-1}, u_{j,k_j}, u_{j,k_j+1}, \dots, u_{j,2k_j-1}, w$ (see Fig. 5 for the labeling of the vertices of a B_4). We note that $u_{i,1}$ and $u_{i,3}$ are twins for each $i \in [z]$.

Case 1: $z \leq 2$.

If $z = 0$ and $x = 0$, then $\cup_{i=1}^m \{\{u_{i,k_i}, u_{i,k_i+1}\}\}$ is a dim-pairing resolving set of B_m . If $z = 0$ and $x \geq 1$, then $\{\{u_{1,k_1-1}, u_{1,k_1+1}\}\} \cup (\cup_{i=2}^x \{\{u_{i,1}, u_{i,k_i-1}\}, \{u_{i,k_i+1}, u_{i,2k_i-1}\}\}) \cup (\cup_{j=x+1}^m \{\{u_{j,k_j}, u_{j,k_j+1}\}\})$ is a dim-pairing resolving set of B_m . If $z = 1$, then

$$\{\{u_{1,1}, u_{1,3}\}\} \cup (\cup_{i=2}^x \{\{u_{i,1}, u_{i,k_i-1}\}, \{u_{i,k_i+1}, u_{i,2k_i-1}\}\}) \cup (\cup_{j=x+1}^m \{\{u_{j,k_j}, u_{j,k_j+1}\}\})$$

is a dim-pairing resolving set of B_m . If $z = 2$, then

$$\begin{aligned} \{\{u_{1,1}, u_{1,3}\}, \{u_{2,1}, u_{2,3}\}, \{u_{1,2}, u_{2,2}\}\} &\cup (\cup_{i=3}^x \{\{u_{i,1}, u_{i,k_i-1}\}, \{u_{i,k_i+1}, u_{i,2k_i-1}\}\}) \\ &\cup (\cup_{j=x+1}^m \{\{u_{j,k_j}, u_{j,k_j+1}\}\}) \end{aligned}$$

is a dim-pairing resolving set of B_m . So, in each case, $o(B_m) = \mathcal{R}$ by Proposition 3.3 and $R_{\text{MB}}(B_m) = R'_{\text{MB}}(B_m) = \dim(B_m)$ by Corollary 3.4.

Case 2: $z = 3$.

Note that $|\cup_{i=1}^3 V(P^i)| = 9$ and, for any resolving set W of B_m , $|W \cap (\cup_{i=1}^3 V(P^i))| \geq 5$ by Lemma 4.9(b). In the S-game, S^* can occupy five vertices of $\cup_{i=1}^3 V(P^i)$ after her fifth move.

Thus, R^* fails to occupy vertices that form a resolving set of B_m . In the R-game, R^* can occupy one vertex of each pair in $(\cup_{i=1}^3 \{\{u_{i,1}, u_{i,3}\}\}) \cup (\cup_{i=4}^x \{\{u_{i,1}, u_{i,k_i-1}\}, \{u_{i,k_i+1}, u_{i,2k_i-1}\}\}) \cup (\cup_{j=x+1}^m \{\{u_{j,k_j}, u_{j,k_j+1}\}\})$ and two vertices of $\cup_{i=1}^3 \{u_{i,2}\}$; thus, R^* can occupy vertices that form a resolving set of B_m . So, $o(B_m) = \mathcal{N}$.

Case 3: $z \geq 4$.

Note that $|\cup_{i=1}^4 V(P^i)| = 12$ and, for any resolving set W of B_m , $|W \cap (\cup_{i=1}^4 V(P^i))| \geq 7$ by Lemma 4.9(b). Regardless of whether S^* plays first or second, S^* can occupy 6 vertices of $\cup_{i=1}^4 V(P^i)$ after her sixth move. So, R^* fails to occupy vertices that form a resolving set of B_m ; thus $o(B_m) = \mathcal{S}$. In determining $S_{\text{MB}}(B_m)$ and $S'_{\text{MB}}(B_m)$, we note that the optimal strategy for R^* is to occupy at least a vertex in each pair of $\cup_{i=1}^z \{\{u_{i,1}, u_{i,3}\}\}$, and the optimal strategy for S^* is to occupy two vertices each in $V(P^i)$ and $V(P^j)$ for distinct $i, j \in [z]$. By relabeling the vertices of $\cup_{i=1}^z V(P^i)$ if necessary, we may assume that the two players occupy the vertices of B_m in the order of $V(P^1), \dots, V(P^z)$. In the S-game, S^* can occupy two vertices of $V(P^1)$ after her second move, R^* would have occupied a vertex in $\{u_{1,1}, u_{1,3}\} \subset V(P^1)$ and a vertex in $\{u_{2,1}, u_{2,3}\} \subset V(P^2)$ after his second move, and S^* can occupy two vertices of $V(P^3)$ on her third and fourth move; thus, S^* wins after her fourth move. In the R-game, R^* occupies a vertex in $\{u_{1,1}, u_{1,3}\}$ after his first move, and S^* can occupy two vertices of $V(P^2)$ after her second move (R^* would have occupied a vertex in $\{u_{2,1}, u_{2,3}\}$ on his second move). If R^* occupies a vertex in $\cup_{i=1}^3 V(P^i)$ that has not yet been taken on his third move, S^* can occupy two vertices of $V(P^4)$ on her third and fourth move. So, in the R-game, S^* wins after her fourth move. Thus, $S_{\text{MB}}(B_m) = S'_{\text{MB}}(B_m) = 4$. \square

Complete multi-partite graphs

The metric dimension of complete multi-partite graphs was determined in [26].

Theorem 4.11. [26] *If $G = K_{a_1, \dots, a_k}$, where $k \geq 2$, $n = \sum_{i=1}^k a_i$, and s is the number of partite sets of G consisting of one element, then*

$$\dim(G) = \begin{cases} n - k; & s = 0, \\ n + s - k - 1; & s \neq 0. \end{cases}$$

For the MBRG we have the following description.

Theorem 4.12. *If $G = K_{a_1, \dots, a_k}$, where $k \geq 2$, and s is the number of partite sets of G consisting of one element, then*

$$o(G) = \begin{cases} \mathcal{S}; & s \geq 4 \text{ or } a_i \geq 4 \text{ for some } i \in [k], \\ & \text{or } s = a_i = 3 \text{ for some } i \in [k], \\ & \text{or } a_i = a_j = 3 \text{ for distinct } i, j \in [k], \\ \mathcal{N}; & s = 3 \text{ and } a_i \leq 2 \text{ for each } i \in [k], \\ & \text{or } s \leq 2 \text{ and } a_i = 3 \text{ for exactly one } i \in [k], \\ \mathcal{R}; & \max\{s, a_i\} \leq 2 \text{ for each } i \in [k]. \end{cases}$$

Moreover, if $o(G) = \mathcal{S}$, then $S_{\text{MB}}(G) = S'_{\text{MB}}(G) = 2$; if $o(G) = \mathcal{R}$, then $R_{\text{MB}}(G) = R'_{\text{MB}}(G) = \dim(G)$.

Proof. Let $V(G)$ be partitioned into V_1, \dots, V_k such that $V_i = \{u_{i,1}, \dots, u_{i,a_i}\}$ with $|V_i| = a_i$, where $i \in [k]$ and $k \geq 2$. We may without loss of generality assume that $a_1 \leq \dots \leq a_k$.

First, suppose that $s \geq 4$ or $a_i \geq 4$ for some $i \in [k]$. If $s \geq 4$, then $\cup_{i=1}^s V_i$ is a twin equivalence class of cardinality at least 4. If $a_i \geq 4$ for some $i \in [k]$, then V_k is a twin equivalence class of cardinality at least 4. By Proposition 3.2(a), $o(G) = \mathcal{S}$ and $S_{\text{MB}}(G) = S'_{\text{MB}}(G) = 2$.

Second, suppose that $\max\{s, a_i\} \leq 3$ for each $i \in [k]$; further, let $s = 3$ or $a_i = 3$ for some $i \in [k]$. If $s = a_x = 3$ for some $x \in [k]$ or $a_y = a_z = 3$ for distinct $y, z \in [k]$, then G has distinct twin equivalence classes of cardinality three; thus, by Proposition 3.2(b), $o(G) = \mathcal{S}$ and $S_{\text{MB}}(G) = S'_{\text{MB}}(G) = 2$.

So, suppose $s = 3$ or $a_i = 3$ for exactly one $i \in [k]$, but not both. Let W be any resolving set of G . By Observation 3.1, we have the following: (1) if $s = 3$, then $|W \cap \{u_{1,1}, u_{2,1}, u_{3,1}\}| \geq 2$; (2) if $a_i = 3$ for exactly one $i \in [k]$, then $a_k = 3$ and $|W \cap V_k| \geq 2$. In the S-game, S^* can occupy two vertices of $\{u_{1,1}, u_{2,1}, u_{3,1}\}$ after her second move (when $s = 3$), or S^* can occupy two vertices of $V_k = \{u_{k,1}, u_{k,2}, u_{k,3}\}$ after her second move (when $a_k = 3$); thus, in each case, R^* fails to occupy vertices that form a resolving set of G . Now, we consider the R-game. If $s = 3$ (and thus $a_i \leq 2$ for each $i \in [k]$), then R^* can occupy two vertices of $\{u_{1,1}, u_{2,1}, u_{3,1}\}$ after his second move, and occupy exactly one vertex of each pair in $\cup_{j=4}^k \{\{u_{j,1}, u_{j,2}\}\}$ thereafter until he completes his $(k-1)^{\text{th}}$ move. If $a_k = 3$ (and thus $s \leq 2$ and $a_i \leq 2$ for each $i \in [k-1]$), then R^* can occupy two vertices of $V_k = \{u_{k,1}, u_{k,2}, u_{k,3}\}$, after his second move, and occupy additional vertices (if any) thereafter as follow: (1) if $s = 0$, then R^* can occupy exactly one vertex of each pair in $\cup_{i=1}^{k-1} \{\{u_{i,1}, u_{i,2}\}\}$; (2) if $s = 1$, then R^* can occupy exactly one vertex of each pair in $\cup_{i=2}^{k-1} \{\{u_{i,1}, u_{i,2}\}\}$; (3) if $s = 2$, then R^* can occupy exactly one vertex of each pair in $\{\{u_{1,1}, u_{2,1}\}\} \cup (\cup_{i=3}^{k-1} \{\{u_{i,1}, u_{i,2}\}\})$. So, in each case of the R-game, the vertices chosen by R^* form a resolving set of G ; thus, R^* wins. Therefore, $o(G) = \mathcal{N}$.

Third, suppose that $\max\{s, a_i\} \leq 2$ for each $i \in [k]$. If $s = 0$, then $\cup_{i=1}^k \{\{u_{i,1}, u_{i,2}\}\}$ is a dim-pairing resolving set of G . If $s = 1$, then $\cup_{i=2}^k \{\{u_{i,1}, u_{i,2}\}\}$ is a dim-pairing resolving set of G . If $s = 2$, then $\{\{u_{1,1}, u_{2,1}\}\} \cup (\cup_{i=3}^k \{\{u_{i,1}, u_{i,2}\}\})$ is a dim-pairing resolving set of G . In each case, $o(G) = \mathcal{R}$ by Proposition 3.3 and $R_{\text{MB}}(G) = R'_{\text{MB}}(G) = \dim(G)$ by Corollary 3.4. \square

Some grid-like graphs

The *Cartesian product* $G \square H$ of graphs G and H is the graph with the vertex set $V(G) \times V(H)$ such that (u, v) is adjacent to (u', v') if and only if either $u = u'$ and $vv' \in E(H)$, or $v = v'$ and $uu' \in E(G)$. Products $P_s \square P_t$ are known as *grid graphs*. For the rest of this section we set $V(P_s) = \{u_1, \dots, u_s\}$ and $V(P_t) = \{v_1, \dots, v_t\}$, see Fig. 6 for the labeling of $P_8 \square P_4$.




Figure 6: Labeling of $P_8 \square P_4$.

Recall the following results of grid graphs.

Proposition 4.13. [5] *If $s, t \geq 2$, then $\dim(P_s \square P_t) = 2$.*

Lemma 4.14. [10] *Let $s, t \geq 2$, and let $W_1 = \{(u_1, v_1), (u_1, v_t)\}$, $W_2 = \{(u_1, v_1), (u_s, v_1)\}$, $W_3 = \{(u_1, v_t), (u_s, v_t)\}$, and $W_4 = \{(u_s, v_1), (u_s, v_t)\}$. Then W is a metric basis of $P_s \square P_t$ if and only if $W = W_i$ for some $i \in [4]$.*

For the MBRG on grid graphs we have:

Proposition 4.15. *If $s, t \geq 2$, then $o(P_s \square P_t) = \mathcal{R}$ and*

$$R_{\text{MB}}(P_s \square P_t) = R'_{\text{MB}}(P_s \square P_t) = \dim(P_s \square P_t) = 2.$$

Proof. Since $\{(u_1, v_1), (u_s, v_t)\}, \{(u_s, v_1), (u_1, v_t)\}$ is a dim-pairing resolving set of $P_s \square P_t$, Proposition 3.3 implies that $o(P_s \square P_t) = \mathcal{R}$. Moreover, $R_{\text{MB}}(P_s \square P_t) = R'_{\text{MB}}(P_s \square P_t) = \dim(P_s \square P_t)$ by Corollary 3.4. \square

Continuing with some grid-related graphs, we next study how the MBRG behaves on the torus grid graphs, that is, the Cartesian product of cycles. We recall the following results that will be used in proving Proposition 4.19.

Theorem 4.16. [5] *If $s, t \geq 3$, then*

$$\dim(C_s \square C_t) = \begin{cases} 3; & s \text{ or } t \text{ is odd,} \\ 4; & s \text{ and } t \text{ are even.} \end{cases}$$

Proposition 4.17. [5] *Let $s, t \geq 3$ be integers.*

- (a) *If s is odd, let $W = \{x, y, z\} \subseteq V(C_s \square C_t)$ such that x, y are diametral in a copy of C_s , and z is adjacent to x in a copy of C_t . Then W is a metric basis for $C_s \square C_t$.*
- (b) *If s and t are even, let $W = \{x, y, z, w\} \subseteq V(C_s \square C_t)$ such that x, y are diametral in a copy of C_s , z is adjacent to x in a copy of C_s , and w is adjacent to x in a copy of C_t . Then W is a metric basis for $C_s \square C_t$.*

Lemma 4.18. [5] *For even integers $s, t \geq 4$, let W be a resolving set of $C_s \square C_t$ with $u \in W$. If u and u' are diametral in $C_s \square C_t$, then $(W - \{u\}) \cup \{u'\}$ is also a resolving set of $C_s \square C_t$.*

Proposition 4.19. *If $s, t \geq 3$, then $o(C_s \square C_t) = \mathcal{R}$ and*

$$R_{\text{MB}}(C_s \square C_t) = R'_{\text{MB}}(C_s \square C_t) = \dim(C_s \square C_t).$$

Proof. Let $s, t \geq 3$ be integers and consider the following two cases.

Case 1: s and t are even.

Let $W = \{x, y, z, w\}$ be a metric basis for $C_s \square C_t$ as described in Proposition 4.17(b); note that no two vertices in W are diametral in $C_s \square C_t$. Let x', y', z' , and w' be diametral vertices of x, y, z , and w , respectively, in $C_s \square C_t$. Then Lemma 4.18 implies that $\{\{x, x'\}, \{y, y'\}, \{z, z'\}, \{w, w'\}\}$ is a dim-pairing resolving set of $C_s \square C_t$, and hence Proposition 3.3 and Corollary 3.4 yield the conclusion.

Case 2: s is odd.

First, we consider the R -game. Suppose R^* selects an arbitrary vertex, say x , in $C_s \square C_t$ on his first move. Note that there are two distinct vertices, say y and y' , that are diametral to x in the copy of C_s containing x and there are two distinct vertices, say z and z' , that are adjacent to x in the copy of C_t containing x . By Proposition 4.17(a), the set $\{x, y^*, z^*\}$, where $y^* \in \{y, y'\}$ and $z^* \in \{z, z'\}$, is a metric basis for $C_s \square C_t$. Since R^* can select a vertex in $\{y, y'\}$ and a vertex in $\{z, z'\}$ in his second and third move, R^* wins the R -game.

Second, we consider the S -game. Suppose S^* selects a vertex, say x' , in $C_s \square C_t$ on her first move. Then R^* can choose a neighbor x of x' in the copy of C_t that contains x' . Note that there are two distinct vertices, say y and y' , that are diametral to x in the copy of C_s containing x and there are two distinct vertices, say z_1 and z'_1 (z_2 and z'_2 , respectively), that are adjacent to y (y' , respectively) in the copy of C_t containing y (y' , respectively). By Proposition 4.17(a), both $\{x, y, z_1^*\}$ and $\{x, y', z_2^*\}$, where $z_1^* \in \{z_1, z'_1\}$ and $z_2^* \in \{z_2, z'_2\}$, form metric bases for $C_s \square C_t$. So, R^* can select a vertex in $\{y, y'\}$ on his second move. If R^* selects y on his second move, he can select a vertex in $\{z_1, z'_1\}$ in his third move; if R^* selects y' on his second move, he can select a vertex in $\{z_2, z'_2\}$ in his third move. In each case, R^* wins the S -game.

Therefore, we conclude that $o(C_s \square C_t) = \mathcal{R}$ and $R_{\text{MB}}(C_s \square C_t) = R'_{\text{MB}}(C_s \square C_t) = \dim(C_s \square C_t)$. \square

We conclude this paper with some open problems.

Question 4.20. *It is known that determining the metric dimension of a general graph is an NP-hard problem (see [11]). What can we say about the computational complexity of determining the outcome of MBRG?*

Question 4.21. *For product graphs G , such as the Cartesian product, the lexicographic product, the corona product, and the direct product, can we determine $o(G)$, as well as $R_{\text{MB}}(G)$ and $R'_{\text{MB}}(G)$ (or $S_{\text{MB}}(G)$ and $S'_{\text{MB}}(G)$)?*

Question 4.22. *It is easy to see that $R_{\text{MB}}(G) = 1$ ($R'_{\text{MB}}(G) = 1$, respectively) if and only if $G = P_n$ for $n \geq 2$. For any positive integer $k \in [\lfloor \frac{n(G)}{2} \rfloor] - \{1\}$, can we characterize graphs G satisfying $R_{\text{MB}}(G) = k$ as well as $R'_{\text{MB}}(G) = k$?*

Acknowledgements

S.K. acknowledges the financial support from the Slovenian Research Agency (research core funding No. P1-0297 and projects J1-9109, J1-1693, N1-0095, N1-0108). I.G.Y. began to conduct research on this project while he was visiting the University of Ljubljana, Slovenia, supported by “Ministerio de Educación, Cultura y Deporte”, Spain, under the “José Castillejo” program for young researchers (reference number: CAS18/00030).

References

- [1] S. Akhter, R. Farooq, Metric dimension of fullerene graphs, *Electron. J. Graph Theory Appl.* 7 (2019) 91–103.
- [2] R. F. Bailey, P. J. Cameron, Base size, metric dimension and other invariants of groups and graphs, *Bull. London Math. Soc.* 43 (2011) 209–242.
- [3] J. Beck, *Combinatorial Games. Tic-Tac-Toe Theory*, Cambridge University Press, Cambridge, 2008.
- [4] B. Brešar, S. Klavžar, D. F. Rall, Domination game and an imagination strategy, *SIAM J. Discrete Math.* 24 (2010) 979–991.
- [5] J. Cáceres, C. Hernando, M. Mora, I. M. Pelayo, M. L. Puertas, C. Seara, D. R. Wood, On the metric dimension of Cartesian products of graphs, *SIAM J. Discrete Math.* 21 (2007) 423–441.
- [6] G. Chartrand, L. Eroh, M. A. Johnson, O. R. Oellermann, Resolvability in graphs and the metric dimension of a graph, *Discrete Appl. Math.* 105 (2000) 99–113.
- [7] G. Chartrand, P. Zhang, The theory and applications of resolvability in graphs. A Survey, *Congr. Numer.* 160 (2003) 47–68.
- [8] E. Duchêne, V. Gledel, A. Parreau, G. Renault, Maker-Breaker domination game, *Discrete Math.* 343 (2020) Article 111955, 12 pp.
- [9] P. Erdős, J. L. Selfridge, On a combinatorial game, *J. Combin. Theory Ser. A* 14 (1973) 298–301.
- [10] L. Eroh, C. X. Kang, E. Yi, The connected metric dimension at a vertex of a graph, *Theoret. Comput. Sci.* 806 (2020) 53–69.
- [11] M. R. Garey, D. S. Johnson, *Computers and Intractability: A Guide to the Theory of NP-completeness*, Freeman, New York, 1979.
- [12] V. Gledel, M. A. Henning, V. Iršič, S. Klavžar, Maker-Breaker total domination game, *Discrete Appl. Math.* (2019) <https://doi.org/10.1016/j.dam.2019.11.004>.

- [13] V. Gledel, V. Iršič, S. Klavžar, Maker-Breaker domination number, *Bull. Malays. Math. Sci. Soc.* 42 (2019) 1773–1789.
- [14] R. Hammack, W. Imrich, S. Klavžar, *Handbook of Product Graphs: Second Edition*, CRC Press, Boca Raton, FL, 2011.
- [15] R. Hancock, The Maker-Breaker Rado game on a random set of integers, *SIAM J. Discrete Math.* 33 (2019) 68–94.
- [16] F. Harary, R. A. Melter, On the metric dimension of a graph, *Ars Combin.* 2 (1976) 191–195.
- [17] D. Hefetz, M. Krivelevich, M. Stojaković, T. Szabó, *Positional Games*, Birkhäuser/Springer, Basel, 2014.
- [18] C. Hernando, M. Mora, I. M. Pelayo, C. Seara, D. R. Wood, Extremal graph theory for metric dimension and diameter, *Electron. J. Combin.* (2010) #R30.
- [19] D. A. Holton, J. Sheehan, *The Petersen Graph*, Cambridge University Press, 1993.
- [20] H. Iswadi, E. T. Baskoro, A. N. M. Salman, R. Simanjuntak, The metric dimension of amalgamation of cycles, *Far East J. Math. Sci.* 41 (2010) 19–31.
- [21] I. Javaid, M. T. Rahim, K. Ali, Families of regular graphs with constant metric dimension, *Util. Math.* 75 (2008) 21–33.
- [22] Z. Jiang, N. Polyanskii, On the metric dimension of Cartesian powers of a graph, *J. Combin. Theory Ser. A* 165 (2019) 1–14.
- [23] S. Khuller, B. Raghavachari, A. Rosenfeld, Landmarks in graphs, *Discrete Appl. Math.* 70 (1996) 217–229.
- [24] M. Mikalački, M. Stojaković, Fast strategies in biased Maker-Breaker games, *Discrete Math. Theor. Comput. Sci.* 20 (2018) Paper No. 6, 25pp.
- [25] C. Poisson, P. Zhang, The metric dimension of unicyclic graphs, *J. Combin. Math. Combin. Comput.* 40 (2002) 17–32.
- [26] S. W. Saputro, E. T. Baskoro, A. N. M. Salman, D. Suprijanto, The metric dimension of a complete n -partite graph and its Cartesian product with a path, *J. Combin. Math. Combin. Comput.* 71 (2009) 283–293.
- [27] P. J. Slater, Leaves of trees, *Congr. Numer.* 14 (1975) 549–559.
- [28] L. du Toit, T. Vetrík, On the metric dimension of circulant graphs with 2 generators, *Kragujevac J. Math.* 43 (2019) 49–58.