

On the variety of general position problems under vertex and edge removal

Jing Tian^{a,*} Pakanun Dokyeesun^{b,†} Sandi Klavžar^{c,b,d,‡}

February 4, 2026

^a School of Science, Zhejiang University of Science and Technology, Hangzhou, Zhejiang 310023, PR China

^b Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia

^c Faculty of Mathematics and Physics, University of Ljubljana, Slovenia

^d Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia

Abstract

Let $gp_t(G)$, $gp_o(G)$, and $gp_d(G)$ be the total, the outer, and the dual general position number of a graph G , respectively. This paper investigates how removing a vertex or removing an edge affects these graph invariants. It is proved that if x is not a cut vertex, then $gp_t(G) - 1 \leq gp_t(G - x) \leq gp_t(G) + \deg_G(x)$. On the other hand, $gp_o(G - x)$ and $gp_d(G - x)$ can be respectively arbitrarily larger/smaller than $gp_o(G)$ and $gp_d(G)$. On the positive side, it is proved that if x lies in some gp_o -set, then $gp_o(G) - 1 \leq gp_o(G - x)$, and that if x is not a cut vertex and lies in some gp_d -set of G , then $gp_d(G) - 1 \leq gp_d(G - x)$. For the edge removal, it is proved that (i) $gp_t(G) - |S(G)_e| \leq gp_t(G - e) \leq gp_t(G) + 2$, where $S(G)_e$ is the set of simplicial vertices adjacent to both endvertices of e , (ii) $gp_o(G)/2 \leq gp_o(G - e) \leq 2gp_o(G)$, and (iii) that $gp_d(G) - gp_d(G - e)$ can be arbitrarily large. All bounds are demonstrated to be sharp.

Keywords: total general position; outer general position; dual general position; vertex-deleted subgraph; edge-deleted subgraph

AMS Subj. Class. (2020): 05C12, 05C69

^{*}jingtian526@126.com

[†]pakanun@gmail.com

[‡]sandi.klavzar@fmf.uni-lj.si

1 Introduction

The concept of a general position set was introduced into graph theory independently in two papers, namely in [4, 13], but in the case of hypercubes, implicitly also much earlier in [10]. It was the article [13] that has stimulated a great deal of interest in this concept. The survey [3] provides a systematic review of the results, in addition we point to the following selected recent publications [2, 5, 8, 9, 11, 12, 15–17, 20].

For a general position set, that is, a set in which no three vertices lie on a common shortest path, one requires that the vertices of the set are in general position. This requirement can be generalized by requiring that the vertices of some set lie in general position with respect to the selected set. Using this idea, and considering the natural selections for sets to be considered, in [18] a variety of general position sets was defined as follows. First, for a given graph $G = (V(G), E(G))$ and a set of vertices $Z \subseteq V(G)$, two vertices $x, y \in V(G)$ are said to be *Z-positionable* if any shortest x, y -path intersects Z at most in x and y , that is, no inner vertex of the path lies in Z . Second, if $\bar{Z} = V(G) \setminus Z$, then the set Z is

- (i) a *general position set*, if every $u, v \in Z$ are *Z-positionable*;
- (ii) a *total general position set*, if every $u, v \in V(G)$ are *Z-positionable*;
- (iii) an *outer general position set*, if every $u, v \in Z$ are *Z-positionable*, and every $u \in Z, v \in \bar{Z}$ are *Z-positionable*; and
- (iv) a *dual general position set*, if every $u, v \in Z$ are *Z-positionable*, and every $u, v \in \bar{Z}$ are *Z-positionable*.

Typically, we are interested in the cardinalities of largest such sets, which are respectively denoted by $gp(G)$, $gp_t(G)$, $gp_o(G)$, and $gp_d(G)$, and called the *general position number*, the *total general position number*, the *outer general position number*, and the *dual general position number* of G , respectively. If $\tau \in \{gp, gp_t, gp_o, gp_d\}$, then a set Z is a τ -set if it has the corresponding property and $|Z| = \tau(G)$. Note that, by definition, $gp_t(G) \leq gp_o(G) \leq gp(G)$ and $gp_t(G) \leq gp_d(G) \leq gp(G)$. After being introduced in [18], this variety has been also investigated on strong and lexicographic products of graphs [6].

The paper [7] investigates how removing a vertex or an edge affects the general position number of a graph. It is proved that $gp(G-x) \leq 2gp(G)$ holds for any vertex x of a connected graph G and that if x lies in some gp -set of G , then $gp(G) - 1 \leq gp(G-x)$. On the other hand, $gp(G-x)$ can be much larger than $gp(G)$ also when $G-x$ is connected. For the edge removal it is proved that $gp(G)/2 \leq gp(G-e) \leq 2gp(G)$ holds for any edge e of G . In this paper we continue this line of research

by investigating how removing a vertex or an edge affects the other three general position numbers.

In the next section we give definitions and recall results needed in the rest of the paper. In Section 3 we consider the vertex removal. For the total general position sets we prove that if x is not a cut vertex, then $gp_t(G) - 1 \leq gp_t(G - x) \leq gp_t(G) + \deg_G(x)$. Section 3.2 addresses the outer general position number of a graph when a vertex is removed and shows that it cannot be bounded in terms of the outer general position number of the original graph. On the other hand, if x lies in some gp_o -set, then $gp_o(G) - 1 \leq gp_o(G - x)$. Moreover, if x is a simplicial vertex, then $gp_o(G - x) \leq gp_o(G) + \deg_G(x) - 1$. Section 3.3 then deals with the dual general position number. Similarly, as for the outer general position number, $gp_d(G - x)$ can be arbitrarily larger/smaller than $gp_d(G)$. On the other hand, if x is not a cut vertex and lies in some gp_d -set of G , then $gp_d(G) - 1 \leq gp_d(G - x)$. In the final section we consider the edge removal operation. For the total and the outer general position number we prove sharp lower and upper bounds, while for the dual general position number we show that the difference $gp_d(G) - gp_d(G - e)$ can be arbitrarily large. All bounds proved in this paper are demonstrated to be sharp.

2 Preliminaries

We consider simple and connected graphs $G = (V(G), E(G))$. For a positive integer k , the set $\{1, \dots, k\}$ is denoted by $[k]$.

Let $u \in V(G)$. Then $N_G(u)$ denotes the set of neighbors of u in G and $N_G[u] = N_G(u) \cup \{u\}$. The *degree* $\deg_G(u)$ of u is $\deg_G(u) = |N_G(u)|$. If $X \subseteq V(G)$, then the subgraph of G induced by X is denoted by $G[X]$. The graph $G - u$ is obtained by deleting u and all incident edges from G , that is, $G - u$ is the subgraph $G[V(G) \setminus \{u\}]$. For an edge $e \in E(G)$, the graph $G - e$ is obtained by deleting the edge e from G . A vertex u of G is *simplicial* if $N_G(u)$ induces a complete subgraph. The set of all simplicial vertices of G will be denoted by $S(G)$ and the cardinality of $S(G)$ by $s(G)$. Moreover, $\omega(G)$ and $\alpha(G)$ stand for the clique number and the independence number of G .

The *distance* $d_G(u, v)$ between vertices u and v of G is the number of edges on a shortest u, v -path. A set $X \subseteq V(G)$ is *convex* if for any vertices $u, v \in X$, any shortest u, v -path contains only vertices from X . By abuse of language, we will also say that a subgraph H of G is convex if $V(H)$ is convex. The vertex u of G is *maximally distant* from a vertex $v \in V(G)$ if every $w \in N_G(u)$ satisfies $d_G(v, w) \leq d_G(u, v)$. If u is maximally distant from v , and v is also maximally distant from u , then u and v are *mutually maximally distant*, MMD for short. Note

that true twins, that is, vertices u and v with $N_G[u] = N_G[v]$, are MMD. The *strong resolving graph* G_{SR} of G has $V(G)$ as the vertex set, two vertices being adjacent in G_{SR} if they are MMD in G . This concept was introduced in [14] for the purpose of better understanding the strong metric dimension of graphs.

The first known result which we need later on describes general position sets in an arbitrary graph. To state it, some more definitions are required. If $\mathcal{P} = \{X_1, \dots, X_t\}$ is a partition of $X \subseteq V(G)$, then \mathcal{P} is *distance-constant* if for any $i, j \in [t]$, $i \neq j$, there exists a constant p_{ij} , such that $d_G(x, y) = p_{ij}$ for every $x \in X_i$, $y \in X_j$. If so, we set $d_G(X_i, X_j) = p_{ij}$. A distance-constant partition \mathcal{P} is *in-transitive* if $p_{ik} \neq p_{ij} + p_{jk}$ holds for $i, j, k \in [t]$.

Theorem 2.1 [1, Theorem 3.1] *Let G be a graph. Then $X \subseteq V(G)$ is a general position set if and only if the components of $G[X]$ are complete subgraphs, the vertices of which form an in-transitive, distance-constant partition of X .*

The following theorem is the most important for us since it contains characterizations of the three variants of general position sets. They were respectively proved in [18, Theorems 2.1, 2.3, 3.1].

Theorem 2.2 *If G is a graph and $X \subseteq V(G)$, then the following hold.*

- (i) *X is a total general position set of G if and only if $X \subseteq S(G)$. Consequently, $\text{gp}_t(G) = s(G)$.*
- (ii) *If $|X| \geq 2$, then X is an outer general position set of G if and only if each pair of vertices from X is MMD. Consequently, $\text{gp}_o(G) = \omega(G_{\text{SR}})$.*
- (iii) *If X is a general position set of G , then X is a dual general position set if and only if $G - X$ is convex in G .*

We conclude the preliminaries with the following straightforward, but very useful observation.

Lemma 2.3 *If G is a graph and $\tau \in \{\text{gp}, \text{gp}_o, \text{gp}_d, \text{gp}_t\}$, then $\tau(G) \geq s(G)$. Moreover, the equality holds if G is a block graph.*

3 Vertex removal

Before we turn our attention to vertex-deleted subgraphs in specific variants of general position sets, we consider the following example which is useful to understand each of the three invariants investigated. Let G_n , $n \geq 2$, be the graph, and x its vertex as shown in Fig. 1.




Figure 1: Graph G_n with $|X_1| = |X_2| = n$.

Proposition 3.1 *If $n \geq 2$ and $\tau \in \{\text{gp}_o, \text{gp}_d, \text{gp}_t\}$, then*

$$\tau(G_n) = 2n \quad \text{and} \quad \tau(G_n - x) = 4n.$$

Proof. Since $s(G_n) = 2n$, Theorem 2.2(i) yields $\text{gp}_t(G_n) = 2n$. Observe that two vertices of G_n are MMD if and only if they are leaves, or they both belong to X_1 , or they both belong to X_2 . Hence the strong resolving graph $(G_n)_{\text{SR}}$ has three non-trivial components, one isomorphic to K_{2n} , and the other two each isomorphic to K_n . Hence $\text{gp}_o(G_n) = 2n$ by Theorem 2.2(ii). By Lemma 2.3 we also have $\text{gp}_d(G_n) \geq 2n$. To prove the reverse inequality, we will apply Theorem 2.2(iii). For this sake we first get by a case analysis that $\text{gp}(G_n) = 2n + 1$. By symmetry, the gp-sets to be considered are $X_1 \cup X_2 \cup \{v\}$, $X_2 \cup Y \cup \{u\}$, $X_2 \cup Y \cup \{v\}$, $X_2 \cup Y \cup \{w\}$, $X_1 \cup Y \cup \{u\}$, $X_1 \cup Y \cup \{v\}$, and $X_1 \cup Y \cup \{w\}$. If Z is any of these sets, then $G_n - Z$ is not connected, hence clearly not convex. Theorem 2.2(iii) thus implies that none of these sets is a dual general position set. It follows that $\text{gp}_d(G_n) \leq 2n$ and we can conclude that $\text{gp}_d(G_n) = 2n$.

Since the graph $G_n - x$ is a tree, Lemma 2.3 gives

$$\text{gp}_t(G_n - x) = \text{gp}_o(G_n - x) = \text{gp}_d(G_n - x) = s(G_n - x) = 4n,$$

and we are done. □

3.1 Total general position sets

In this subsection we show that the total general position set of $G - x$ can be bounded below and above when x is not a cut vertex of a graph G .

Theorem 3.2 *If x is not a cut vertex of a graph G , then*

$$\text{gp}_t(G) - 1 \leq \text{gp}_t(G - x) \leq \text{gp}_t(G) + \deg_G(x).$$

Moreover, if x is a simplicial vertex, then

$$\text{gp}_t(G - x) \leq \text{gp}_t(G) + \deg_G(x) - 1.$$

In addition, all three bounds are sharp.

Proof. The bounds clearly hold if G is complete, hence assume in the rest of the proof that G is not complete.

Let X be a gp_t -set of G . Then $X = S(G)$ by Theorem 2.2(i).

To prove the lower bound, it suffices to show that $X \setminus \{x\} \subseteq S(G - x)$. Suppose to the contrary that $y \in X \setminus \{x\}$ is not simplicial in $G - x$. Then there exist $z, w \in N_{G-x}(y)$ such that z and w are not adjacent in $G - x$. The vertices z and w are also not adjacent in G , but then y is not simplicial in G , a contradiction. Hence $X \setminus \{x\} \subseteq S(G - x)$ and by applying Theorem 2.2(i) we get $\text{gp}_t(G) - 1 \leq |X \setminus \{x\}| \leq \text{gp}_t(G - x)$. This proves the lower bound.

For the upper bound observe that if $y \in V(G) \setminus N_G[x]$ is a simplicial vertex of $G - x$, then y is a simplicial vertex of G . Using Theorem 2.2(i) once more we can argue as follows:

$$\text{gp}_t(G - x) = s(G - x) \leq s(G) + \deg_G(x) = \text{gp}_t(G) + \deg_G(x).$$

This proves the upper bound in the general case. Assume now that x is a simplicial vertex. Since G is not complete, at least one vertex in $N_G(x)$ is not simplicial in $G - x$, hence

$$\text{gp}_t(G - x) = s(G - x) \leq s(G) + (\deg_G(x) - 1) = \text{gp}_t(G) + \deg_G(x) - 1.$$

To show that the lower bound is sharp, consider the stars $K_{1,n}$, $n \geq 3$. Clearly, $s(K_{1,n}) = n$ and then $s(K_{1,n} - x) = n - 1$ for any leaf x of $K_{1,n}$. To demonstrate that the upper bound is tight, consider complete bipartite graphs $K_{2,n}$, $n \geq 3$. If x is a vertex of $K_{2,n}$ of degree n , then we have $\text{gp}_t(K_{2,n}) = 0$ and $\text{gp}_t(K_{2,n} - x) = n = \text{gp}_t(K_{2,n}) + \deg_{K_{2,n}}(x)$. Finally, to show that the upper bound is sharp in the case when x is a simplicial vertex, consider the edge deleted complete graph $K_n - e$ and let x_1 and x_2 be the vertices of $K_n - e$ of degree $n - 2$. Then $S(K_n - e) = \{x, y\}$, so that $\text{gp}_t(K_n - e) = 2$. On the other hand, $(K_n - e) - x_1 \cong K_{n-1}$. Hence $\text{gp}_t((K_n - e) - x_1) = n - 1 = \text{gp}_t(K_n - e) + \deg_{K_n - e}(x_1) - 1$. \square

Note that in view of Theorem 2.2(i), we can rephrase Theorem 3.2 by saying that if x is not a cut vertex of a graph G , then

$$s(G) - 1 \leq s(G - x) \leq s(G) + \deg_G(x),$$

and if x is a simplicial vertex, then

$$s(G - x) \leq s(G) + \deg_G(x) - 1.$$

3.2 Outer general position sets

In this subsection we show that anything can happen with outer general position sets when a vertex of a graph G is removed.

To demonstrate that $\text{gp}_o(G - x)$ can be arbitrarily larger than $\text{gp}_o(G)$, consider the graph G_n , $n \geq 2$, from Fig. 1. By Proposition 3.1, $\text{gp}_o(G_n) = 2n$ and $\text{gp}_o(G_n - x) = 4n$.

To show that $\text{gp}_o(G - x)$ can also be arbitrarily smaller than $\text{gp}_o(G)$, consider the *fan graph* F_n , $n \geq 3$, of order $n + 1$, which is obtained from P_n and a vertex adjacent to all the vertices of P_n . Let x be the vertex of F_n of degree n . In [6, Proposition 3.2] it is proved that if G is a diameter 2 graph with no true twins, then $\text{gp}_o(G) = \alpha(G)$. Since $F_n - x \cong P_n$, we thus have

$$\text{gp}_o(F_n) = \lceil n/2 \rceil \quad \text{and} \quad \text{gp}_o(F_n - x) = 2.$$

On the other hand, in the case when a vertex x lies in some gp_o -set, we can bound $\text{gp}_o(G - x)$ from below as follows.

Theorem 3.3 *If x is not a cut vertex and lies in some gp_o -set of a graph G , then $\text{gp}_o(G) - 1 \leq \text{gp}_o(G - x)$ and the bound is sharp.*

Proof. Let X be a gp_o -set of G such as $x \in X$. Then $|X| = \text{gp}_o(G)$ and by Theorem 2.2(ii), every pair of vertices in X are MMD in G .

To show the bound, it suffices to prove the assertion that $X \setminus \{x\}$ is an outer general position set of $G - x$. Given Theorem 2.2(ii) this is equivalent to proving that every two vertices in $X \setminus \{x\}$ are MMD in $G - x$. Suppose to the contrary that there exist $u, v \in X \setminus \{x\}$ such that u and v are not MMD in $G - x$. By symmetry we may assume that there exists $u' \in N_{G-x}(u)$ such that $d_{G-x}(u', v) = d_{G-x}(u, v) + 1$. Since u and v are MMD in G , we have $d_G(u', v) \leq d_G(u, v)$. This implies that x lies on some shortest u', v -path in G . But then v and x are not MMD in G , a contradiction. This contradiction implies that every pair of vertices in $X \setminus \{x\}$ are MMD in $G - x$. We can conclude that $\text{gp}_o(G - x) \geq |X \setminus \{x\}| = \text{gp}_o(G) - 1$.

For sharpness of the bound, consider the stars $K_{1,n}$, $n \geq 3$. Then $\text{gp}_o(K_{1,n}) = n$ and $\text{gp}_o(K_{1,n} - x) = n - 1$ for any leaf x of $K_{1,n}$. \square

By Theorem 3.3 we can thus bound $\text{gp}_o(G - x)$ from below when x is not a cut vertex and lies in some gp_o -set of G . As a possible general upper bound on $\text{gp}_o(G - x)$ we pose:

Conjecture 3.4 *If x is not a cut vertex of a graph G , then*

$$\text{gp}_o(G - x) \leq \text{gp}_o(G) + \deg_G(x).$$

If Conjecture 3.4 holds true, then it is sharp as justified by Proposition 3.1. For the case when x is a simplicial vertex, the conjecture holds true, and even more is true:

Proposition 3.5 *If x is a simplicial vertex of a graph G , then*

$$\text{gp}_o(G - x) \leq \text{gp}_o(G) + \deg_G(x) - 1$$

and the bound is sharp.

Proof. Let u and v be arbitrary vertices of $G - x$. Since x is a simplicial vertex of G , no shortest u, v -path in G contains x which in turn implies that a u, v -path is shortest in G if and only if it is shortest in $G - x$. Consequently, if $u, v \in V(G) \setminus N_G[x]$, then u, v are MMD in G if and only if they are MMD in $G - x$.

Let $\text{gp}_o(G) = k$ and let X be a largest set of pairwise MMD vertices of G . Note that $x \in X$. Let further Y be a largest set of pairwise MMD vertices of $G - x$. Then by the above, $|Y \cap (V(G) \setminus N_G[x])| \leq k - 1$. This in turn implies that

$$|Y| \leq (k - 1) + |N_G(x)| = \text{gp}_o(G) - 1 + \deg_G(x),$$

which proves the bound.

Let $G_{n,k}$, $k < \frac{n+1}{2}$, be the graph obtained from K_n and another vertex x which is adjacent to k vertices of K_n . Then $\text{gp}_o(G_{n,k}) = \max\{n - k + 1, k\} = n - k + 1$, where the last equality holds because we have assumed that $k < \frac{n+1}{2}$. On the other hand, $G_{n,k} - x \cong K_n$ and hence $\text{gp}_o(G_{n,k} - x) = n$. \square

3.3 Dual general position sets

In this subsection we first show that $gp_d(G - x)$ can be arbitrarily larger/smaller than $gp_d(G)$. After that we prove that if x is not a cut vertex and lies in some gp_d -set of G , then $gp_d(G) - 1 \leq gp_d(G - x)$. We conclude the subsection by demonstrating that for such a vertex x , the value $gp_d(G - x)$ cannot be bounded from above by $gp_d(G)$.

Consider the fan graphs F_n ($n \geq 4$) defined in Subsection 3.2, where x denotes the vertex of degree n . It was proved in [19, Corollary 2.9] that if $n \geq 4$, then $gp(F_n) = \left\lfloor \frac{2(n+1)}{3} \right\rfloor$. Moreover, a gp-set of F_n consists of the end-vertices of a largest set of independent edges of the path P_n in F_n . By Theorem 2.2(iii) it is then straightforward to check that every largest general position set of F_n is a dual general position set. Hence $gp(F_n) = gp_d(F_n)$. Moreover, since $gp(F_n) = \left\lfloor \frac{2(n+1)}{3} \right\rfloor$ for $n \geq 4$, and because $F_n - x \cong P_n$, we have

$$gp_d(F_n) = \left\lfloor \frac{2(n+1)}{3} \right\rfloor \quad \text{and} \quad gp_d(F_n - x) = 2.$$

On the other hand, consider the family of graphs defined as follows. The *wheel* W_n , $n \geq 4$, is the graph of order $n + 1$ obtained from C_n by adding one more vertex and connecting it to all vertices of the cycle. The *mushroom* M_k , $k \geq 4$, is the graph obtained from the disjoint union of W_{k+4} and K_k by adding a matching between the vertices of K_k and k consecutive vertices of the $(k + 4)$ -cycle of W_{k+4} . See Fig. 2 for M_4 .




Figure 2: The mushroom M_4 .

Proposition 3.6 *If $k \geq 4$, then $\text{gp}_d(M_k) = k + 2$ and $\text{gp}_d(M_k - x) = 0$.*

Proof. Let $k \geq 4$, and set

$$V(M_k) = \{w_1, \dots, w_{k+4}, x, v_1, \dots, v_k\},$$

where $V(W_{k+4}) = \{w_1, \dots, w_{k+4}, x\}$ with x being the center of W_{k+4} , $V(K_k) = \{v_1, \dots, v_k\}$, and $v_i w_i \in E(M_k)$ for $i \in [k]$, see Fig. 2 again.

Let $Y = V(K_k) \cup \{w_{k+2}, w_{k+3}\}$. Then it follows from Theorem 2.1 that Y is a general position set of M_k . Moreover, $M_k - Y$ is convex, hence Theorem 2.2(iii) implies that Y is a dual general position set of M_k . Thus $\text{gp}_d(M_k) \geq k + 2$.

To prove that $\text{gp}_d(M_k) \leq k + 2$, suppose to the contrary that there exists a dual general position set X of M_k with $|X| > k + 2$. We first claim that $x \notin X$ by using the argument that if x would belong to X , then $G - X$ would not be convex. So suppose that $x \in X$. Then at least one of w_1 and w_4 must belong to X , say $w_1 \in X$. Then $w_4 \notin X$. Since $w_1 \in X$, exactly one of w_2 and w_{k+4} lies in X . If $w_{k+4} \in X$, then w_2 and w_4 are not X -positionable. And if $w_2 \in X$, then w_{k+4} and w_4 are not X -positionable. We analogously get a contradiction in the case when $w_4 \in X$. We can conclude that $x \notin X$. We next claim that $w_i \notin X$ for each $i \in [k]$. Suppose that $w_i \in X$ for some $i \in [k]$. Since $x \notin X$, we get that $v_i \in X$ because x, w_i, v_i is the unique shortest x, v_i -path. Further, exactly one of the neighbors of w_i on the $(k + 4)$ -cycle of W_{k+4} lies in X . But this is not possible, as X is a general position set.

From our assumption, we have $|X \cap \{w_{k+1}, w_{k+2}, w_{k+3}, w_{k+4}\}| \geq 3$. If $w_{k+1} \in X$, then $w_{k+2} \in X$ because $w_k \notin X$. Otherwise, the vertices w_k and w_{k+2} are not X -positionable. Since w_{k+2} lies on the shortest w_{k+1}, w_{k+3} -path, it follows that $w_{k+3} \notin X$ and $w_{k+4} \in X$. If $w_{k+4} \in X$, then we must have $w_1 \in X$ which is not possible because we have proved that $w_i \notin X$ for each $i \in [k]$. This contradiction implies that $w_{k+1} \notin X$. Similarly, we get that $w_{k+4} \notin X$. Therefore, $|X \cap \{w_{k+1}, w_{k+2}, w_{k+3}, w_{k+4}\}| \leq 2$, which leads to $|X| \leq k + 2$, the final contradiction. We can conclude that $\text{gp}_d(M_k) = k + 2$.

We now prove that $\text{gp}_d(M_k - x) = 0$. Let X be an arbitrary dual general position set of $M_k - x$. Then we first infer that $w_i \notin X$ for each $i \in [k]$. Indeed, if $w_i \in X$ for some $i \in [k]$, then exactly one of w_{i-1}, w_{i+1} must belong to X , but this leads to a contradiction that X is a dual general position set. Similarly we get that none of w_{k+j} , $j \in [4]$, belong to X . Finally, if some $v_i \in X$, then we get that all v_i , $i \in [k]$ belong to X , but then w_1 and w_k are not X -positionable. We can conclude that $X = \emptyset$. \square

Proposition 3.6 shows that $\text{gp}_d(G - x)$ can be arbitrarily smaller than $\text{gp}_d(G)$. Next, we show that $\text{gp}_d(G - x)$ can also be arbitrarily larger than $\text{gp}_d(G)$ by the

following example. Let T_k , $k \geq 3$, be a graph obtained from the disjoint union of $K_{1,2k}$ with the central vertex u and leaves v_1, \dots, v_{2k} , and an isolated vertex x , by adding the edges xv_i , $i \in [k]$.

Proposition 3.7 *If $k \geq 3$, then $\text{gp}_d(T_k) = k$ and $\text{gp}_d(T_k - x) = 2k$.*

Proof. Since $s(T_k) = k$, Lemma 2.3 gives $\text{gp}_d(T_k) \geq k$. To prove the reverse inequality, consider an arbitrary gp_d -set X of T_k . Suppose first that $x \in X$. Then $v_i, v_j \in X$, where $i, j \in [k]$, is not possible since then v_i and v_j are not X -positionable. Analogously we see that also $v_i, v_j \notin X$, where $i, j \in [k]$, cannot happen. Since $k \geq 3$, we can conclude that $x \notin X$. By an analogous reason, $u \notin X$. This in turn implies that $v_i \notin X$ for each $i \in [k]$. Hence $|X| \leq k$, and then $\text{gp}_d(T_k) = k$.

Since $s(T_k - x) = 2k$, Lemma 2.3 implies that $\text{gp}_d(T_k - x) = 2k$, and we are done. \square

We also have a result parallel to Theorem 3.3 for the dual general position number in the following.

Theorem 3.8 *If x is not a cut vertex and lies in some gp_d -set of a graph G , then $\text{gp}_d(G) - 1 \leq \text{gp}_d(G - x)$ and the bound is sharp.*

Proof. Let X be a gp_d -set of G such that $x \in X$. Then $|X| = \text{gp}_d(G)$ and by Theorem 2.2(ii), $G - X$ is convex. By the proof of [7, Proposition 3.3], the set $X \setminus \{x\}$ is a general position set of $G - x$. It suffices to show that $(G - x) - (X \setminus \{x\})$ is convex in $G - x$, because then by Theorem 2.2(iii) the set $X \setminus \{x\}$ is a dual general position set of $G - x$ and hence $\text{gp}_d(G - x) \geq |X \setminus \{x\}| = \text{gp}_d(G) - 1$.

Suppose to the contrary that this is not the case. There are $u, v \notin X$ such that a shortest u, v -path P contains a vertex $w \in X \setminus \{x\}$. Since $G - X$ is convex, $d_G(u, v) < d_G(u, w) + d_G(w, v)$. Based on this, we can conclude the following:

$$\begin{aligned} d_{G-x}(u, v) &= d_{G-x}(u, w) + d_{G-x}(w, v) \\ &\geq d_G(u, w) + d_G(w, v) \\ &> d_G(u, v). \end{aligned}$$

This in turn implies that x lies on the shortest u, v -path in G which contradicts with $G - X$ being convex. We can conclude that $(G - x) - (X \setminus \{x\})$ is convex in $G - x$.

To prove the sharpness of the bound, consider a graph $K_{1,n}$, $n \geq 3$, and let x be an arbitrary leaf of it. Then $\text{gp}_d(K_{1,n}) = n$ and $\text{gp}_d(K_{1,n} - x) = n - 1$. \square

In Theorem 3.8 we have bounded $\text{gp}_d(G - x)$ from below by $\text{gp}_d(G) - 1$ when x is not a cut vertex and lies in some gp_d -set of G . We next demonstrate that for such a vertex x the value $\text{gp}_d(G - x)$ cannot be bounded from above by $\text{gp}_d(G)$.

Consider the graph Y_k , which is obtained from the complete graph K_{k+3} by adding two vertices x and y , and connecting each of them to the same k vertices of K_{k+3} as illustrated in Fig. 3.




Figure 3: Graph Y_k .

Proposition 3.9 *If $k \geq 2$, then $\text{gp}_d(Y_k) = 5$ and $\text{gp}_d(Y_k - x) = k + 3$.*

Proof. Let $\{v_1, \dots, v_k, u, v, w\}$ be the set of vertices of the induced complete graph K_{k+3} in Y_k as indicated in Fig. 3. Then $V(Y_k) = \{v_1, \dots, v_k, u, v, w, x, y\}$ and $\deg_{Y_k}(u) = \deg_{Y_k}(v) = \deg_{Y_k}(w) = k + 2$. It is straightforward to check that $\{x, y, u, v, w\}$ is a general position set of Y_k . Hence $\{x, y, u, v, w\}$ is also a dual general position set of Y_k by Theorem 2.2(iii). Therefore, $\text{gp}_d(Y_k) \geq 5$.

To prove that $\text{gp}_d(Y_k) \leq 5$, consider an arbitrary gp_d -set X of Y_k . Suppose to the contrary that $v_i \in X$ for some $i \in [k]$. Then exactly one of the vertices x and y must belong to X , for otherwise v_i would lie on a shortest x, y -path, which is a contradiction. Without loss of generality assume that $x \in X$ and $y \notin X$. We claim that there is no vertex from $\{u, v, w\}$ which is contained in X . If $z \in \{u, v, w\} \cap X$, then v_i lies on a shortest x, z -path, but this contradicts the fact that X is a gp_d -set of Y_k . If a vertex of $\{u, v, w\}$ does not contain in X , say u , then v_i lies on a shortest y, u -path, which implies that y and u are not X -positionable. This contradiction yields $v_i \notin X$ for each $i \in [k]$. Therefore, $|X| \leq 5$, and we conclude that $\text{gp}_d(Y_k) = 5$.

If x is removed from Y_k , by Theorem 2.1, the set $\{v_1, \dots, v_k, u, v, w\}$ is a largest general position set of $Y_k - x$. Using Theorem 2.2(iii), we get that $\{v_1, \dots, v_k, u, v, w\}$ is also a dual general position set of $Y_k - x$. Note that $\text{gp}_d(Y_k - x) \leq \text{gp}(Y_k - x) = k + 3$. We can conclude that $\text{gp}_d(Y_k - x) = k + 3$. \square

4 Edge removal

Impact of removing an edge on the general position number of a graph was investigated in [7]. In this section, we complement this by examining the effect of edge removal on the other three general position invariants.

We begin with the total general position number, for which the following notation is needed. If G is a graph and $e = uv \in E(G)$, then let $S(G)_e$ denote the set of simplicial vertices in G which are adjacent to both u and v .

Proposition 4.1 *If e is an edge of a graph G , then*

$$\text{gp}_t(G) - |S(G)_e| \leq \text{gp}_t(G - e) \leq \text{gp}_t(G) + 2.$$

Moreover, the bounds are sharp.

Proof. Let $e = uv$. If x is a simplicial vertex which is adjacent to at most one of u and v , then x remains simplicial in $G - e$. Thus $\text{gp}_t(G - e) = s(G - e) \geq s(G) - |S(G)_e| = \text{gp}_t(G) - |S(G)_e|$. If $x \neq u, v$ is a simplicial vertex of $G - e$, it remains simplicial in G . Hence $\text{gp}_t(G) \geq \text{gp}_t(G - e) - 2$.

For the sharpness, note first that $s(K_n) = n$ and $s(K_n - e) = 2$, hence the lower bound is sharp. For the upper bound, let X_n , $n \geq 2$, be the graph obtained from the disjoint union of two copies of K_n , say K and K' and a vertex x , by adding an edge $e = uv$ between K and K' , and joining x to a vertex u' in K and a vertex v' in K' , where $\{u, v\} \cap \{u', v'\} = \emptyset$. Then we infer that $\text{gp}_t(X_n) = 2(n - 2)$ and $\text{gp}_t(X_n - e) = 2(n - 1)$. \square

For the outer general position number, the following holds.

Theorem 4.2 *If e is an edge in graph G , then*

$$\frac{\text{gp}_o(G)}{2} \leq \text{gp}_o(G - e) \leq 2\text{gp}_o(G).$$

Moreover, both bounds are sharp.

Proof. Let $e = uv$ and let X be a gp_o -set of G . Then consider the sets of vertices

$$\begin{aligned} X_{uv} &= \{w \in X : d_G(u, w) < d_G(v, w)\}, \\ X_{vu} &= \{w \in X : d_G(v, w) < d_G(u, w)\}, \\ {}_v X_u &= \{w \in X : d_G(u, w) = d_G(v, w)\}. \end{aligned}$$

Clearly, $X = X_{uv} \cup X_{vu} \cup {}_v X_u$. Let

$$X_u = X_{uv} \cup {}_v X_u \quad \text{and} \quad X_v = X_{vu} \cup {}_v X_u.$$

Then we recall from the proof of [7, Theorem 6.2] that X_u and X_v are general position sets of $G - e$. Moreover, by a parallel argument we prove that X_u and X_v remain to be MMD in $G - e$ provided they were MMD in G . Since $|X|/2 \leq \max\{|X_u|, |X_v|\}$, Theorem 2.2(ii) yields the lower bound. The argument for the upper bounds proceeds along parallel lines.

To demonstrate the sharpness of the lower bound, consider first graphs Y'_n , $n \geq 3$, constructed as follows. First, take two disjoint copies of K_n , say K and K' and add an edge between them, say $e = uv$, where $u \in K$ and $v \in K'$. Second, add two new vertices u' and v' , the edge $u'v'$, all the edges between u' and $V(K) \setminus \{u\}$, and all the edges between v' and $V(K') \setminus \{v\}$, see Fig. 4.




Figure 4: Graphs Y'_4 and $Y'_4 - e$, and their gp_o -sets

We can observe that $gp_o(Y'_n) = 2(n - 1)$ and that $gp_o(Y'_n - e) = n - 1$. This demonstrates the sharpness of the lower bound.

For the sharpness of the upper bound, let Z_n , $n \geq 2$, be the graph obtained from two disjoint copies of $K_{2,n}$ by adding an edge $e = uv$ between them, where u and v are vertices of degree n in different copies of $K_{2,n}$. Then $gp_o(Z_n) = n$ and $gp_o(Z_n - e) = 2n$. \square

The sharpness of the upper bound in Theorem 4.2 is demonstrated by graphs that contain bridges. Consider next the graphs Z_n , $n \geq 3$, constructed as follows. Take two disjoint complete graphs K_n , add an edge e between them, and add a path of length 3 between them disjoint from e . Then we can infer that $gp_o(Z_n) = 2n - 4$ and that $gp_o(Z_n - e) = 2n - 2$. In the case $n = 3$, we thus also have a graph without bridges for which the upper bound in Theorem 4.2 is sharp. It remains to be seen

whether the upper bound in Theorem 4.2 is sharp also on graphs without bridges with an arbitrary large other general position number.

To conclude the paper, we consider the dual general position number under edge removal for which we have the following.

Theorem 4.3 *The difference $\text{gp}_d(G) - \text{gp}_d(G - e)$ can be arbitrarily large.*

Proof. Consider the graphs H_n , $n \geq 1$, obtained from n disjoint triangles and one C_6 , by selecting an edge in each of these $n + 1$ graphs and identifying them into a single edge. See Fig. 5 from which the vertex labelling of H_n should be clear.




Figure 5: The graph H_n

Let $X = \{w_1, \dots, w_n\}$. It is straightforward to check that X is a general position set of H_n . Since there is a unique shortest path between x and y that does not contain any vertex from X , it follows that $G - X$ is convex. In addition, since $\{x, y, v_1, v_2, v_3, v_4\}$ induces a convex C_6 , using Theorem 2.2(iii) we infer that none of these vertices lies in a dual general position set. We can conclude that X is the largest dual general position set of H_n , therefore $\text{gp}_d(H_n) = n$. On the other hand, removing the edge $e = xy$, every edge of $H_n - e$ is the middle edge of some isometric path P_4 . In view of [18, Proposition 3.3] we have $\text{gp}_d(H_n - e) = 0$. \square

Acknowledgements

This work was supported by the Slovenian Research and Innovation Agency (ARIS) under the grants P1-0297, N1-0355, N1-0285, N1-0431, and J1-70045.

Data Availability Statement

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

Conflict of Interest Statement

Sandi Klavžar is an Associate Editor of the Discrete Applied Mathematics journal and was not involved in the review and decision-making process of this article. In addition, the authors declare no other conflict of interest.

References

- [1] B.S. Anand, U. Chandran S.V., M. Changat, S. Klavžar, E.J. Thomas, Characterization of general position sets and its applications to cographs and bipartite graphs, *Appl. Math. Comput.* 359 (2019) 84–89.
- [2] J. Araujo, M.C. Dourado, F. Protti, R. Sampaio, The iteration time and the general position number in graph convexities, *Appl. Math. Comput.* 487 (2025) Paper 129084.
- [3] U. Chandran S.V., S. Klavžar, J. Tuite, The general position problem: a survey, [arXiv:2501.19385](https://arxiv.org/abs/2501.19385) (2025).
- [4] U. Chandran S.V., G.J. Parthasarathy, The geodesic irredundant sets in graphs, *Int. J. Math. Combin.* 4 (2016) 135–143.
- [5] G. Di Stefano, S. Klavžar, A. Krishnakumar, J. Tuite, I.G. Yero, Lower general position sets in graphs, *Discuss. Math. Graph Theory* 45 (2025) 509–531.
- [6] P. Dokyeesun, S. Klavžar, D. Kuziak, J. Tian, General position problems in strong and lexicographic products of graphs, *Comput. Appl. Math.* 45 (2026) Paper 97.

- [7] P. Dokyeesun, S. Klavžar, J. Tian, The general position number under vertex and edge removal, *Quaest. Math.* 48 (2025) 1277–1290.
- [8] V. Iršič, S. Klavžar, G. Rus, J. Tuite, General position polynomials, *Results Math.* 79 (2024) Paper 110.
- [9] S. Klavžar, A. Krishnakumar, D. Kuziak, E. Shallcross, J. Tuite, I.G. Yero, Moving through Cartesian products, coronas and joins in general position, *Discrete Appl. Math.* 379 (2026) 768–780.
- [10] J. Körner, On the extremal combinatorics of the Hamming space, *J. Comb. Theory Ser. A* 71 (1995) 112–126.
- [11] D. Korže, A. Vesel, Mutual-visibility and general position sets in Sierpiński triangle graphs, *Bull. Malays. Math. Sci. Soc.* 48 (2025) Paper 106.
- [12] E. Kruft Welton, S. Khudairi, J. Tuite, Lower general position in Cartesian products, *Commun. Comb. Optim.* 10 (2025) 110–125.
- [13] P. Manuel, S. Klavžar, A general position problem in graph theory, *Bull. Aust. Math. Sci. Soc.* 98 (2018) 177–187.
- [14] O.R. Oellermann, J. Peters-Fransen, The strong metric dimension of graphs and digraphs, *Discrete Appl. Math.* 155 (2007) 356–364.
- [15] D. Roy, S. Klavžar, A.S. Lakshmanan, Mutual-visibility and general position in double graphs and in Mycielskians, *Appl. Math. Comput.* 488 (2025) Paper 129131.
- [16] M. Thankachy, U. Chandran S.V., J. Tuite, E. Thomas, G. Di Stefano, G. Erskine, On the vertex position number of graphs, *Discuss. Math. Graph Theory* 44 (2024) 1169–1188.
- [17] E.J. Thomas, U. Chandran S.V., J. Tuite, G. Di Stefano, On the general position number of Mycielskian graphs, *Discrete Appl. Math.* 353 (2024) 29–43.
- [18] J. Tian, S. Klavžar, Variety of general position problems in graphs, *Bull. Malays. Math. Sci. Soc.* 48 (2025) Paper 5.
- [19] J. Tian, K. Xu, D. Chao, On the general position numbers of maximal outerplane graphs, *Bull. Malays. Math. Sci. Soc.* 46 (2023) Paper 198.
- [20] E.K. Welton, S. Khudairi, J. Tuite, Lower general position in Cartesian products, *Commun. Comb. Optim.* 10 (2025) 110–125.