

## ON GRAPHS WITH SMALL GAME DOMINATION NUMBER

*Sandi Klavžar, Gašper Košmrlj, Simon Schmidt*

The domination game is played on a graph  $G$  by Dominator and Staller. The game domination number  $\gamma_g(G)$  of  $G$  is the number of moves played when Dominator starts and both players play optimally. Similarly,  $\gamma'_g(G)$  is the number of moves played when Staller starts. Graphs  $G$  with  $\gamma_g(G) = 2$ , graphs with  $\gamma'_g(G) = 2$ , as well as graphs extremal with respect to the diameter among these graphs are characterized. In particular,  $\gamma'_g(G) = 2$  and  $\text{diam}(G) = 3$  hold for a graph  $G$  if and only if  $G$  is a so-called gamburger. Graphs  $G$  with  $\gamma_g(G) = 3$  and  $\text{diam}(G) = 6$ , as well as graphs  $G$  with  $\gamma'_g(G) = 3$  and  $\text{diam}(G) = 5$  are also characterized.

### 1. INTRODUCTION

The domination game is played on an arbitrary graph  $G$  by *Dominator* and *Staller*. The two players take turns choosing a vertex from  $G$  such that at least one previously undominated vertex becomes dominated. The game ends when no move is possible and the score of the game is the total number of vertices chosen. Dominator wants to minimize the score, while Staller wants to maximize it. By *D-game* we mean a game in which Dominator has the first move and by *S-game* a game started by Staller. Assuming that both players play optimally, the *game domination number*  $\gamma_g(G)$  of a graph  $G$  denotes the score of D-game played on  $G$ . Similarly, the *Staller-start game domination number*  $\gamma'_g(G)$  is defined as the score of optimal S-game.

---

2010 Mathematics Subject Classification. 05C57, 05C69, 05C12.

Keywords and Phrases. Domination game, game domination number, diameter, gamburger.

The game was introduced in [4] and already received considerable attention. One of the reasons for this interest is the so-called 3/5-conjecture from [18] asserting that  $\gamma_g(G) \leq 3n/5$  holds for any isolate free graph of order  $n$ . Trees that attain this bound were investigated in [5], while recently BUJTAS [8] made a breakthrough by proving that the conjecture holds for all graphs with the minimum degree at least 3. In order to achieve this result she further developed the proof technique introduced in [6, 7] where the conjecture is verified for all trees in which no two leaves are at distance 4. HENNING and KINNERSLEY [13] further established the truth of the 3/5-conjecture over the class of graphs with minimum degree at least 2, hence the 3/5-conjecture remains open only for graphs with pendant vertices. We point out that Bujtas' proof technique already turned out to be useful elsewhere [9].

Recently, two closely related games were introduced. The total version of the domination game was investigated in [14] and further studied in [15] where it was proved that for any graph of order  $n$  in which every component contains at least three vertices, the corresponding total invariant is bounded above by  $4n/5$ , see also [16]. The second related game, named the *disjoint domination game*, was studied in [11]. Among the additional investigations of the domination game we mention the complexity studies from [1], the behaviour of the game played on the disjoint union of graphs [12], domination game critical graphs [10], and a characterization of trees  $T$  for which  $\gamma_g(T) = \gamma(T)$  holds [20], where  $\gamma(T)$  is the usual domination number of  $T$ .

Motivated in part by the characterization of graphs with  $\gamma_g = 3$  and  $\gamma'_g = 2$  from [2] and by the complexity studies from [1], we study in this paper graphs that have small game domination number. In the next section we introduce notations needed, recall some results, and bound the diameter of a graph from above in terms of the game domination number (see [3, Corollary 4.1] for a closely related result). In Section 3 we first characterize graphs  $G$  with  $\gamma_g(G) = 2$  and graphs with  $\gamma'_g(G) = 2$ . We also characterize graphs extremal with respect to the diameter among these graphs. In particular, we introduce the concept of the so-called gamburger and prove that  $G$  has  $\gamma'_g(G) = 2$  and  $\text{diam}(G) = 3$  if and only if  $G$  is a gamburger. In Section 4 we then characterize extremal graphs (w.r.t. the diameter) in the class of graphs with the game domination number equal 3. In particular,  $\gamma'_g(G) = 3$  and  $\text{diam}(G) = 5$  hold for a graph  $G$  if and only if  $G$  is the so-called double-gamburger.

## 2. PRELIMINARIES

If  $x$  is a vertex of a graph  $G$ , then  $N_G(x)$  (resp.  $N_G[x]$ ) denotes the neighborhood (resp. closed neighborhood) of  $x$ . Let  $d_G(x, y)$  be the standard shortest-path distance between vertices  $x$  and  $y$  of  $G$ . The eccentricity  $\text{ecc}_G(x)$  of  $x$  is  $\max\{d(x, y) : y \in V(G)\}$  and the diameter  $\text{diam}(G)$  of  $G$  is the maximum eccentricity of its vertices. If  $\text{ecc}_G(x) = \text{diam}(G)$ , then  $x$  is called a diametrical vertex. A pair of vertices  $x, y$  such that  $d_G(x, y) = \text{diam}(G)$  holds is called a diametrical pair.  $S_r^G(x) = \{y \in V(G) : d_G(x, y) = r\}$  is called the sphere with center  $x$  and radius  $r$  and  $B_r^G(x) = \{y \in V(G) : d_G(x, y) \leq r\}$  is called the ball with center  $x$  and radius

$r$ . If  $G$  will be clear from the context, we will simplify the above notations to  $N(x)$ ,  $N[x]$ ,  $d(x, y)$ ,  $S_r(x)$ ,  $B_r(x)$ , and  $\text{ecc}(x)$ .

We next collect some known results to be used later on. A fundamental result about the domination game is the following theorem for which the fact that  $\gamma_g(G) \leq \gamma'_g(G) + 1$  holds was proved in [4], while the inequality  $\gamma'_g(G) \leq \gamma_g(G) + 1$  was later established in [18].

**Theorem 2.1.** [4, 18] *For any graph  $G$ ,  $|\gamma_g(G) - \gamma'_g(G)| \leq 1$ .*

If  $(\gamma_g(G), \gamma'_g(G)) = (k, \ell)$  then we say that  $G$  *realizes*  $(k, \ell)$ . By Theorem 2.1, if  $G$  realizes the pair  $(k, \ell)$ , then  $|k - \ell| \leq 1$ .

Vertices  $u$  and  $v$  of a graph  $G$  are called *twins* if  $N[u] = N[v]$ . Note that twins are necessarily adjacent. A graph is called *twin-free* if it contains no twins. The following result is not difficult to prove and was implicitly or explicitly (cf. [2]) used earlier and is also stated in [20].

**Proposition 2.2.** *If  $u$  and  $v$  are twins in a graph  $G$ , then  $\gamma_g(G) = \gamma_g(G - u)$  and  $\gamma'_g(G) = \gamma'_g(G - u)$ .*

A subgraph  $H$  of a graph  $G$  is *guarded in  $G$*  if for any vertex  $x$  in  $G$  there exists a vertex  $y \in V(H)$  such that  $N[x] \cap V(H) \subseteq N[y] \cap V(H)$ . The vertex  $y$  is called a *guard* of  $x$  in  $H$ . (If  $x \in V(H)$ , then  $x$  is a guard of itself.) The concept of a guarded subgraph was introduced in [3] where the following result was proved:

**Theorem 2.3.** *If  $H$  is guarded in  $G$ , then  $\gamma_g(H) \leq \gamma_g(G)$  and  $\gamma'_g(H) \leq \gamma'_g(G)$ .*

To bound the diameter of a graph with its game domination number we recall the following result proved in [17], cf. also [19].

**Proposition 2.4.** *If  $n \geq 1$ , then*

$$(i) \quad \gamma_g(P_n) = \begin{cases} \left\lceil \frac{n}{2} \right\rceil - 1; & n \equiv 3 \pmod{4}, \\ \left\lceil \frac{n}{2} \right\rceil; & \text{otherwise.} \end{cases}$$

$$(ii) \quad \gamma'_g(P_n) = \left\lceil \frac{n}{2} \right\rceil.$$

We are now ready for the announced bound that was not earlier reported (at least explicitly) in the literature, hence we include its proof.

**Proposition 2.5.** *If  $G$  is a graph, then*

$$(i) \quad \text{diam}(G) \leq \begin{cases} 2\gamma_g(G); & \gamma_g(G) \text{ odd}, \\ 2\gamma_g(G) - 1; & \text{otherwise.} \end{cases}$$

$$(ii) \quad \text{diam}(G) \leq 2\gamma'_g(G) - 1.$$

*Moreover, the bounds are tight.*

**Proof.** Let  $G$  be a graph and let  $P$  be a diametrical path of  $G$ . Then  $P$  is a shortest path isomorphic to  $P_{\text{diam}(G)+1}$  and by the proof of [3, Corollary 4.1],  $P$  is guarded in  $G$ . Hence by Theorem 2.3,  $\gamma_g(G) \geq \gamma_g(P)$  and  $\gamma'_g(G) \geq \gamma'_g(P)$ . Using the latter inequality and Proposition 2.4 we get

$$\gamma'_g(G) \geq \gamma'_g(P_{\text{diam}(G)+1}) = \left\lceil \frac{\text{diam}(G)+1}{2} \right\rceil \geq \frac{\text{diam}(G)+1}{2},$$

which proves the assertion (ii).

Let  $\gamma_g(G)$  be odd. Assume by way of contradiction that  $\text{diam}(G) \geq 2\gamma_g(G) + 1$ . Then  $G$  contains a shortest path  $P'$  of order  $2\gamma_g(G) + 2$  and hence  $\gamma_g(G) \geq \gamma_g(P') = \lceil (2\gamma_g(G) + 2)/2 \rceil = \gamma_g(G) + 1$ , a contradiction. Hence  $\text{diam}(G) \leq 2\gamma_g(G)$  holds when  $\gamma_g(G)$  is odd.

Let  $\gamma_g(G)$  be even and suppose that  $\text{diam}(G) \geq 2\gamma_g(G)$  holds. Then  $G$  contains a shortest path  $P''$  of order  $2\gamma_g(G) + 1$ . Since  $\gamma_g(G)$  is even,  $2\gamma_g(G) + 1 \equiv 1 \pmod{4}$ , and therefore,  $\gamma_g(G) \geq \gamma_g(P'') = \lceil (2\gamma_g(G) + 1)/2 \rceil = \gamma_g(G) + 1$ , a contradiction. We conclude that  $\text{diam}(G) \leq 2\gamma_g(G) - 1$  holds when  $\gamma_g(G)$  is even.

That the bounds are tight follows by considering the path graphs. More precisely, for any  $k \geq 1$  we have

- $\gamma_g(P_{4k}) = 2k$  and  $\text{diam}(P_{4k}) = 4k - 1 = 2\gamma_g(P_{4k}) - 1$ ,
- $\gamma_g(P_{4k+3}) = 2k + 1$  and  $\text{diam}(P_{4k+3}) = 4k + 2 = 2\gamma_g(P_{4k+3})$ , and
- $\gamma'_g(P_{2k}) = k$  and  $\text{diam}(P_{2k}) = 2k - 1 = 2\gamma'_g(P_{2k}) - 1$ . □

### 3. GRAPHS $G$ WITH $\gamma_g(G) = 2$ OR $\gamma'_g(G) = 2$

In this section we study the structure of graphs  $G$  with  $\gamma_g(G) = 2$  or  $\gamma'_g(G) = 2$ . Before doing it, we observe that

- $\gamma_g(G) = 1$  if and only if  $\Delta(G) = n - 1$ , and
- $\gamma'_g(G) = 1$  if and only if  $G$  is a complete graph.

Note also that if  $\gamma_g(G) = 2$  and  $G$  is not connected, then  $G$  consists of two components, one containing a universal vertex and the other being complete. Moreover, if  $\gamma'_g(G) = 2$  and  $G$  is not connected, then  $G$  consists of two complete components. To characterize connected graphs with  $\gamma_g(G) = 2$  we introduce the following concept. We say that a vertex  $u$  of a (connected) graph  $G$  is *2-dense*, if

- $\text{ecc}(u) = 2$ ,
- there is a join between  $S_2(u)$  and the neighborhood of  $S_2(u)$  in  $S_1(u)$ , and
- $S_2(u)$  induces a clique.

Let  $u$  be a 2-dense vertex of  $G$  and let  $L(u)$  be the neighbors of  $u$  that are at distance 3 from  $S_2(u)$ , see Fig. 1. We point out that it is possible that  $L(u) = \emptyset$ .

**Proposition 3.6.** *If  $G$  is a connected graph, then the following hold.*

- (i)  $\gamma_g(G) = 2$  if and only if  $G$  contains a 2-dense vertex.
- (ii)  $\gamma'_g(G) = 2$  if and only if  $G$  is not complete and every vertex lies in a dominating set of order 2.




Figure 1. A 2-dense vertex  $u$

**Proof.** (i) Suppose first that  $\gamma_g(G) = 2$  and let  $u$  be an optimal first move of Dominator. Then  $\text{ecc}(u) \geq 2$ , for otherwise Dominator could play a universal vertex and

finish the game in one move. Since Staller, by playing a vertex in  $S_1(u)$ , dominates no vertex in  $S_3(u)$  it follows that  $\text{ecc}(u) = 2$ . If there exists two nonadjacent vertices in  $S_2(u)$ , then Staller could play any one of them in order to prolong the game for one more move. Hence  $S_2(u)$  must induce a clique. Assume finally that there is a vertex  $x \in S_1(u)$  that is adjacent to  $y \in S_2(u)$  and not adjacent to  $z \in S_2(u)$ . Then Staller can play  $x$  in order to prolong the game for one more move. Therefore there is no such vertex  $z$ , that is, there is a join between  $S_2(u)$  and the neighborhood of  $S_2(u)$  in  $S_1(u)$ . We conclude that  $u$  is a 2-dense vertex.

Conversely, suppose that  $G$  contains a 2-dense vertex  $u$ . Then it is straightforward to see that Dominator forces Staller to finish the game in the next move by playing  $u$ . Hence  $\gamma_g(G) \leq 2$ . On the other hand,  $\text{ecc}(u) = 2$  implies that  $\text{diam}(G) \geq 2$  which in turn implies that  $\gamma_g(G) \geq 2$ .

(ii) Suppose that  $\gamma'_g(G) = 2$ . Then  $G$  is clearly not complete. Let next  $u$  be an arbitrary vertex of  $G$ . There is nothing to prove if  $u$  is a universal vertex (just add any vertex to  $u$  to form a dominating set). If  $u$  is not universal, then if Staller plays  $u$  as the first move, Dominator has a reply  $v$  such that the game is over after this move. But then  $\{u, v\}$  is a dominating set.

Conversely, suppose that  $G$  is not complete and every vertex lies in a dominating set of order 2. Then  $\gamma'_g(G) \geq 2$  because  $G$  is not complete. Moreover,  $\gamma'_g(G) \leq 2$  because after an arbitrary first move  $u$  of Staller, Dominator can play  $v$ , where  $\{u, v\}$  is a dominating set.  $\square$

If  $\gamma_g(G) = 2$  or  $\gamma'_g(G) = 2$ , then  $G$  realizes one of the pairs  $(1, 2)$ ,  $(2, 2)$ ,  $(3, 2)$ , or  $(2, 3)$ . Among the twin-free graphs, the classes of graphs that characterize the first three pairs can be described in the following simple way.

**Observation 3.7.** *Let  $G$  be a connected, twin-free graph with  $\gamma'_g(G) = 2$ . Then*

- (i)  $G$  realizes  $(1, 2)$  if and only if  $\Delta(G) = n - 1$ .
- (ii)  $G$  realizes  $(2, 2)$  if and only if  $\Delta(G) = n - 2$ .

(iii)  $G$  realizes  $(3, 2)$  if and only if  $\Delta(G) \leq n - 3$ .

**Proof.** The statement (i) is clear, (iii) follows from [2, Corollary 3.2], and then (ii) characterizes the remaining connected, twin-free graphs  $G$  with  $\gamma'_g(G) = 2$ .  $\square$

The last pair to consider is  $(2, 3)$ . It is straightforward to see that  $G$  realizes the pair  $(2, 3)$  if and only if  $\gamma_g(G) = 2$  and  $G$  contains a vertex that is in no minimum dominating set.

By Proposition 2.5, if  $\gamma_g(G) = 2$  or  $\gamma'_g(G) = 2$ , then  $\text{diam}(G) \leq 3$ . In the rest of the section we give an explicit description of the structure of related extremal classes of graphs, that is, those with diameter equal to 3. For  $\gamma_g$  Proposition 3.6 immediately implies:

**Corollary 3.8.** *Let  $G$  be a connected graph. Then  $\gamma_g(G) = 2$  and  $\text{diam}(G) = 3$  if and only if  $G$  contains a 2-dense vertex  $u$  such that  $L(u) \neq \emptyset$ .*

In order to characterize graphs with  $\gamma'_g(G) = 2$  and  $\text{diam}(G) = 3$ , we introduce the following concept. We say that a connected graph  $G$  is a *gamburger* with a gamburger structure  $Q_1, T_1, T_2, Q_2$ , if  $G$  is the disjoint union of non-empty subgraphs  $Q_1, T_1, T_2$ , and  $Q_2$ , where  $Q_1$  and  $Q_2$  induce cliques with no edges between them and, in addition, the following hold for any  $i \in \{1, 2\}$  (see Fig. 2).

- There is a join between  $Q_i$  and  $T_i$  and there are no edges between  $Q_i$  and  $T_{3-i}$ .
- For any vertex  $x \in T_i$  there exists a vertex  $x' \in T_{3-i} \cup Q_{3-i}$  such that  $T_1 \cup T_2 \subseteq N[x] \cup N[x']$ .




Figure 2. Gamburger (with cliques  $Q_1$  and  $Q_2$ )

We first show that every gamburger can be presented in a canonical form.

**Lemma 3.9.** *If  $G$  is a gamburger with a gamburger structure  $Q_1, T_1, T_2, Q_2$ , then there exists a gamburger structure  $Q'_1, T'_1, T'_2, Q'_2$  for  $G$  such that any vertex from  $T'_i, 1 \leq i \leq 2$ , has at least one neighbor in  $T'_{3-i}$ .*

**Proof.** Let  $Q_1, Q_2, T_1, T_2$  be a gamburger structure for  $G$ . For  $i \in \{1, 2\}$ , we define  $W_i$  as the subset of vertices of  $T_i$  whose neighborhood is contained in  $T_i \cup Q_i$ . We put  $Q'_i = Q_i \cup W_i$  and  $T'_i = T_i \setminus W_i$ . Note that, since  $G$  is connected  $W_i \neq T_i$  and then  $T'_i$  is not empty. To show that this defines a new gamburger structure for  $G$ , we only need to show that  $N[w] = Q_i \cup T_i$  holds for all  $w \in W_i$ . Suppose, this is not the case. Since there is a join between  $Q_i$  and  $T_i$ , this means that there exists  $x \in T_i$ , which is not a neighbor of  $w$ . Since  $G$  is a gamburger, by the definition

there exists a vertex  $x' \in G \setminus (T_i \cup Q_i)$  such that  $w$  belongs to  $N(x')$ . This is a contradiction with the definition of  $W_i$ . In conclusion,  $N[w] = Q_i \cup T_i$  and  $Q'_i$  induces a clique.  $\square$

**Lemma 3.10.** *A gamburger has diameter 3.*

**Proof.** Since  $Q_1$  and  $Q_2$  are not empty,  $\text{diam}(G) \geq 3$ . By Lemma 3.9 we can assume that every vertex in  $T_1$  (resp.  $T_2$ ) has a neighbor in  $T_2$  (resp.  $T_1$ ) which in turn implies  $\text{diam}(G) \leq 3$ .  $\square$

We can now prove that gamburgers are precisely the graphs  $G$  with  $\gamma'_g(G) = 2$  and extremal diameter.

**Theorem 3.11.** *A graph  $G$  has  $\gamma'_g(G) = 2$  and  $\text{diam}(G) = 3$  if and only if  $G$  is a gamburger.*

**Proof.** Assume first that  $G$  is a gamburger. By Lemma 3.10,  $\text{diam}(G) = 3$ . We prove that  $\gamma'_g(G)$  is 2. Since  $\text{diam}(G) = 3$ ,  $\gamma'_g(G)$  is clearly greater or equal to 2. First, suppose Staller selects a vertex in one of the two cliques, say  $Q_1$ . Dominator ends the game at his turn, by playing any vertex in  $Q_2$ . Suppose next that Staller does not play in one of the two cliques, but say on  $x \in T_1$ . There is a vertex  $x' \in T_2 \cup Q_2$  such that  $N[x] \cup N[x']$  contains  $T_1 \cup T_2$ . Since no vertex in  $Q_2$  is dominated, playing  $x'$  is a legal move, which enables Dominator to end the game at his turn.

Conversely, suppose that  $\gamma'_g(G) = 2$  and  $\text{diam}(G) = 3$ . Let  $u$  be an arbitrary diametrical vertex of  $G$ . We set  $Q_1 = \{u\}$ ,  $T_1 = S_1(u)$ ,  $T_2 = S_2(u)$  and  $Q_2 = S_3(u)$ . These four sets obviously form a partition of  $V(G)$ . We claim that  $Q_1, T_1, T_2, Q_2$  is a gamburger structure for  $G$ . It is clear that there are no edges that would violate the gamburger structure, that is, there are no edges between  $Q_1$  and  $Q_2$  and no edges between  $Q_i$  and  $T_{3-i}$  for  $1 \leq i \leq 2$ . It is also clear that  $Q_1$  induces a (one-vertex) clique and hence that there is a join between  $Q_1$  and  $T_1$ . Whatever Staller plays as her first move, Dominator has to be able to end the game in the next turn. If Staller plays in  $Q_2$ , Dominator will not be able at his turn to simultaneously dominate vertices in  $Q_1$  and  $Q_2$ . Hence, all the vertices of  $Q_2$  must be dominated by such a Staller's first move. It follows that  $Q_2$  induces a clique. Assume next that Staller plays a vertex  $y \in T_2$ . In order to dominate the vertex in  $Q_1$ , Dominator has to play in  $T_1 \cup Q_1$ . Hence, all the vertices of  $Q_2$  are adjacent to  $y$ . Since  $y$  was an arbitrary vertex of  $T_2$ , it follows that there is a join between  $Q_2$  and  $T_2$ . Consider now an optimal move  $y'$  of Dominator. As already observed,  $y'$  belongs to  $T_1 \cup Q_1$ . Since the game ends with this move, it follows that  $T_1 \cup T_2 \subseteq N[y] \cup N[y']$ . Similarly, for all  $x \in T_1$ , there is a vertex  $x' \in T_2 \cup Q_2$  such that  $T_1 \cup T_2 \subseteq N[x] \cup N[x']$ . In conclusion,  $G$  is a gamburger and  $Q_1, T_1, T_2, Q_2$  is a gamburger structure for  $G$ , which completes the proof.  $\square$

If a gamburger  $G$  with a gamburger structure  $Q_1, T_1, T_2, Q_2$  contains a vertex  $x \in T_1$  such that  $T_1 \cup T_2 \subseteq N[x]$ , then we say that  $G$  is a *full-gamburger*. We will also say that the vertex  $x$  is a *full vertex* of  $G$ .

With this definition we can specialize Theorem 3.11 to  $(2, 2)$ -graphs as follows.

**Corollary 3.12.** *A graph  $G$  is a  $(2, 2)$ -graph with  $\text{diam}(G) = 3$  if and only if  $G$  is a full-gamburger.*

**Proof.** Assume first that  $G$  is a full-gamburger. By Theorem 3.11,  $\gamma'_g(G) = 2$  and  $\text{diam}(G) = 3$ . Note that a full vertex  $x$  from  $T_1$  is 2-dense. Hence  $\gamma_g(G) = 2$  holds by Proposition 3.6.

Conversely, suppose  $G$  is a  $(2, 2)$ -graph with diameter 3. By Theorem 3.11,  $G$  has a gamburger structure  $Q_1, T_1, T_2, Q_2$ . Moreover, by Lemma 3.9 we may without loss of generality assume that every vertex in  $T_i$  has a neighbor in  $T_{3-i}$ ,  $1 \leq i \leq 2$ . Let  $x$  be an optimal first move of Dominator in D-game. Then  $x$  belongs to  $T_1 \cup T_2$ , for if  $x$  would belong to  $Q_i$ , Staller could play in  $T_i$  without ending the game. Assume without loss of generality that  $x \in T_1$  and suppose that  $N[x]$  does not contain  $T_1 \cup T_2$ . Let  $y$  be a vertex of  $T_1 \cup T_2$  which is not in  $N[x]$ . If  $y$  belongs to  $T_1$ , Staller can play  $y$  and this will not end the game. If  $y$  belongs to  $T_2$ , we know that  $y$  has a neighbor in  $T_1$ , say  $y'$ . Clearly,  $y'$  is distinct from  $x$ , hence playing  $y'$  is a legal move because it newly dominates  $y$ . In both cases, we thus have a contradiction with  $\gamma_g(G) = 2$ . We conclude that  $T_1 \cup T_2 \subseteq N[x]$  and hence  $x$  is a full vertex of the full-gamburger  $G$ .  $\square$

#### 4. ON GRAPHS WITH GAME DOMINATION NUMBER 3

The class of graphs  $G$  with  $\gamma_g(G) = 3$  seems too rich to allow some nice characterization. In [2] the subclass of these graphs with the property that  $\gamma'_g(G) = 2$  was characterized. For instance, the cycles  $C_5$  and  $C_6$  both belong to this class. In view of Proposition 2.5, in the first part of this section we characterize the subclass of  $\gamma_g = 3$  graphs extremal with respect to the diameter, that is graphs with  $\gamma_g(G) = 3$  and  $\text{diam}(G) = 6$ . For instance, this class contains  $P_7$  and is disjoint from the above class of graphs realizing  $(3, 2)$ . That the two classes are disjoint follows from facts that  $\gamma(G) = 2$  holds for all graphs  $G$  from the first class and  $\gamma(G) \geq 3$  for all graphs  $G$  from the second class. In Fig. 3 the situation is demonstrated.




Figure 3. Subclasses of  $\gamma_g = 3$  graphs

In the second part of the section we then characterize the graphs  $G$  with  $\gamma'_g(G) = 3$  and  $\text{diam}(G) = 5$ .



#### 4.1. Graphs with $\gamma_g = 3$ and $\text{diam} = 6$

Full-gamburgers turn out to be useful also for the first main result of this section. In addition to this concept, we also introduce the following. If  $G$  is a connected graph, then a vertex  $u$  of  $G$  is *nice* if the following conditions are fulfilled.

- There exists  $v_1 \in S_1(u)$  such that  $N[v_1] = B_2(u)$ .
- There is a join between  $N(S_3(u)) \cap S_2(u)$  and  $S_3(u)$ .
- $S_3(u), S_4(u), S_5(u), S_6(u)$  is a full-gamburger structure with a full vertex in  $S_5(u)$ .
- $S_7(u) = \emptyset$ .

With these notations, the announced result reads as follows.

**Theorem 4.13.** *If  $G$  is a connected graph, then the following statements are equivalent.*

- (i)  $\gamma_g(G) = 3$  and  $\text{diam}(G) = 6$ .
- (ii) *Any diametrical pair of vertices contains at least one nice vertex.*
- (iii) *There exists a nice diametrical vertex.*

**Proof.** We first prove that (i) implies (ii). Let  $u_1$  and  $u_2$  be vertices of  $G$  with  $d(u_1, u_2) = 6$  and let  $P$  be a shortest  $u_1, u_2$ -path. Since  $\gamma_g(G) = 3 = \gamma_g(P_7)$ , and because the two neighbors of pendant vertices of  $P_7$  are the only optimal start moves for Dominator when playing D-game on  $P_7$ , we infer that Dominator's first move must be either in  $S_1(u_1)$  or in  $S_1(u_2)$ . Indeed, for otherwise Staller could guarantee (by considering the game restricted to  $P$ ) the game to last at least four moves. Let  $i \in \{1, 2\}$  be such that we have an optimal move for Dominator in  $S_1(u_i)$ . For convenience, we will write  $S_j$ ,  $0 \leq j \leq 6$ , instead of  $S_j(u_i)$  and set  $u = u_i$ .

Since the diameter of the graph is 6, the spheres  $S_j$  are non-empty and form a partition of  $V(G)$ . We are going to prove several claims that will establish the structure as described in (ii). Let  $x \in S_1$  be an optimal first move of Dominator.

**Claim 1.** There exists a vertex  $v_1 \in S_1$  such that  $N[v_1] = B_2(u)$ .

Staller's move could be in  $S_4$ . To finish the game in three turns, Dominator must then play in  $S_5 \cup S_6$ . Hence, his first move  $x$  must dominate the whole  $B_2(u)$  and therefore  $v_1 = x$  fulfills the claim.

**Claim 2.** There is a join between  $N(S_3) \cap S_2$  and  $S_3$ .

Staller could play her first move anywhere in  $N(S_3) \cap S_2$ . Such moves are actually legal because they dominate at least one new vertex in  $S_3$ . In that case Dominator has to answer in  $S_5$ . Hence, all the vertices in  $S_3$  must have been dominated by the Staller's move. Since this move of Staller in  $N(S_3) \cap S_2$  was arbitrary, we have a join between  $N(S_3) \cap S_2$  and  $S_3$ .

It remains to prove that  $S_3, S_4, S_5, S_6$  form a full-gamburger structure with a full vertex in  $S_5$ .

**Claim 3.**  $S_3$  and  $S_6$  are cliques.

Assume Staller plays first in  $S_3$  (resp.  $S_6$ ). Dominator must play next in  $S_5 \cup S_6$  (resp.  $S_3 \cup S_4$ ). Since this move of Dominator has to end the game, all the vertices of  $S_3$  (resp.  $S_6$ ) must have been dominated by the move of Staller. This move could be arbitrary in the sphere  $S_3$  (resp.  $S_6$ ), because  $x$  dominates no vertex in  $S_3$ . It follows that  $S_3$  (resp.  $S_6$ ) induces a clique.

**Claim 4.** There is a join between  $S_3$  and  $S_4$  and between  $S_5$  and  $S_6$ . Moreover, for any vertex  $y$  in  $S_4$  (resp. in  $S_5$ ) there is a vertex  $y'$  in  $S_5 \cup S_6$  (resp. in  $S_3 \cup S_4$ ) such that  $N[y] \cup N[y']$  contains  $S_4 \cup S_5$ .

Suppose Staller plays a vertex  $y$  in  $S_4$ . Dominator must play next in  $S_5 \cup S_6$ . Hence, all the vertices of  $S_3$  must have been dominated by the move of Staller. Because  $y$  is an arbitrary vertex from  $S_4$ , there must be a join between  $S_3$  and  $S_4$ . Let now  $y'$  in  $S_5 \cup S_6$  be an optimal answer of Dominator. Vertices of  $S_4 \cup S_5$  could only be dominated by the last two moves of the game, that is, by  $y$  and  $y'$ . Therefore, these two spheres are contained in  $N[y] \cup N[y']$ . Similarly, when Staller plays as her first move a vertex  $y$  from  $S_5$ , Dominator replies by playing a vertex  $y' \in S_3 \cup S_4$  and we get the same conclusion.

Until now we have already seen that  $S_3, S_4, S_5, S_6$  defines a gamburger structure. Hence it remains to prove the following.

**Claim 5.** The gamburger on  $S_3, S_4, S_5, S_6$  has a full vertex in  $S_5$ .

As in Claim 2, assume that Staller plays in  $S_2$ . Dominator has to answer in  $S_5$  and must dominate all the vertices in  $S_4, S_5$  and  $S_6$  with this move. Therefore, such a reply of Dominator in  $S_5$  is a full vertex of the gamburger with the gamburger structure  $S_3, S_4, S_5, S_6$ .

In conclusion we have proved that  $u$  is a nice vertex.

Since (ii) trivially implies (iii), it remains to prove that (iii) implies (i). Assume that there exists a nice diametrical vertex  $u$ . As above, we simply write  $S_i$  instead of  $S_i(u)$ , for any  $i \in \{0, \dots, 6\}$ . Let  $v_1$  be a vertex in  $S_1$  such that  $N[v_1] = B_2(u)$ .

We prove first that  $\text{diam}(G) = 6$ . Since  $S_3, S_4, S_5, S_6$  induce a gamburger structure, none of these spheres is empty. Hence  $\text{ecc}(u) = 6$  and  $\text{diam}(G) \geq 6$ . By Lemma 3.10, the subgraph induced by  $S_3, S_4, S_5, S_6$  has diameter 3. Moreover, using the vertex  $v_1 \in S_1$ , it is easy to show that all the vertices from  $B_2(u)$  are at distance at most 3 from any vertex in  $S_3$ . In conclusion,  $G$  has diameter 6.

Since  $\text{diam}(G) = 6$ , we have  $\gamma_g(G) \geq 3$ . Hence, to conclude the proof we need to show that Dominator could ensure the game to end in three turns. His strategy is to play  $v_1$  as the first move. After that, all the vertices in  $B_2(u)$  are dominated and the remaining legal moves for Staller are in  $G \setminus B_1(u)$ . First assume Staller plays in  $G \setminus B_2(u)$ . Since these vertices induce a gamburger, Theorem 3.11 implies that Dominator has an answer in  $G \setminus B_2(u)$ , such that all the vertices of this subgraph

are dominated after this move that clearly ends the game. Second, if Staller plays in  $S_2$ , then to be legal, this move has to belong to  $N(S_3)$ . Because of the join between  $N(S_3) \cap S_2$  and  $S_3$ , all the vertices in  $S_3$  are dominated. Dominator can then finish the game by playing a full vertex in  $S_5$ .  $\square$

We point out that both characterizations from Theorem 4.13 of graphs with  $\gamma_g(G) = 3$  and  $\text{diam}(G) = 6$  are useful if we wish to give a fast recognition algorithm for these graphs. Indeed, we only need to select a pair of diametrical vertices and check if one of them leads to the structure as described in (iii). If not, then we know from (ii) that no other diametrical vertex can give us the desired structure.

#### 4.2. Graphs with $\gamma'_g = 3$ and $\text{diam} = 5$

To characterize the graphs from the title, we introduce one more concept. We say that a connected graph  $G$  is a *double-gamburger*, if  $G$  is the disjoint union of non-empty subgraphs  $Q_1, R_1, T_1, T_2, R_2, Q_2$ , for which the following hold for any  $i \in \{1, 2\}$ .

- $Q_i$  and  $T_i$  induce cliques.
- There are joins between  $Q_i$  and  $R_i$ , and between  $R_i$  and  $T_i$ .
- $R_i$  induces a clique minus a matching  $M_i$ .
- If  $M_i$  is a perfect matching of  $R_i$ , then  $M_{3-i}$  is empty and there is a join between  $T_1$  and  $T_2$ .
- There exists a vertex  $x_i \in T_i$ , such that  $T_{3-i} \subseteq N(x_i)$ .
- There are no edges between  $Q_i$  and  $G \setminus R_i$  and between  $R_i$  and  $G \setminus (Q_i \cup T_i)$ .

See Fig. 4. We say that  $Q_1, R_1, T_1, T_2, R_2, Q_2$  is a *double-gamburger structure* for  $G$ .




Figure 4. Double-gamburger

**Lemma 4.14.** *A double-gamburgen has diameter 5.*

**Proof.** The graphs  $H_1$  and  $H_2$ , respectively induced by  $Q_1, R_1, T_1$  and  $T_2, R_2, Q_2$  have diameter 2. For any  $i \in \{1, 2\}$ , we have a vertex  $x \in T_i$ , such that  $T_{3-i} \subseteq N(x)$ . Using these vertices, it is straightforward to see that the distance between a vertex in  $H_i$  and the subgraph  $H_{3-i}$  is 3. In conclusion, a double-gamburgen has diameter 5.  $\square$

**Theorem 4.15.** *If  $G$  is a connected graph, then the following statements are equivalent.*

- (i)  $\gamma'_g(G) = 3$  and  $\text{diam}(G) = 5$ .
- (ii) *For any diametrical vertex  $u$ , defining  $Q_1 = \{u\} \cup W_1$ ,  $R_1 = S_1(u) \setminus W_1$ ,  $T_1 = S_2(u)$ ,  $T_2 = S_3(u)$ ,  $R_2 = S_4(u)$  and  $Q_2 = S_5(u)$ , where  $W_1 = S_1(u) \setminus N(S_2(u))$ , is a double-gamburgen structure for  $G$ .*
- (iii)  *$G$  is a double-gamburgen.*

**Proof.** We first prove that (i) implies (ii). Let  $u$  be a vertex of  $G$  with  $\text{ecc}(u) = 5$ . We are going to prove four claims which together establish that the subgraphs  $Q_1, R_1, T_1, T_2, R_2, Q_2$  as defined in (ii) form a double-gamburgen structure for  $G$ . Note first that these subgraphs are clearly not empty and there are no edges between  $Q_i$  and  $G \setminus R_i$ , as well as no edges between  $R_i$  and  $G \setminus (Q_i \cup T_i)$ .

**Claim 1.** For  $i \in \{1, 2\}$ ,  $Q_i$  induces a clique, there is a join between  $Q_i$  and  $R_i$  and there exists  $u \in T_{3-i}$  such that  $N(u)$  contains  $T_i$ .

Assume  $x \in Q_i$  is the first move of Staller. If Dominator plays in  $R_i \cup T_i$ , Staller will be able to play her second move in  $T_{3-i}$ . In that case,  $Q_{3-i}$  will remain undominated at the end of the third turn. If he plays in  $Q_{3-i}$ , she could answer in  $R_{3-i}$  and  $T_i$  will not be dominated. Hence Dominator's optimal answer is either in  $T_{3-i}$  or in  $R_{3-i}$ . In both cases, there is still a legal move for Staller in  $T_{3-i} \cup R_{3-i}$ . Since the game must end in three turns, this implies that  $Q_i \cup R_i$  has been entirely dominated by  $x$ . Hence,  $Q_i$  is a clique and there is a join between  $Q_i$  and  $R_i$ . Now assume, that Dominator's move was in  $T_{3-i}$ . Staller can play her last move in  $R_{3-i}$ . So, the game could end in three turns only if Dominator's move dominates all vertices in  $T_i$ . On the other hand, if Dominator played in  $R_{3-i}$ , Staller can play her last move in  $T_{3-i}$ . Once more, this move must dominate all  $T_i$ . In conclusion, we have at least one vertex in  $T_{3-i}$  whose neighborhood contains  $T_i$ .

**Claim 2.** For  $i \in \{1, 2\}$ ,  $T_i$  induces a clique and there is a join between  $T_i$  and  $R_i$ . By way of contradiction, assume there exists  $x \in T_i$ , such that  $N[x]$  does not contain  $T_i \cup R_i$  and let  $y$  be a vertex in  $(T_i \cup R_i) \setminus N[x]$ . Staller can play the vertex  $x$ . If Dominator plays in  $Q_i \cup R_i \cup T_i$ , Staller will be able to play her second move in  $T_{3-i}$ . If he plays in  $T_{3-i}$ , she could answer in  $Q_i$ . In both cases, the vertices in  $Q_{3-i}$  will not be dominated at the end of the third turn. Therefore, Dominator has to play in  $R_{3-i} \cup Q_{3-i}$ . Now, if  $y \in T_i$ , playing  $y$  is a legal move for Staller and after this move no vertex in  $Q_i$  is dominated. Otherwise, if  $y$  is in  $R_i$ , then by

the way we define  $Q_1$ ,  $T_i$  and  $R_i$ ,  $y$  has a neighbor  $y'$  in  $T_i$ . Since playing  $y'$  will dominate the new vertex  $y$ , it is a legal move. By playing it, Staller ensures that no vertex in  $Q_i$  is already dominated. In both cases, the game is not over at the end of the third turn. This is a contradiction with the assumption  $\gamma'_g(G) = 3$ .

**Claim 3.** For  $i \in \{1, 2\}$ ,  $R_i$  induces a clique from which a matching  $M_i$  has been removed.

If  $|R_i| \leq 2$ , the claim is trivially true. Assume now that  $|R_i| \geq 3$ . By way of contradiction, suppose that  $R_i$  does not induced a clique minus a matching. In other words, there exist distinct vertices  $y$ ,  $y'$  and  $x$  in  $R_i$  such that  $x$  is neither adjacent to  $y$  nor to  $y'$ . If Staller starts by playing  $y$ , Dominator's optimal answer is in  $T_{3-i} \cup R_{3-i} \cup Q_{3-i}$ . Otherwise, Staller could play her second move in  $T_{3-i}$ , and  $Q_{3-i}$  would remain undominated. After the move of Dominator,  $y'$  is still undominated. Hence Staller can play this vertex as her second move. Finally, after these three turns,  $x$  is not yet dominated, again contradicting  $\gamma'_g(G) = 3$ .

**Claim 4.** For  $i \in \{1, 2\}$ , if the matching  $M_i$  is perfect, then  $M_{3-i}$  is empty and there is a join between  $T_1$  and  $T_2$ .

Assume first there is not a join between  $T_1$  and  $T_2$ . Then Staller can choose a vertex in  $T_{3-i}$ , such that at least one vertex in  $T_i$  is not dominated. As in Claim 2, Dominator must play in  $R_i \cup Q_i$ . Since  $T_i$  is not completely dominated after Staller's first move, Dominator has no choice but to play in  $R_i$ . Moreover, Staller could play her last move in  $Q_{3-i} \cup R_{3-i}$ . So all the vertices in  $T_i \cup R_i \cup Q_i$  must be dominated by Dominator's move. Therefore we must have a vertex in  $R_i$  whose closed neighborhood contains at least  $R_i$ . In conclusion,  $M_i$  is not a perfect matching.

Suppose now that  $M_{3-i}$  is not empty (i.e.  $R_{3-i}$  does not induce a clique). Staller can play a vertex in  $R_{3-i}$ , whose closed neighborhood does not contain  $R_{3-i}$ . As in Claim 3, Dominator has to play in  $T_i \cup R_i \cup Q_i$ . But, Staller would be in all cases able to make her last move in  $R_{3-i}$ . Hence, the game could end in three turns only if Dominator can dominate all  $T_i \cup R_i \cup Q_i$  in one move. As above, it implies that the matching  $M_i$  is not perfect.

Statement (ii) obviously implies (iii). It remains to prove that (iii) leads to (i). Assume that  $G$  has a double-gamburger structure. By Lemma 4.14,  $\text{diam}(G) = 5$  which in turn implies that  $\gamma'_g(G) \geq 3$ . So, we only have to give a strategy for Dominator which ensures that the game ends in at most three turns. By symmetry, we have three cases.

**Case 1.** Staller's first move is in  $Q_1$ . Then Dominator chooses to play a vertex from  $T_2$  whose neighborhood contains  $T_1$ . Clearly, after such a move, all the vertices of  $G \setminus Q_2$  are dominated. Moreover, all the remaining legal moves for Staller are in  $R_2 \cup Q_2$ . Since there is a join between these two sets, whatever she plays, all the vertices in  $Q_2$  will be dominated and the game will be over in three turns.

**Case 2.** Staller's first move is in  $R_1$ . They are two possibilities. First, the matching  $M_2$  removed from a clique to get  $R_2$  is perfect. Therefore  $R_1$  is a clique. Dominator

selects a vertex in  $R_2$ . All the vertices of  $G$ , except one in  $R_2$ , are dominated after this move. Staller has no choice but to end the game by selecting a vertex in the closed neighborhood of this last undominated vertex. Second, the matching  $M_2$  is not perfect. Hence, there is a vertex  $x \in R_2$  whose neighborhood contains  $R_2$ . Dominator chooses this vertex. Now, there is at most one undominated vertex in  $G$ , which is actually in  $R_1$ . Staller has to dominate this vertex and the game is over after her move.

**Case 3.** Staller's first move is in  $T_1$ . Suppose first that we have a join between  $T_1$  and  $T_2$ . Dominator selects any vertex in  $Q_2$ . On the other hand, if there is no join between  $T_1$  and  $T_2$ , then  $M_2$  is not a perfect matching and we have a vertex  $x \in R_2$  such that  $N[x]$  contains  $T_2 \cup R_2 \cup Q_2$ . Dominator plays this vertex. In both situations, all the vertices in  $G \setminus Q_1$  are now dominated. The legal moves for Staller are only in  $Q_1 \cup R_1$ . Since there is a join between these two sets, whatever she chooses the game will end after her move.  $\square$

As for Theorem 4.13, both characterizations of graphs with  $\gamma'_g(G) = 3$  and  $\text{diam}(G) = 5$  are useful for a fast recognition algorithm.

To conclude the paper we show that a double-gamburger  $G$  always has a property  $\gamma_g(G) = 3$ . In other words, the following is true.

**Corollary 4.16.** *A graph  $G$  is a  $(3, 3)$ -graph of diameter 5 if and only if  $G$  is a double-gamburger. In particular, there is no  $(4, 3)$ -graph with diameter 5.*

**Proof.** By Theorem 4.15 we only need to prove that if  $G$  is a double-gamburger, then  $\gamma_g(G) = 3$ . Since  $\text{diam}(G) = 5$ , [3, Corollary 4.1] implies that  $\gamma_g(G) \geq 3$ . Hence we only have to prove that Dominator can end the game in at most three turns.

Let  $M_1$  and  $M_2$  be the two matchings defined by the double-gamburger structure of  $G$ . Assume first that both are not perfect. Then Dominator can start by playing a vertex in  $R_1$  which dominates all  $R_1$ . As his second move, he selects a vertex in  $R_2$  which dominates all  $R_2$ . Such a vertex is still available, because Staller plays optimally and hence she did not select such a vertex. Playing this way, all the vertices will be dominated at the end of the third turn. On the other hand, if one of the matchings, say  $M_1$  is perfect, by the double-gamburger structure of  $G$ ,  $R_2$  is a clique and we have a join between  $T_1$  and  $T_2$ . Dominator starts the game with any vertex of  $Q_1$ . If Staller answers in  $R_1 \cup T_1 \cup T_2$ , Dominator plays his second move in  $R_2$ . If Staller plays in  $R_2 \cup Q_2$ , then he plays in  $T_2$ . Note that these two cases are exhaustive, because the vertices in  $Q_1$  are not legal moves for Staller (since all the vertices in  $Q_1 \cup R_1$  are already dominated by Dominator's move). Finally, it is straightforward to show that in both cases all the vertices are dominated at the end of the third turn.  $\square$

**Acknowledgements.** The research was in part financed by Slovenian Research Agency under the grants N1-0043, P1-0297 and L7-5554, by the ANR-14-CE25-0006 project of the French National Research Agency, and by the grant CMIRA Explo'RA Doc from La Région Rhône Alpes. The authors thank the two referees

for carefully reading the manuscript and offering many suggestions that improved the article.

#### REFERENCES

1. B. BREŠAR, P. DORBEC, S. KLAVŽAR, G. KOŠMRLJ: *Complexity of the game domination problem*. Manuscript, 2014.
2. B. BREŠAR, P. DORBEC, S. KLAVŽAR, G. KOŠMRLJ: *How long one can bluff in the domination game*. Discuss. Math. Graph Theory, to appear.
3. B. BREŠAR, S. KLAVŽAR, G. KOŠMRLJ, D. F. RALL: *Guarded subgraphs and the domination game*. Discrete Math. Theor. Comput. Sci., **17** (2015), 161–168.
4. B. BREŠAR, S. KLAVŽAR, D. F. RALL: *Domination game and an imagination strategy*. SIAM J. Discrete Math., **24** (2010), 979–991.
5. B. BREŠAR, S. KLAVŽAR, G. KOŠMRLJ, D. F. RALL: *Domination game: extremal families of graphs for the 3/5-conjectures*. Discrete Appl. Math., **161** (2013), 1308–1316.
6. CS. BUJTÁS: *Domination game on trees without leaves at distance four*. Proceedings of the 8th Japanese-Hungarian Symposium on Discrete Mathematics and Its Applications (A. Frank, A. Recski, G. Wiener, eds.), June 4–7, 2013, Veszprém, Hungary, 73–78.
7. CS. BUJTÁS: *Domination game on forests*. Discrete Math., **338** (2015), 2220–2228.
8. CS. BUJTÁS: *On the game domination number of graphs with given minimum degree*. Electron. J. Combin., **22** (2015), paper #P3.29.
9. CS. BUJTÁS, S. KLAVŽAR: *Improved upper bounds on the domination number of graphs with minimum degree at least five*. Graphs Combin., **32** (2016), 511–519.
10. CS. BUJTÁS, S. KLAVŽAR, G. KOŠMRLJ: *Domination game critical graphs*. Discuss. Math. Graph Theory, **35** (2015), 781–796.
11. CS. BUJTÁS, Z. TUZA: *The disjoint domination game*. Discrete Math., to appear. DOI: doi:10.1016/j.disc.2015.04.028.
12. P. DORBEC, G. KOŠMRLJ, G. RENAULT: *The domination game played on unions of graphs*. Discrete Math., **338** (2015), 71–79.
13. M. A. HENNING, W. B. KINNERSLEY: *Domination game: A proof of the 3/5-conjecture for graphs with minimum degree at least two*. SIAM J. Discrete Math., **30** (2016), 20–35.
14. M. A. HENNING, S. KLAVŽAR, D. F. RALL: *Total version of the domination game*. Graphs Combin., **31** (2015), 1453–1462.
15. M. A. HENNING, S. KLAVŽAR, D. F. RALL: *The 4/5 upper bound on the game total domination number*. Combinatorica, to appear.
16. M. A. HENNING, D. F. RALL: *Progress towards the total domination game 3/4-conjecture*. ArXiv:1512.02916 [math.CO].
17. W. B. KINNERSLEY, D. B. WEST, R. ZAMANI: *Game domination for grid-like graphs*. unpublished manuscript, 2012.
18. W. B. KINNERSLEY, D. B. WEST, R. ZAMANI: *Extremal problems for game domination number*. SIAM J. Discrete Math., **27** (2013), 2090–2107.

- 
19. G. KOŠMRLJ: *Realizations of the game domination number*. J. Comb. Optim., **28** (2014), 447–461.
  20. M. J. NADJAFI-ARANI, M. SIGGERS, H. SOLTANI: *Characterisation of forests with trivial game domination numbers*. J. Comb. Optim., to appear. DOI: 10.1007/s10878-015-9903-9.

Faculty of Mathematics and Physics,  
University of Ljubljana, Ljubljana  
Slovenia

and

Faculty of Natural Sciences and Mathematics,  
University of Maribor, Maribor  
Slovenia

and

Institute of Mathematics, Physics and Mechanics,  
Ljubljana  
Slovenia

E-mail: [sandi.klavzar@fmf.uni-lj.si](mailto:sandi.klavzar@fmf.uni-lj.si)

Abelium, R&D d.o.o., Ljubljana  
Slovenia

and

Institute of Mathematics, Physics and Mechanics,  
Ljubljana  
Slovenia

E-mail: [gasper.kosmrlj@abelium.eu](mailto:gasper.kosmrlj@abelium.eu)

Institut Fourier, SFR Maths à Modeler,  
Université Grenoble Alpes, Grenoble  
France

E-mail: [simon.schmidt@ujf-grenoble.fr](mailto:simon.schmidt@ujf-grenoble.fr)

(Received May 13, 2015)

(Revised January 13, 2016)