



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Applied Mathematics Letters xx (xxxx) xxx–xxx

**Applied  
Mathematics  
Letters**

[www.elsevier.com/locate/aml](http://www.elsevier.com/locate/aml)

# A theorem on Wiener-type invariants for isometric subgraphs of hypercubes

Sandi Klavžar<sup>a,b,\*</sup>, Ivan Gutman<sup>c</sup>

<sup>a</sup> Department of Mathematics and Computer Science, PeF, University of Maribor, Koroška cesta 160, 2000 Maribor, Slovenia

<sup>b</sup> Institute of Mathematics, Physics and Mechanics, Jadranska 19, 1000 Ljubljana, Slovenia

<sup>c</sup> Faculty of Science, University of Kragujevac, P. O. Box 60, 34000 Kragujevac, Serbia and Montenegro

Received 22 November 2005; accepted 11 December 2005

## Abstract

Let  $d(G, k)$  be the number of pairs of vertices of a graph  $G$  that are at distance  $k$ ,  $\lambda$  a real (or complex) number, and  $W_\lambda(G) = \sum_{k \geq 1} d(G, k) k^\lambda$ . It is proved that for a partial cube  $G$ ,  $W_{\lambda+1}(G) = |\mathcal{F}| W_\lambda(G) - \sum_{F \in \mathcal{F}} W_\lambda(G \setminus F)$ , where  $\mathcal{F}$  is the partition of  $E(G)$  induced by the Djoković–Winkler relation  $\Theta$ . This result extends a previously known result for trees and implies several relations for distance-based topological indices.

© 2006 Published by Elsevier Ltd

**Keywords:** Graph distance; Hypercube; Partial cube; Wiener number; Hyper-Wiener index

## 1. Introduction

The *Wiener number* (or *Wiener index*)  $W(G)$  of a connected graph  $G$  is the sum of distances between all pairs of vertices of  $G$ , that is,

$$W(G) = \sum_{\{u,v\} \subseteq V(G) \times V(G)} d(u, v).$$

In the case of trees the Wiener number was introduced back in 1947 by Wiener in [25], hence the name of this graph invariant. Right up to today, it has been extensively investigated, above all in mathematical chemistry; see special issues of journals devoted to the topic [13,14], recent surveys [5,6], and recent papers [7–9].

The Wiener number can be extended to disconnected graphs as follows [12]. Denote by  $d(G, k)$  the number of pairs of vertices of  $G$  that are at distance  $k$ . Note that  $d(G, 0)$  and  $d(G, 1)$  represent the number of vertices and edges, respectively. Then  $W$  can be extended to disconnected graphs as  $W(G) = \sum_{k \geq 1} d(G, k) k$ . Moreover, this definition can be further generalized in the following natural way [11,12]:

$$W_\lambda(G) = \sum_{k \geq 1} d(G, k) k^\lambda,$$

\* Corresponding author at: Department of Mathematics and Computer Science, PeF, University of Maribor, Koroška cesta 160, 2000 Maribor, Slovenia. Tel.: +386 2 22 93 604; fax: +386 1 25 172 81.

E-mail addresses: sandi.klavzar@uni-mb.si (S. Klavžar), gutman@knez.uis.kg.ac.yu (I. Gutman).

where  $\lambda$  is some real (or complex) number. Several particular instances of the invariant  $W_\lambda$  have been previously studied. For instance,  $W_{-2}$ ,  $W_{-1}$ ,  $\frac{1}{2} W_2 + \frac{1}{2} W_1$ , and  $\frac{1}{6} W_3 + \frac{1}{2} W_2 + \frac{1}{3} W_1$  are the so-called Harary index, reciprocal Wiener index, hyper-Wiener index, and Tratch–Stankevich–Zefirov index; cf. [12] and references therein. In the chemical literature also  $W_{1/2}$  [27] as well as the general case  $W_\lambda$  were examined [10,11,15].

Let  $T$  be a tree; then in [12] the following recursive formula for  $W_\lambda$  has been obtained:

$$W_{\lambda+1}(T) = (n-1)W_\lambda(T) - \sum_{e \in E(T)} W_\lambda(T-e). \quad (1)$$

In this note we prove that if  $G$  is a partial cube and  $\mathcal{F}$  the partition of  $E(G)$  induced by the Djoković–Winkler relation  $\Theta$ , then

$$W_{\lambda+1}(G) = |\mathcal{F}|W_\lambda(G) - \sum_{F \in \mathcal{F}} W_\lambda(G \setminus F). \quad (2)$$

Since trees are partial cubes in which the partition  $\mathcal{F}$  is trivial, that is, every edge of a tree forms a class of the partition, (1) immediately follows from (2). In addition we will demonstrate that some known relations between distance-based topological indices follow from formula (2).

## 2. The main result

For  $u, v \in V(G)$ , let  $d_G(u, v)$  denote the length of a shortest path (also called a *geodesic*) in  $G$  from  $u$  to  $v$ . A subgraph  $H$  of a graph  $G$  is called *isometric* if  $d_H(u, v) = d_G(u, v)$  for all  $u, v \in V(H)$ . Isometric subgraphs of hypercubes are called *partial cubes*. Clearly, hypercubes are partial cubes, as well as trees and median graphs. Partial cubes form a well studied class of graphs; we refer the reader to classical references [1,4,26], the book [16], the recent paper [20] and references therein. For applications of partial cubes to mathematical chemistry see [3,17–19,21].

The *Djoković–Winkler relation*  $\Theta$  is defined on the edge set of a graph in the following way [4,26]. Edges  $e = xy$  and  $f = uv$  of a graph  $G$  are in relation  $\Theta$  if

$$d_G(x, u) + d_G(y, v) \neq d_G(x, v) + d_G(y, u).$$

Winkler [26] proved that among bipartite graphs,  $\Theta$  is transitive precisely for partial cubes; hence  $\Theta$  partitions the edge set of a partial cube. Let  $G$  be a partial cube and  $\mathcal{F} = \{F_1, F_2, \dots, F_r\}$  the partition of its edge set induced by the relation  $\Theta$ . Then we say that  $\mathcal{F}$  is the  $\Theta$ -partition of  $G$ .

For the proof of our main theorem we need the following facts about  $\Theta$ ; cf. [16,20].

**Lemma 1.** *Let  $G$  be a partial cube.*

- (i) *A path  $P$  in  $G$  is a geodesic if and only if no two different edges of  $P$  are in relation  $\Theta$ .*
- (ii) *Let  $F$  be a class of the  $\Theta$ -partition of  $G$ . Then  $G \setminus F_i$  consists of two connected components.*

We are now ready for our main result.

**Theorem 2.** *Let  $G$  be a partial cube and  $\mathcal{F}$  its  $\Theta$ -partition. Then for any real (or complex) number  $\lambda$ ,*

$$W_{\lambda+1}(G) = |\mathcal{F}|W_\lambda(G) - \sum_{F \in \mathcal{F}} W_\lambda(G \setminus F).$$

**Proof.** Let  $s$  be the diameter of  $G$ ; then

$$W_\lambda(G) = \sum_{k=1}^s d(G, k) k^\lambda.$$

Let  $\mathcal{F} = \{F_1, F_2, \dots, F_r\}$  and set

$$X = \sum_{i=1}^r W_\lambda(G \setminus F_i).$$

Let  $u$  and  $v$  be arbitrary vertices of  $G$ , where  $d(u, v) = k$ ,  $1 \leq k \leq s$ . Let  $P$  be a  $u, v$ -geodesic. By Lemma 1(i), the edges of  $P$  belong to pairwise different classes of  $\mathcal{F}$ . We may assume without loss of generality that they belong to  $F_1, F_2, \dots, F_k$ . By Lemma 1(ii),  $u$  and  $v$  belong to different connected components of  $G \setminus F_i$  for  $i = 1, \dots, k$ . On the other hand,  $u$  and  $v$  are in the same connected component of  $G \setminus F_i$  for  $i = k + 1, \dots, r$ . Clearly, in the latter case,  $d_{G \setminus F_i}(u, v) = k$ . It follows that the pair  $\{u, v\}$  contributes  $(r - k)$  times to  $X$ . Thus,

$$\begin{aligned} X &= \sum_{k=1}^s (r - k) d(G, k) k^\lambda \\ &= r \sum_{k=1}^s d(G, k) k^\lambda - \sum_{k=1}^s d(G, k) k^{\lambda+1} \\ &= r W_\lambda(G) - W_{\lambda+1}(G). \quad \square \end{aligned}$$

If  $F$  is a  $\Theta$ -class of the hypercube  $Q_n$ , then  $Q_n \setminus F$  consists of two disjoint copies of  $Q_{n-1}$ . Thus, by Theorem 2,  $W_{\lambda+1}(Q_n) = n W_\lambda(Q_n) - 2n W_\lambda(Q_{n-1})$ . By this recurrence relation it follows that  $W_\lambda(Q_n) = p_\lambda(n) 4^n$ , where  $p_\lambda(n)$  is a polynomial. This can also be seen from the formula  $W_\lambda(Q_n) = 2^{n-1} \sum_{k=1}^n \binom{n}{k} k^\lambda$ .

### 3. Applications

In this section we give two applications of Theorem 2. The first one is the following result for the Wiener number, first given in [19], and extended to the so-called  $L_1$ -graphs in [2].

Let  $G$  be a partial cube,  $\mathcal{F}$  its  $\Theta$ -partition, and  $F \in \mathcal{F}$ . Then we will denote the connected components of  $G \setminus F$  by  $G_1(F)$  and  $G_2(F)$ . Set  $n_1(F) = |G_1(F)|$  and  $n_2(F) = |G_2(F)|$ .

**Corollary 3.** *Let  $G$  be a partial cube and  $\mathcal{F}$  its  $\Theta$ -partition. Then*

$$W_1(G) = W(G) = \sum_{F \in \mathcal{F}} n_1(F) n_2(F).$$

**Proof.** Let  $n = |V(G)|$ ; then for any  $F \in \mathcal{F}$ ,  $n_1(F) + n_2(F) = n$ . Using Theorem 2 we can compute as follows:

$$\begin{aligned} W_1(G) &= |\mathcal{F}| W_0(G) - \sum_{F \in \mathcal{F}} W_0(G \setminus F) \\ &= |\mathcal{F}| \binom{n}{2} - \sum_{F \in \mathcal{F}} \left[ \binom{n_1(F)}{2} + \binom{n_2(F)}{2} \right] \\ &= |\mathcal{F}| \binom{n}{2} - \frac{1}{2} \sum_{F \in \mathcal{F}} [n^2 - n - 2n_1(F)n_2(F)] \\ &= |\mathcal{F}| \binom{n}{2} - \frac{1}{2} \sum_{F \in \mathcal{F}} (n^2 - n) + \sum_{F \in \mathcal{F}} n_1(F)n_2(F) \\ &= \sum_{F \in \mathcal{F}} n_1(F)n_2(F). \quad \square \end{aligned}$$

For the second application some more concepts are needed. The hyper-Wiener index  $WW$  is a topological index proposed by Randić [24] for trees and extended to all graphs by Klein et al. [22] as

$$WW(G) = \frac{1}{2} W_1(G) + \frac{1}{2} W_2(G).$$

Let  $G$  be a partial cube,  $\mathcal{F}$  its  $\Theta$ -partition, and  $F, F' \in \mathcal{F}$ ,  $F \neq F'$ . Then we will define  $n_{11}(F, F') = |G_1(F) \cap G_1(F')|$ ,  $n_{12}(F, F') = |G_1(F) \cap G_2(F')|$ ,  $n_{21}(F, F') = |G_2(F) \cap G_1(F')|$ , and  $n_{22}(F, F') = |G_2(F) \cap G_2(F')|$ . We say that the classes  $F$  and  $F'$  cross if  $n_{k\ell}(F, F') \neq 0$  for  $1 \leq k, \ell \leq 2$ , and write  $F \# F'$  to denote the fact that  $F$  and  $F'$  cross; see [20,23]. Now we can deduce from Theorem 2 the following result given in [17].




Fig. 1. Non-crossing classes  $F_i$  and  $F_j$ .

**Corollary 4.** Let  $G$  be a partial cube and  $\mathcal{F} = \{F_1, F_2, \dots, F_r\}$  its  $\Theta$ -partition. Then

$$WW(G) = W(G) + \sum_{i < j} [n_{11}(F_i, F_j)n_{22}(F_i, F_j) + n_{12}(F_i, F_j)n_{21}(F_i, F_j)].$$

**Proof.** By Theorem 2,  $W_2(G) = rW(G) - \sum_{i=1}^r W(G \setminus F_i)$ . On the other hand,  $WW(G) = W(G)/2 + W_2(G)/2$ .

Combining these two equalities we get

$$WW(G) = W(G) + \frac{1}{2} \left[ (r-1)W(G) - \sum_{i=1}^r W(G \setminus F_i) \right]. \quad (3)$$

By Corollary 3 we have

$$(r-1)W(G) = \sum_{j=1}^{r-1} \sum_{i=1}^r n_1(F_i)n_2(F_i) = \sum_{i=1}^r \sum_{j=1}^{r-1} n_1(F_i)n_2(F_i), \quad (4)$$

while on the other hand

$$\sum_{i=1}^r W(G \setminus F_i) = \sum_{i=1}^r [W(G_1(F_i)) + W(G_2(F_i))]. \quad (5)$$

Combining (4) and (5) with (3) we obtain

$$WW(G) = W(G) + \frac{1}{2} \sum_{i=1}^r \left[ \sum_{j=1}^{r-1} n_1(F_i)n_2(F_i) - W(G_1(F_i)) - W(G_2(F_i)) \right]. \quad (6)$$

Having in mind Corollary 3 we now consider the contribution of a fixed pair of classes  $F_i$  and  $F_j$  to the right-hand side sum in (6). For the rest of the proof let  $n_{11}$ ,  $n_{12}$ ,  $n_{21}$ , and  $n_{22}$  denote  $n_{11}(F_i, F_j)$ ,  $n_{12}(F_i, F_j)$ ,  $n_{21}(F_i, F_j)$ , and  $n_{22}(F_i, F_j)$ , respectively.

Suppose first that  $F_i$  and  $F_j$  cross. Then the contribution of the pair  $F_i, F_j$  is

$$[(n_{11} + n_{12})(n_{21} + n_{22}) + (n_{11} + n_{21})(n_{12} + n_{22})] - [(n_{11}n_{12} + n_{21}n_{22}) + (n_{11}n_{21} + n_{12}n_{22})] \\ = 2n_{11}n_{22} + 2n_{12}n_{21}.$$

If  $F_i, F_j$  do not cross, then there are four possibilities for how  $F_i$  and  $F_j$  are related; the possibilities are shown in Fig. 1.

Then the contributions of the classes  $F_i$  and  $F_j$  are, respectively,

$$(i) (n_{11} + n_{12})n_{22} + n_{11}(n_{12} + n_{22}) - (n_{11}n_{12} + n_{12}n_{22}) = 2n_{11}n_{22},$$

- (ii)  $(n_{11} + n_{12})n_{21} + n_{12}(n_{11} + n_{21}) - (n_{12}n_{11} + n_{11}n_{21}) = 2n_{12}n_{21}$ ,
- (iii)  $(n_{21} + n_{22})n_{11} + n_{22}(n_{11} + n_{21}) - (n_{21}n_{22} + n_{21}n_{11}) = 2n_{11}n_{22}$ ,
- (iv)  $(n_{21} + n_{22})n_{12} + n_{21}(n_{12} + n_{22}) - (n_{21}n_{22} + n_{22}n_{21}) = 2n_{12}n_{21}$ .

Since in cases (i), (ii), (iii), and (iv) we have  $n_{21} = 0$ ,  $n_{22} = 0$ ,  $n_{12} = 0$ , and  $n_{11} = 0$ , respectively, in all cases the contribution of  $F_i$  and  $F_j$  to the right-hand side sum in (6) can be written as

$$2n_{11}n_{22} + 2n_{12}n_{21}$$

which completes the argument.  $\square$

## Acknowledgments

The first author was supported in part by the Ministry of Science of Slovenia under the grant P1-0297. We would also like to thank a referee for pointing out two missing cases in the original version of the proof of Corollary 4.

## References

- [1] V. Chepoi,  $d$ -Convexity and isometric subgraphs of Hamming graphs, *Cybernetics* 1 (1988) 6–9.
- [2] V. Chepoi, M. Deza, V. Grishukhin, Clin d'oeil on  $L_1$ -embeddable planar graphs, *Discrete Appl. Math.* 80 (1997) 3–19.
- [3] V. Chepoi, S. Klavžar, The Wiener index and the Szeged index of benzenoid systems in linear time, *J. Chem. Inf. Comput. Sci.* 37 (1997) 752–755.
- [4] D. Djoković, Distance preserving subgraphs of hypercubes, *J. Combin. Theory Ser. B* 14 (1973) 263–267.
- [5] A.A. Dobrynin, R. Entringer, I. Gutman, Wiener index of trees: theory and applications, *Acta Appl. Math.* 66 (2001) 211–249.
- [6] A.A. Dobrynin, I. Gutman, S. Klavžar, P. Žigert, Wiener index of hexagonal systems, *Acta Appl. Math.* 72 (2002) 247–294.
- [7] A.A. Dobrynin, L.S. Mel'nikov, Trees and their quadratic line graphs having the same Wiener index, *MATCH Commun. Math. Comput. Chem.* 50 (2004) 146–164.
- [8] A.A. Dobrynin, L.S. Mel'nikov, Wiener index, line graphs and the cyclomatic number, *MATCH Commun. Math. Comput. Chem.* 53 (2005) 209–214.
- [9] A.A. Dobrynin, L.S. Mel'nikov, Wiener index for graphs and their line graphs with arbitrary large cyclomatic numbers, *Appl. Math. Lett.* 18 (2005) 307–312.
- [10] B. Furtula, I. Gutman, Ž. Tomović, A. Vesel, I. Pesek, Wiener-type topological indices of phenylenes, *Indian J. Chem.* 41A (2002) 1767–1772.
- [11] I. Gutman, A property of the Wiener number and its modifications, *Indian J. Chem.* 36A (1997) 128–132.
- [12] I. Gutman, A.A. Dobrynin, S. Klavžar, L. Pavlović, Wiener-type invariants of trees and their relation, *Bull. Inst. Combin. Appl.* 40 (2004) 23–30.
- [13] I. Gutman, S. Klavžar, B. Mohar (Eds.), Fifty years of the Wiener index, *MATCH Commun. Math. Comput. Chem.* 35 (1997) 1–259.
- [14] I. Gutman, S. Klavžar, B. Mohar (Eds.), Fiftieth anniversary of the Wiener index, *Discrete Appl. Math.* 80 (1) (1997) 1–113.
- [15] I. Gutman, D. Vidović, L. Popović, On graph representation of organic molecules — Cayley's plerograms vs. his kenograms, *J. Chem. Soc. Faraday Trans.* 94 (1998) 857–860.
- [16] W. Imrich, S. Klavžar, *Product Graphs: Structure and Recognition*, John Wiley & Sons, New York, 2000.
- [17] S. Klavžar, Applications of isometric embeddings to chemical graphs, *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.* 51 (2000) 249–259.
- [18] S. Klavžar, I. Gutman, Wiener number of vertex-weighted graphs and a chemical application, *Discrete Appl. Math.* 80 (1997) 73–81.
- [19] S. Klavžar, I. Gutman, B. Mohar, Labeling of benzenoid systems which reflects the vertex-distance relations, *J. Chem. Inf. Comput. Sci.* 35 (1995) 590–593.
- [20] S. Klavžar, H.M. Mulder, Partial cubes and crossing graphs, *SIAM J. Discrete Math.* 15 (2002) 235–251.
- [21] S. Klavžar, A. Vesel, P. Žigert, I. Gutman, Binary coding of Kekulé structures of catacondensed benzenoid hydrocarbons, *Comput. Chem.* 25 (2001) 569–575.
- [22] D.J. Klein, I. Lukovits, I. Gutman, On the definition of the hyper-Wiener index for cycle-containing structures, *J. Chem. Inf. Comput. Sci.* 35 (1995) 50–52.
- [23] F.R. McMorris, H.M. Mulder, F.R. Roberts, The median procedure on median graphs, *Discrete Appl. Math.* 84 (1998) 165–181.
- [24] M. Randić, Novel molecular descriptor for structure–property studies, *Chem. Phys. Lett.* 211 (1993) 478–483.
- [25] H. Wiener, Structural determination of paraffin boiling points, *J. Amer. Chem. Soc.* 69 (1947) 17–20.
- [26] P. Winkler, Isometric embeddings in products of complete graphs, *Discrete Appl. Math.* 7 (1984) 221–225.
- [27] H.Y. Zhu, D.J. Klein, I. Lukovits, Extensions of the Wiener number, *J. Chem. Inf. Comput. Sci.* 36 (1996) 420–428.