

Comparing Wiener complexity with eccentric complexity

Kexiang Xu^a, Vesna Iršič^{b,c}, Sandi Klavžar^{b,c,d}, Huimin Li^a

^a College of Science, Nanjing University of Aeronautics & Astronautics,
Nanjing, Jiangsu 210016, PR China

^b Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia

^c Faculty of Mathematics and Physics, University of Ljubljana, Slovenia

^d Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia

kexxu1221@126.com (K. Xu)

vesna.irsic@fmf.uni-lj.si (V. Iršič)

sandi.klavzar@fmf.uni-lj.si (S. Klavžar)

1213966965@qq.com (H. Li)

Abstract

The transmission of a vertex v of a graph G is the sum of distances from v to all the other vertices in G . The Wiener complexity of G is the number of different transmissions of its vertices. Similarly, the eccentric complexity of G is defined as the number of different eccentricities of its vertices. In this paper these two complexities are compared. The complexities are first studied on Cartesian product graphs. Transmission indivisible graphs and arithmetic transmission graphs are introduced to demonstrate sharpness of upper and lower bounds on the Wiener complexity, respectively. It is shown that for almost all graphs the Wiener complexity is not smaller than the eccentric complexity. Several families of graphs in which the complexities are equal are constructed. Using the Cartesian product, it is also proved that the eccentric complexity can be arbitrarily larger than the Wiener complexity.

Keywords: graph distance; Wiener complexity; eccentric complexity; Cartesian product of graphs; graph of diameter 2

AMS Math. Subj. Class. (2010): 05C12, 05C76

1 Introduction

If $G = (V(G), E(G))$ is a graph, then $d_G(u, v)$ denotes the shortest-path distance between vertices $u, v \in V(G)$. The *transmission* $\text{Tr}_G(v)$ of a vertex $v \in V(G)$ is the sum of distances from v to the vertices in G , that is,

$$\text{Tr}_G(v) = \sum_{u \in V(G)} d_G(u, v).$$

The Wiener index $W(G)$ of G can then be defined as $W(G) = \frac{1}{2} \sum_{v \in V(G)} \text{Tr}_G(v)$. The *eccentricity* $\text{ec}_G(v)$ of a vertex $v \in V(G)$ is the maximum distance from v to other vertices of G :

$$\text{ec}_G(v) = \max_{u \neq v} d_G(u, v).$$

Eccentricity is a central concept of metric graph theory and has many applications elsewhere, in particular in location theory and in chemical graph theory. In the latter area, important eccentricity-based graph invariants (alias topological indices in mathematical chemistry) include the first and the second Zagreb eccentricity indices [22], eccentric connectivity index [21, 27], and connective eccentricity index [11]. For mathematical properties of these invariants see [9, 20, 23, 24, 26].

The *Wiener complexity* $C_W(G)$ of a graph G was introduced in [1] (under the name Wiener dimension) as the number of different transmission of vertices in G :

$$C_W(G) = |\{\text{Tr}_G(v) : v \in V(G)\}|.$$

The Wiener complexity of graphs has been further investigated in [4, 13, 14]. In the same spirit as the Wiener complexity is defined, the connective eccentric complexity [3] and the eccentric complexity [2] have been recently introduced. The *eccentric complexity* $C_{\text{ec}}(G)$ of a graph G is the number of different eccentricities in G . Equivalently,

$$C_{\text{ec}}(G) = \text{diam}(G) - \text{rad}(G) + 1,$$

where $\text{diam}(G) = \max_{v \in V(G)} \text{ec}_G(v)$ is the *diameter* of G and $\text{rad}(G) = \min_{v \in V(G)} \text{ec}_G(v)$ is the *radius* of G .

In view of the conceptional similarities between the Wiener complexity and the eccentric complexity, we compare in this paper these two complexities and proceed as follows. In the rest of this section we list definitions, concepts, and known results needed. In Section 2 we consider the Wiener complexity and the eccentric complexity of Cartesian products. In particular, a new lower bound on the Wiener complexity is proved and shown to be sharp using the so-called arithmetic transmission graphs.

To demonstrate that the Wiener complexity of a Cartesian product can be equal to the product of the complexities of the factors, transmission indivisible graphs are introduced. In Section 3 we first prove that $C_{ec}(G) \leq C_W(G)$ holds for almost all graphs G . Consequently, in Subsections 3.1 and 3.2, we consider the graphs G for which $C_{ec}(G) = C_W(G)$ and $C_{ec}(G) > C_W(G)$ holds, respectively. We construct several families of graphs, among them two families of product graphs, for which the equality holds. Using the Cartesian product, we find an infinite family of graphs G with the property $C_{ec}(G) > C_W(G)$. Moreover, the construction shows that the difference can be arbitrarily large.

1.1 Preliminaries

If k is a positive integer, then $[k] = \{1, \dots, k\}$. The degree of a vertex v of a graph G is denoted by $\deg_G(v)$. If G is a graph, then $n(G)$ denotes the order of G .

The *center* $C(G)$ of a graph G is the set of vertices of G of the minimum eccentricity, these vertices being called *central*. G is *self-centered* [5] if all its vertices have the same eccentricity, that is, if and only if $C_{ec}(G) = 1$.

Let G be a graph. The *transmission set* $\text{Tr}(G)$ of G is the set of the transmissions of its vertices, that is, $\text{Tr}(G) = \{\text{Tr}_G(v) : v \in V(G)\}$. The *eccentricity set* $\text{Ec}(G)$ of a graph G is the set of the eccentricities of its vertices, that is, $\text{Ec}(G) = \{\text{ec}_G(v) : v \in V(G)\}$. A graph G is *transmission regular* [16] if all its vertices have the same transmission. In other words, transmission regular graphs are precisely the graphs G with $C_W(G) = 1$. In addition, G is *transmission irregular* [4] if all its vertices have pairwise different transmissions, that is, if and only if $C_W(G) = n(G)$. We will make use of the following easy result on the transmission.

Proposition 1.1 ([1]) *Let G be a graph with $\text{diam}(G) = 2$. If $v \in V(G)$ with $\text{ec}_G(v) = 2$, then $\text{Tr}_G(v) = 2n(G) - 2 - \deg_G(v)$.*

The *Cartesian product* $G \square H$ of graphs G and H is the graph with $V(G \square H) = V(G) \times V(H)$, vertices (g, h) and (g', h') being adjacent if $gg' \in E(G)$ and $h = h'$, or $g = g'$ and $hh' \in E(H)$.

2 Complexities on Cartesian products

While comparing the Wiener complexity and the eccentric complexity we will extensively use the Cartesian product operation. In this section we hence recall known, and derive new related results.

It is well known that the distance function is additive on Cartesian product graphs. More precisely, if G and H are graphs, then

$$d_{G \square H}((g, h), (g', h')) = d_G(g, g') + d_G(h, h') \quad (1)$$

holds for arbitrary vertices $(g, h), (g', h') \in V(G \square H)$, cf. [12, Proposition 5.1]. This, in particular, implies that the diameter and radius are additive functions on Cartesian product graphs, which in turn gives the following closed formula for the eccentric complexity of Cartesian products.

Theorem 2.1 ([2, Theorem 11]) *If G and H are connected graphs, then we have $C_{ec}(G \square H) = C_{ec}(G) + C_{ec}(H) - 1$.*

The Wiener complexity of Cartesian products is more involved. The distance formula (1) yields

$$\begin{aligned} \text{Tr}_{G \square H}((g, h)) &= \sum_{(g', h') \in V(G \square H)} d_{G \square H}((g, h), (g', h')) \\ &= \sum_{(g', h') \in V(G \square H)} (d_G(g, g') + d_G(h, h')) \\ &= n(H) \sum_{g' \in V(G)} d_G(g, g') + n(G) \sum_{h' \in V(H)} d_G(h, h') \\ &= n(H) \text{Tr}_G(g) + n(G) \text{Tr}_H(h), \end{aligned} \quad (2)$$

a result deduced earlier in [1]. Consequently,

$$C_W(G \square H) = |\{n(H) \text{Tr}_G(g) + n(G) \text{Tr}_H(h) : g \in V(G), h \in V(H)\}|, \quad (3)$$

from which we immediately get:

$$\max\{C_W(G), C_W(H)\} \leq C_W(G \square H) \leq C_W(G)C_W(H). \quad (4)$$

The lower bound in (4) can be improved as follows.

Proposition 2.2 *If G and H are graphs, then*

$$C_W(G \square H) \geq C_W(G) + C_W(H) - 1.$$

Proof. Let $\text{Tr}(G) = \{x_1, \dots, x_s\}$ and $\text{Tr}(H) = \{y_1, \dots, y_t\}$, where $x_1 < \dots < x_s$ and $y_1 < \dots < y_t$. Then the set X defined as

$$\{x_1 n(H) + y_1 n(G), \dots, x_1 n(H) + y_t n(G), x_2 n(H) + y_1 n(G), \dots, x_s n(H) + y_t n(G)\}$$

contains pairwise different integers, and $X \subseteq \text{Tr}(G \square H)$ by (2). Since

$$|X| = s + t - 1 = C_W(G) + C_W(H) - 1,$$

the result follows. \square

For the Cartesian product of a graph by itself, (3) simplifies as follows:

$$\begin{aligned} C_W(G \square G) &= |\{n(G)\text{Tr}_G(g) + n(G)\text{Tr}_G(g') : g, g' \in V(G)\}| \\ &= |\{\text{Tr}_G(g) + \text{Tr}_G(g') : g, g' \in V(G)\}|. \end{aligned} \quad (5)$$

If G is a graph and H a graph with $C_W(H) = 1$, then $C_W(G \square H) = C_W(G)$, a result first reported in [1]. Hence the lower bound in (4) is best possible, it coincides with that in Proposition 2.2. On the other hand, the sharpness of the upper bound in (4) was not discussed in [1]. To establish the sharpness also for the upper bound we introduce the following notion. A graph G is *transmission indivisible* if $n(G) \nmid (\text{Tr}_G(u) - \text{Tr}_G(v))$ for every two distinct vertices $u, v \in V(G)$.

Theorem 2.3 *Let G and H be graphs. If at least one of G and H is transmission indivisible, and $\gcd(n(G), n(H)) = 1$, then $C_W(G \square H) = C_W(G)C_W(H)$.*

Proof. Let $\text{Tr}(G) = \{p_i : i \in [s]\}$ and $\text{Tr}(H) = \{q_j : j \in [t]\}$. Then in view of (3),

$$C_W(G \square H) = |\{n(H)p_i + n(G)q_j : i \in [s], j \in [t]\}|.$$

To prove the assertion of the theorem we need to show that $n(H)p_i + n(G)q_j \neq n(H)p_{i'} + n(G)q_{j'}$ for every $\{i, i'\} \in \binom{[s]}{2}$ and every $\{j, j'\} \in \binom{[t]}{2}$. Suppose on the contrary that for some such pairs $\{i, i'\}$ and $\{j, j'\}$ we have $n(H)p_i + n(G)q_j = n(H)p_{i'} + n(G)q_{j'}$, that is, $n(H)(p_i - p_{i'}) = n(G)(q_{j'} - q_j)$. Because $\gcd(n(G), n(H)) = 1$ we infer that $n(G) \mid (p_i - p_{i'})$ and $n(H) \mid (q_{j'} - q_j)$. But this means that neither G nor H is transmission indivisible, a contradiction. \square

Note that if G is transmission indivisible, then the transmissions of all the vertices of G are pairwise different, that is, G is transmission irregular. Although almost all graphs are not transmission irregular, an infinite family of transmission irregular trees was constructed in [4] and an infinite family of transmission irregular trees of even order in [7]. Moreover, an infinite family of transmission irregular 2-connected graphs was constructed in [6] and an infinite family of 3-connected cubic transmission irregular graphs in [8].

A transmission irregular graph need not be transmission indivisible. In Fig. 1 a transmission irregular but non-transmission indivisible tree of order 7 is shown,




Figure 1: A transmission irregular but non-transmission indivisible tree




Figure 2: Two of the smallest interval irregular graphs

where the transmission is given for each vertex. Sporadic transmission indivisible graphs of order 7 and 8 are shown in Fig. 2, where along with each vertex its transmission is stated.

Note that the transmission of the vertices in each of the two examples from Fig. 2 are consecutive integers from the intervals $[7..13]$ and $[8..15]$, respectively, which makes these examples particularly interesting. Such graphs were named *interval (transmission) irregular graphs* in [6]. Interval irregular graphs are transmission indivisible and hence Theorem 2.3 applies. We have checked by computer that there are 1, 2, 13, and 0 interval irregular graphs on 7, 8, 9, and 10 vertices, respectively. In addition, Dobrynin [6] reports that there are no less than 207 interval irregular 2-connected graphs of order 11. Their respective intervals of transmissions are $[13..23]$ (154 graphs), $[15..25]$ (51 graphs), and $[17..27]$ (2 graphs). The existence of an infinite family of interval irregular graphs is an open problem.

With interval irregular graphs in hands (and hence with transmission indivisible ones) the following result makes sense.

Corollary 2.4 *If G is a transmission indivisible graph, then there exists a family of graphs $\{H_i\}_{i \geq 1}$ such that $C_W(G \square H_i) = C_W(G)C_W(H_i)$.*

Proof. Let $\{p_i\}_{i \geq 1}$ be a set of primes each larger than $n(G)$ and let H_i be a graph of order p_i . Then $\gcd(n(G), n(H_i)) = 1$ and the result follows from Theorem 2.3. \square

By a computer search (using [17]) we have checked that the class of transmission indivisible graphs is strictly larger than the class of interval irregular graphs. There are no such examples on up to and including 10 vertices. However, a bit surprisingly, there are 221 graphs on 11 vertices that are transmission indivisible but not interval irregular. Among them there are no trees, but one finds 14 graphs which are 2-connected, an example can be seen in Fig. 3.




Figure 3: A 2-connected transmission indivisible graph which is not interval irregular

A graph G is *arithmetic transmission* if the ordered elements of $\text{Tr}(G)$ form an arithmetic progression. Moreover, if $\text{Tr}(G)$ has step a , we say that G has step a . (In Subsection 3.2 see an example of arithmetic transmission graph with step 4.) Below we present two results in which the lower bound is attained in Proposition 2.2.

Proposition 2.5 *If G is an arithmetic transmission graph, then $C_W(G \square G) = 2C_W(G) - 1$.*

Proof. Let $\text{Tr}(G) = \{x_1, \dots, x_k\}$, where $x_{i+1} - x_i = a$ for $i \in [k - 1]$. From (5) we get that $\text{Tr}(G \square G)$ is the set

$$\{2n(G)x_1, n(G)(x_1 + x_2), n(G)(x_1 + x_3), \dots, n(G)(x_{k-1} + x_k), 2n(G)x_k\},$$

which is an arithmetic progression of cardinality $2k - 1$ with step $n(G)a$. \square

Proposition 2.6 *Let G and H be arithmetic transmission graphs. If $\text{Tr}(H) \subseteq \text{Tr}(G)$ and $n(G) = n(H)$, then $C_W(G \square H) = C_W(G) + C_W(H) - 1$.*

Proof. Set $n = n(G) = n(H)$ and $\text{Tr}(G) = \{x_1, x_2, \dots, x_k\}$. We may without loss of generality assume that $\text{Tr}(H) = \{x_1, x_2, \dots, x_j\}$, where $j \leq k$. Then $|\text{Tr}(G)| = k = \frac{x_k - x_1}{a} + 1$ and $|\text{Tr}(H)| = j = \frac{x_j - x_1}{a} + 1$. By (5), $\text{Tr}(G \square H) = \{n(x_1 + x_1), \dots, n(x_k + x_j)\}$ and hence

$$\begin{aligned} |\text{Tr}(G \square H)| &= \frac{(x_k + x_j) - (x_1 + x_1)}{a} + 1 \\ &= \frac{x_k - x_1}{a} + \frac{x_j - x_1}{a} + 1 + 1 - 1 \\ &= |\text{Tr}(G)| + |\text{Tr}(H)| - 1. \end{aligned}$$

□

In Fig. 4 graphs G and H are shown which satisfy the conditions of Proposition 2.6.




Figure 4: Graphs G (right) and H (left) with $\text{Tr}(H) \subset \text{Tr}(G)$

Clearly, any interval irregular graph is arithmetic transmission as a special case. Hence Propositions 2.5 and 2.6 also apply for interval irregular graphs. Next we provide a result on the Cartesian product of arithmetic transmission graphs for which the upper bound in (4) is attained.

Theorem 2.7 *Let G and H be arithmetic transmission graphs with steps a and b , respectively. If $n(H)(C_W(G) - 1)a < n(G)b$, then $C_W(G \square H) = C_W(G)C_W(H)$.*

Proof. Let $\text{Tr}(G) = \{x_1, \dots, x_k\}$ and $\text{Tr}(H) = \{y_1, \dots, y_j\}$. Thus $C_W(G) = k$, $C_W(H) = j$ and

$$C_W(G \square H) = |\{n(H)x_p + n(G)y_q : p \in [k], q \in [j]\}|$$

by (3). Let $A_i = \{n(H)x_1 + n(G)y_i, \dots, n(H)x_k + n(G)y_i\}$ for $i \in [j]$. Therefore, we have $\text{Tr}(G \square H) = \bigcup_{i=1}^j A_i$. From the assumption, we have $n(H)(k - 1)a < n(G)b$.

It follows that

$$\begin{aligned}
n(H)x_k + n(G)y_i &= n(H)(x_1 + (k-1)a) + n(G)(y_{i+1} - b) \\
&= (n(H)x_1 + n(G)y_{i+1}) + (n(H)(k-1)a - n(G)b) \\
&< n(H)x_1 + n(G)y_{i+1}.
\end{aligned}$$

This implies that $\max\{s : s \in A_i\} < \min\{s : s \in A_{i+1}\}$, hence $A_s \cap A_t = \emptyset$ for any $s, t \in [k]$, $s \neq t$. We conclude that $C_W(G \square H) = kj$. \square

3 Comparing $C_{\text{ec}}(G)$ with $C_W(G)$

It was proved in [2] that $C_{\text{ec}}(G) \leq \lceil \frac{n(G)}{2} \rceil$ for any graph G . Therefore, if G is a transmission irregular graph, then $C_W(G) > C_{\text{ec}}(G)$. Actually, this is a phenomena that is very common as the next result shows.

Proposition 3.1 *For almost all graphs G , we have $C_{\text{ec}}(G) \leq C_W(G)$.*

Proof. It is well-known that almost all graphs have diameter 2. So let G be a graph with $\text{diam}(G) = 2$. Then $C_{\text{ec}}(G) \leq 2$. There is nothing to show if $C_{\text{ec}}(G) = 1$, so let $C_{\text{ec}}(G) = 2$ in which case we have $\text{Ec}(G) = \{1, 2\}$. But then G contains at least one vertex of degree $n(G) - 1$, and at least one vertex of smaller degree. Since their transmissions are different by Proposition 1.1, $C_W(G) \geq 2$. \square

We next show that the difference $C_W(G) - C_{\text{ec}}(G)$ can be arbitrarily large. For this recall that the d -cube Q_d , $d \geq 1$, has the vertex set $\{0, 1\}^d$, two vertices in Q_d are adjacent if they differ in precisely one coordinate. Let Q_d^- be the graph obtained from Q_d by removing an arbitrary vertex.

Proposition 3.2 *If $d \geq 2$, then $C_W(Q_d^-) - C_{\text{ec}}(Q_d^-) = d - 2$.*

Proof. We may without loss of generality assume that $V(Q_d^-) = V(Q_d) \setminus \{0^d\}$. Then $\text{ec}_{Q_d^-}(1^d) = d - 1$ and $\text{ec}_{Q_d^-}(v) = d$ for all other vertices v of Q_d^- . Thus $C_{\text{ec}}(Q_d^-) = 2$.

Let $x \in V(Q_d^-)$. Then $\text{Tr}_{Q_d^-}(x) = \text{Tr}_{Q_d}(x) \setminus \{d_{Q_d}(x, 0^d)\}$. Since there are precisely d different values of $d_{Q_d}(x, 0^d)$, we conclude that $\text{Tr}(Q_d^-) = d$. \square

Corollary 3.3 *If $d \geq 2$, then there exists a graph G with $\text{diam}(G) = d$ such that $C_W(G) > C_{\text{ec}}(G)$.*

Proof. If $d = 2$, then consider the paw graph P for which we have $\text{Ec}(P) = \{1, 2\}$ and $\text{Tr}(P) = \{3, 4, 5\}$. For $d \geq 3$, apply Proposition 3.2. \square

3.1 Graphs G with $C_W(G) = C_{ec}(G)$

Let G be a graph. Then, by definition, $C_{ec}(G) = C_W(G) = 1$ if and only if G is a self-centered, transmission regular graph. In the next result we present several additional classes of graphs G for which $C_W(G) = C_{ec}(G)$ holds. To state the result, we need some further definitions. If the eccentricity of the vertices of a self-centered graph G is k , we say that G is *k-self-centered*. A graph G is *bidegreed* if all vertices of G have one of two possible degrees, cf. [18]. (If x is a vertex of a bidegreed graph G , then $\deg_G(x) \in \{\delta(G), \Delta(G)\}$.) Let finally $G^{(k*)}$, $k \geq 1$, denote the graph obtained from G by attaching k pendant vertices to each vertex of G .

Proposition 3.4 (i) *If G is a regular, 2-self-centered graph, then $C_W(G) = C_{ec}(G) = 1$.*

(ii) *If G be a bidegreed, non-self-centered graph with $\text{diam}(G) = 2$, then $C_W(G) = C_{ec}(G) = 2$.*

(iii) *If G is a regular or bidegreed graph obtained from K_n by removing k edges, $k \in [\lfloor \frac{n}{2} \rfloor]$, then $C_W(G) = C_{ec}(G)$.*

(iv) *If G is a vertex-transitive graph and $k \geq 1$, then $C_{ec}(G^{(k*)}) = C_W(G^{(k*)}) = 2$.*

Proof. (i) As G is self-centered, $C_{ec}(G) = 1$. Combining Proposition 1.1 and the fact that G is regular, we have $C_W(G) = 1$.

(ii) Since G is not self-centered and $\text{diam}(G) = 2$, we have $C_{ec}(G) = 2$ and at least one vertex must have eccentricity 1, that is, $\Delta(G) = n(G) - 1$. As G is bidegreed, all the vertices that have degree smaller than $n(G) - 1$ must have the same degree. In view of Proposition 1.1, we have $C_W(G) = 2$.

(iii) If G is regular, then since $k \leq \lfloor \frac{n}{2} \rfloor$, G is obtained from K_n by removing a perfect matching (in which case n is even and $k = n/2$). Then the assertion follows by (i). Otherwise G is not regular. But then G is bidegreed and hence G fulfils the assumption of (ii).

(iv) Since G is vertex-transitive, G is a transmission regular, self-centered graph. Assume that $\text{diam}(G) = d$, $n = n(G)$, and let $V(G) = \{v_1, \dots, v_n\}$. Let $V(G^{(k*)}) = V(G) \cup \{v_i^{(j)} : i \in [n], j \in [k]\}$ with $v_i v_i^{(j)} \in E(G^{(k*)})$ for $j \in [k]$. From the structure of $G^{(k*)}$ we have $\text{ec}_{G^{(k*)}}(v_i) = d + 1$ and $\text{ec}_{G^{(k*)}}(v_i^{(j)}) = d + 2$ for every $i \in [n]$, $j \in [k]$. Hence $C_{ec}(G^{(k*)}) = 2$.

On the other hand, $\text{Tr}_{G^{(k*)}}(v_i)$ is the same for all $i \in [n]$. Since $v_i^{(j)}$ is a pendant vertex, we see that $\text{Tr}_{G^{(k*)}}(v_i^{(j)}) = \text{Tr}_{G^{(k*)}}(v_i) + (n + 1)k - 2$ is the same for every $j \in [k]$. Thus $C_W(G^{(k*)}) = 2$. \square

The class of graphs from Proposition 3.4 (i) contains vertex-transitive graphs as a proper subclass. For instance, if G is an arbitrary regular graph that is not vertex-transitive, then the join of two copies of G is a regular, 2-self-centered graph, but not vertex-transitive. (The *join* of graphs G and H is obtained from the disjoint union of G and H by adding all possible edges between vertices of G and vertices of H .)

To show that there exist graphs that have the same Wiener complexity and eccentric complexity which is arbitrary large, Cartesian and lexicographic product graphs can be used. We have already defined the Cartesian product. The *lexicographic product* $G \circ H$ of graphs G and H also has the vertex set $V(G) \times V(H)$, vertices (g, h) and (g', h') being adjacent if either $gg' \in E(G)$, or $g = g'$ and $hh' \in E(H)$.

Theorem 3.5 (i) *If G is a graph with $C_W(G) = C_{ec}(G)$, and H is a self-centered, transmission regular graph, then*

$$C_{ec}(G \square H) = C_W(G \square H) = C_W(G).$$

(ii) *If H is a regular graph and $n \geq 4$, then*

$$C_W(P_n \circ H) = C_{ec}(P_n \circ H) = \left\lceil \frac{n}{2} \right\rceil.$$

Proof. (i) The assumption that H is a self-centered, transmission regular graph, means that $C_{ec}(H) = C_W(H) = 1$. Then Theorem 2.1 implies that

$$C_{ec}(G \square H) = C_{ec}(G) + C_{ec}(H) - 1 = C_{ec}(G) = C_W(G).$$

On the other hand, from (4) we get $C_W(G \square H) = C_W(G)$ because $C_W(H) = 1$.

(ii) Let $V(P_n) = \{v_1, \dots, v_n\}$ with natural adjacency relation.

Consider first the case when $H = K_1$. Then $P_n \circ H = P_n \circ K_1 = P_n$. Clearly, $ec_{P_n}(v_i) = |i|_n$, where $|i|_n = \max\{i - 1, n - i\}$. Consequently $C_{ec}(P_n) = \left\lceil \frac{n}{2} \right\rceil$. Moreover, if $i \in [n]$, then $\text{Tr}_{P_n}(v_i) = \binom{i}{2} + \binom{n-i+1}{2}$. In particular, if $i, j \leq \left\lceil \frac{n}{2} \right\rceil$, $i \neq j$, then $\text{Tr}_{P_n}(v_i) \neq \text{Tr}_{P_n}(v_j)$, and $\text{Tr}_{P_n}(v_i) = \text{Tr}_{P_n}(v_{n-i+1})$. It follows that $C_W(P_n) = \left\lceil \frac{n}{2} \right\rceil$.

Let now H be an arbitrary regular graph and consider the lexicographic product $P_n \circ H$. Note first that $d_{P_n \circ H}((v_i, h), (v_i, h')) \leq 2$ for any vertices $h, h' \in V(H)$. Moreover, $d_{P_n \circ H}((v_i, h), (v_j, h')) = d_{P_n}(v_i, v_j)$ for $i \neq j$. Since $n \geq 4$ it follows that $ec_{P_n \circ H}((v_i, h)) = ec_{P_n}(v_i) = |i|_n$ and hence $C_{ec}(P_n \circ H) = \left\lceil \frac{n}{2} \right\rceil$.

Consider now vertices (v_i, h) and (v_i, h') of $P_n \circ H$ for some $i \in [n]$ and $h, h' \in V(H)$, $h \neq h'$. Let $V_i = \{(v_i, x) : x \in V(H)\}$. Since H is regular, both vertices (v_i, h) and (v_i, h') have the same number of neighbors in V_i . Moreover, the distance

between them and their non-neighbors in V_i is 2. Since we already observed that $d_{P_n \circ H}((v_i, h), (v_j, h'')) = d_{P_n}(v_i, v_j) = d_{P_n \circ H}((v_i, h'), (v_j, h''))$ for every $h'' \in V(H)$, we get that $\text{Tr}_{P_n \circ H}((v_i, h)) = \text{Tr}_{P_n \circ H}((v_i, h'))$, that is, the vertices of V_i have the same transmission. Moreover, by the argument from the case $H = K_1$ we also get that if $i, j \leq \lceil \frac{n}{2} \rceil$, $i \neq j$, and $h \in V(H)$, then $\text{Tr}_{P_n \circ H}((v_i, h)) \neq \text{Tr}_{P_n \circ H}((v_j, h))$, and $\text{Tr}_{P_n \circ H}((v_i, h)) = \text{Tr}_{P_n \circ H}((v_{n-i+1}, h))$. We conclude that $C_W(P_n \circ H) = \lceil \frac{n}{2} \rceil$. \square

For a connected graph G , the set of vertices at the given distance from $C(G)$ is called a *distance-level* of G . A tree T is *center-regular* if the vertices in a given distance-level have the same degree. Note that this in particular implies that if T is bicentered, then the central vertices have the same degree.

Proposition 3.6 *If T is a center-regular tree, then $C_W(T) = C_{\text{ec}}(T)$.*

Proof. Let L_i , $0 \leq i \leq k$, be the sets of vertices at distance i from $C(T)$, where $k = \text{rad}(T)$. Let u and v be arbitrary vertices from L_i . Then since T is center-regular, there exists an automorphism $\varphi \in \text{Aut}(T)$ such that $\varphi(u) = v$. This implies that all the vertices of L_i have the same eccentricity as well as the same transmission. Hence, $C_{\text{ec}}(T) \leq k + 1$ and $C_W(T) \leq k + 1$. On the other hand, it is obvious that $C_{\text{ec}}(T) \geq k + 1$ and $C_W(T) \geq k + 1$. \square

The family of center-regular trees includes as a special case the recently introduced *degree-eccentricity regular (DE-regular for short)* trees [25]. The degree-eccentricity regular trees are defined as the trees T in which $\deg_T(v) + \text{ec}_T(v)$ is a fixed constant for every vertex $v \in V(T)$. Clearly, such a tree is center-regular. In [25], among other results, all the molecular *DE*-regular trees were completely characterized.

Remark 3.7 *The result from Proposition 3.6 can be generalized as follows. Let G be a graph such that for every two vertices in a distance-level with respect to $C(G)$, there exists an automorphism mapping one vertex to the other. If, in addition, transmissions are pairwise different for distance-levels, then $C_W(G) = C_{\text{ec}}(G)$.*

3.2 Graphs G with $C_{\text{ec}}(G) > C_W(G)$

In this section, we construct graphs G in which the Wiener complexity is arbitrarily smaller than the eccentric complexity. The existence of such graphs is not obvious at the first sight. Consider the following example.

Let Z_k , $k \geq 1$, be the graph obtained by attaching a pendant vertex to each of two diametrical vertices in a cycle C_{2k+2} . Let u and v be the degree 3 vertices of Z_k ,

let u' and v' be its respective neighbors of degree 1, and let x_1, \dots, x_k and y_1, \dots, y_k be the u, v -paths in Z_k . Since the transmission of a vertex in C_n is $\lfloor \frac{n^2}{4} \rfloor$, see [19], we have

$$\begin{aligned} \text{Tr}_{Z_k}(x_i) &= \text{Tr}_{Z_k}(y_i) \\ &= \text{Tr}_{C_{2k+2}}(x_i) + (i+1) + (k+1-i+1) \\ &= (k+1)^2 + k + 3, \end{aligned}$$

for $i \in [k]$. Moreover, $\text{Tr}_{Z_k}(u) = \text{Tr}_{Z_k}(v) = (k+1)^2 + k + 3$, and $\text{Tr}_{Z_k}(u') = \text{Tr}_{Z_k}(v') = (k+1)^2 + 3k + 5$. On the other hand, $\text{Ec}(Z_k) = \{k+1, k+2, k+3\}$. Thus $C_{\text{ec}}(Z_k) - C_W(Z_k) = 1$.

Each of the graphs Z_k leads to another infinite family of graphs for which the eccentricity complexity exceeds the Wiener complexity. For this sake we recall the following result.

Corollary 3.8 ([1, Corollary 3.2]) *If G is a graph and H a graph with $C_W(H) = 1$, then $C_W(G \square H) = C_W(G)$.*

Proposition 3.9 *If G is a graph with $C_{\text{ec}}(G) > C_W(G)$ and $k \geq 1$, then*

$$C_{\text{ec}}(G \square Q_d) > C_W(G \square Q_d).$$

Proof. By Corollary 3.8, $C_W(G \square Q_d) = C_W(G)$. On the other hand, Theorem 2.1 implies that $C_{\text{ec}}(G \square Q_d) = C_{\text{ec}}(G) + C_{\text{ec}}(Q_d) - 1 = C_{\text{ec}}(G)$. \square

Note that in the proof of Proposition 3.9, the family of hypercubes could be replaced by an arbitrary family of graphs $\{H_i\}_{i \geq 1}$ with $C_W(H_i) = 1$.

We have thus seen that there are infinitely many graphs with eccentric complexity larger than the Wiener complexity. In the above families, this difference was 1. We now demonstrate that the difference can be arbitrarily large. Let $Z = Z_1$ and let Z^m be m^{th} power of Z with respect to Cartesian product, that is, the Cartesian product of m copies of Z .

Theorem 3.10 *If $k \geq 0$, then $C_{\text{ec}}(Z^{2^k}) - C_W(Z^{2^k}) = 2^k$.*

Proof. The proof proceeds by induction on k . The induction hypothesis is that $C_{\text{ec}}(Z^{2^k}) = 2^{k+1} + 1$, $C_W(Z^{2^k}) = 2^k + 1$, and that $\text{Tr}(Z^{2^k})$ is an arithmetic progression with step 4.

As we have observed above, $\text{Ec}(Z) = \{2, 3, 4\}$ and $\text{Tr}(Z) = \{8, 12\}$. Hence, the hypothesis holds for $k = 0$.

Suppose the hypothesis holds for $k - 1$. By Theorem 2.1,

$$C_{ec}(Z^{2^k}) = C_{ec}(Z^{2^{k-1}} \square Z^{2^{k-1}}) = (2^k + 1) + (2^k + 1) - 1 = 2^{k+1} + 1.$$

Let x be the smallest number from the set $\text{Tr}(Z^{2^{k-1}})$. By (5) and the induction hypothesis that $\text{Tr}(Z^{2^k})$ is an arithmetic progression with step 4, we have

$$\begin{aligned} \text{Tr}(Z^{2^k}) &= \{2x, 2x + 4, 2x + 8, \dots, 2 \cdot (x + 4 \cdot 2^{k-1})\} \\ &= \{2x, 2x + 4, \dots, 2x + 2^{k+2}\}. \end{aligned}$$

Therefore, $C_W(Z^{2^k}) = |\text{Tr}(Z^{2^k})| = \frac{2x + 2^{k+2} - 2x}{4} + 1 = 2^k + 1$. The conclusion now follows immediately. \square

Based on Theorem 2.1 and Proposition 2.5, Theorem 3.10 can be generalized as follows.

Corollary 3.11 *Let G be an arithmetic transmission graph with $C_{ec}(G) > C_W(G)$. Then $C_{ec}(G^2) > C_W(G^2)$.*

Using the same argument as in the proof of Proposition 3.9, we infer that for any positive integer N there exists an infinite family of graphs $\{H_i\}_{i \geq 1}$ such that $C_{ec}(H_i) - C_W(H_i) > N$.

Denote by Y_k the graph of order $2k + 4$ consisting of P_{2k+3} and an additional vertex which is adjacent to the two neighbors of the central vertex of P_{2k+3} . Note that $Y_1 = Z_1$. After a straightforward calculation we get that

$$\text{Tr}(Y_k) = \{(k + 1)(k + 2) + i(i + 1) : i \in [k + 1]\}$$

and

$$\text{Ec}(Y_k) = \{k + 1, k + 2, \dots, 2k + 2\}.$$

Hence $C_{ec}(Y_k) = k + 2$ and $C_W(Y_k) = k + 1$.

Now we have the following natural question. Is it true that if $C_{ec}(G) > C_W(G)$ and $C_{ec}(H) > C_W(H)$, then $C_{ec}(G \square H) > C_W(G \square H)$ holds? The answer is negative, as can be seen from the following result:

$$C_{ec}(Y_k \square Z_k) - C_W(Y_k \square Z_k) = \begin{cases} 1, & k = 2; \\ 0, & k \in \{3, 4\}; \\ -2, & k = 5. \end{cases}$$

The graphs G with $C_{ec}(G) > C_W(G)$ found in this paper have diameter at least 4. On the other hand, it follows from the proof of Proposition 3.1 that there are no such graphs of diameter 2. Hence, we pose:

Problem 3.12 *Does there exist a graph G with $\text{diam}(G) = 3$ and $C_{\text{ec}}(G) > C_W(G)$?*

We have checked by computer that there is no such graph of order at most 10. Since $1 \leq C_{\text{ec}}(G) \leq 2$ for any graph G with $\text{diam}(G) = 3$, the key point for solving the above problem is to determine the existence of transmission regular but non-self-centered graphs with diameter 3.

Finally, we have also checked by computer that there is no such tree T with $C_{\text{ec}}(T) > C_W(T)$ on up to 24 vertices. Hence we pose:

Conjecture 3.13 *There is no tree T with $C_{\text{ec}}(T) > C_W(T)$.*

Acknowledgements

Kexiang Xu is supported by supported by NNSF of China (grant No. 11671202, and the China-Slovene bilateral grant 12-9). Sandi Klavžar acknowledges the financial support from the Slovenian Research Agency (research core funding P1-0297, projects J1-9109, N1-0095, and the bilateral grant BI-CN-18-20-008).

References

- [1] Y. Alizadeh, V. Andova, S. Klavžar, R. Škrekovski, Wiener dimension: Fundamental properties and (5,0)-nanotubical fullerenes, MATCH Commun. Math. Comput. Chem. 72 (2014) 279–294.
- [2] Y. Alizadeh, T. Došlić, K. Xu, On the eccentric complexity of graphs, Bull. Malays. Math. Sci. Soc. (2017) <https://doi.org/10.1007/s40840-017-0564-y>.
- [3] Y. Alizadeh, S. Klavžar, Complexity of topological indices: The case of connective eccentric index, MATCH Commun. Math. Comput. Chem. 76 (2016) 659–667.
- [4] Y. Alizadeh, S. Klavžar, On graphs whose Wiener complexity equals their order and on Wiener index of asymmetric graphs, Appl. Math. Comput. 328 (2018) 113–118.
- [5] F. Buckley, Self-centered graphs, Ann. New York Acad. Sci. 576 (1989) 71–78.
- [6] A. A. Dobrynin, Infinite family of 2-connected transmission irregular graphs, Appl. Math. Comput. 340 (2019) 1–4.

- [7] A. A. Dobrynin, Infinite family of transmission irregular trees of even order, *Discrete Math.* 342 (2019) 74–77.
- [8] A. A. Dobrynin, Infinite family of 3-connected cubic transmission irregular graphs, *Discrete Appl. Math.* 257 (2019) 151–157.
- [9] Z. Du, B. Zhou, N. Trinajstić, Extremal properties of the Zagreb eccentricity indices, *Croat. Chem. Acta* 85 (2012) 359–362.
- [10] P. C. Fishburn, Packing graphs with odd and even trees, *J. Graph Theory* 7 (1983) 369–383.
- [11] S. Gupta, M. Singh, A. K. Madan, Connective eccentricity index: a novel topological descriptor for predicting biological activity, *J. Mol. Graph. Model.* 18 (2000) 18–25.
- [12] R. Hammack, W. Imrich and S. Klavžar, *Handbook of Product Graphs*, Second Edition, CRC Press, Boca Raton, FL, 2011.
- [13] D. A. Jemilet, I. Rajasingh, Wiener dimension of spiders, k -ary trees and binomial trees, *Int. J. Pure Appl. Math.* 109 (2016) 143–149.
- [14] S. Klavžar, D. A. Jemilet, I. Rajasingh, P. Manuel, N. Parthiban, General transmission lemma and Wiener complexity of triangular grids, *Appl. Math. Comput.* 338 (2018) 115–122.
- [15] S. Klavžar, H. Liu, P. Singh, K. Xu, Constructing almost peripheral and almost self-centered graphs revisited, *Taiwanese J. Math.* 21 (2017) 805–817.
- [16] H. Lin, K. Ch. Das, B. Wu, Remoteness and distance eigenvalues of a graph, *Discrete Appl. Math.* 215 (2016) 218–224.
- [17] B. D. McKay, A. Piperno, Practical Graph Isomorphism, II, *J. Symbolic Comp.* 60 (2014) 94–112.
- [18] W. L. Myrvold, M. N. Ellingham, D. G. Hoffman, Bidegreed graphs are edge reconstructible, *J. Graph Theory* 11 (1987) 281–302.
- [19] J. Plesník, On the sum of all distances in a graph or digraph, *J. Graph Theory* 8 (1984) 1–21.
- [20] X. Qi, B. Zhou, J. Li, Zagreb eccentricity indices of unicyclic graphs, *Discrete Appl. Math.* 233 (2017) 166–174.

- [21] V. Sharma, R. Goswami, A. K. Madan, Eccentric connectivity index: A novel highly discriminating topological descriptor for structure-property and structure-activity studies, *J. Chem. Inf. Comput. Sci.* 37 (1997) 273–282.
- [22] D. Vukičević, A. Graovac, Note on the comparison of the first and second normalized Zagreb eccentricity indices, *Acta Chim. Slov.* 57 (2010) 524–528.
- [23] K. Xu, Y. Alizadeh, K. Ch. Das, On two eccentricity-based topological indices of graphs, *Discrete Appl. Math.* 233 (2017) 240–251.
- [24] K. Xu, K. Ch. Das, H. Liu, Some extremal results on the connective eccentricity index of graphs, *J. Math. Anal. Appl.* 433 (2016) 803–817.
- [25] K. Xu, X. Gu, I. Gutman, Relations between total irregularity and non-self-centrality of graphs, *Appl. Math. Comput.* 337 (2018) 461–468.
- [26] K. Xu, X. Li, Comparison between two eccentricity-based topological indices of graphs, *Croat. Chem. Acta* 89 (2016) 499–504.
- [27] H. Zhang, S. Li, B. Xu, Extremal graphs of given parameters with respect to the eccentricity distance sum and the eccentric connectivity index, *Discrete Appl. Math.* 254 (2019) 204–221.