

## 1-FACTORIZATION OF THE COMPOSITION OF REGULAR GRAPHS

Tomaž Pisanski, John Shawe-Taylor, Bojan Mohar

**Abstract.** 1-factorability of the composition of graphs is studied. The followings sufficient conditions are proved:  $G[H]$  is 1-factorable if  $G$  and  $H$  are regular and at least one of the following holds: (i) Graphs  $G$  and  $H$  both contain a 1-factor, (ii)  $G$  is 1-factorable (iii)  $H$  is 1-factorable. It is also shown that the tensor product  $G \otimes H$  is 1-factorable, if at least one of two graphs is 1-factorable. This result in turn implies that the strong tensor product  $G \otimes' H$  is 1-factorable, if  $G$  is 1-factorable.

**1.0 Introduction.** The source of inspiration for this paper is rightfully Kotzig's [3]. His simple sufficient conditions for the cartesian product of graphs to be 1-factorable naturally raise the question of when other well known products of graphs are 1-factorable.

In this paper we give analogous results for the composition of graphs and partial results for the tensor and strong tensor products, which extend those announced in [5].

We will leave the basic definitions of graph theory to any standard textbook, for example Harary's Graph Theory [2], and will limit ourselves to defining only lesser known terms and those which may cause confusion.

**2.0 Definitions.** If  $u$  and  $v$  are adjacent vertices of a graph, then we write  $u \sim v$  and denote with  $uv$  the edge joining them.

For a graph  $G$ , let  $V(G)$  denote the vertex set of  $G$  and  $E(G)$  denote its edge set.

The composition, also known as the lexicographical product, of graphs  $G$  and  $H$  is defined as the graph  $G[H]$  with the vertex set  $V(G[H]) = V(G) \times V(H)$  and the edge set  $E(G[H]) = \{(u, v)(u', v') : \text{either } (u = u' \text{ and } v \sim v') \text{ or } u \sim u'\}$ .

---

*AMS Subject Classification* (1980): 05C15

*Key words and phrases:* Regular graph, edge-colouring, 1-factorization.

This work was supported in part by the Boris Kidrič Fund, Yugoslavia.

The tensor product of graphs  $G$  and  $H$  is defined as the graph  $G \otimes H$  with vertex set  $V(G) \times V(H)$  and the edge set

$$E(G \otimes H) = \{(u, v)(u', v') : u \sim u' \text{ and } v \sim v'\}$$

If by  $\deg(v)$  we denote the degree of a vertex  $v$ , then for  $(u, v)$  a vertex in  $G \otimes H$  we have  $\deg((u, v)) = \deg(u) \cdot \deg(v)$ . Thus if  $G$  and  $H$  are regular, so is  $G \otimes H$ .

The graph  $G\{m\}$  is defined as  $G \otimes K_m$ , where  $K_m$  is the complete graph on  $m$  vertices.

If  $G$  and  $H$  have the same vertex set  $V = V(G) = V(H)$ , and disjoint edge sets,  $E(G) \cap E(H) = \emptyset$ , then the sum  $G \oplus H$  is the graph having the vertex set  $V(G \oplus H) = V$  and the edge set  $E(G \oplus H) = E(G) \cup E(H)$ .

Several authors have defined  $G(m)$  as  $G[mK_1]$  [1,4]. Mohar and Pisanski studied 1-factorability of  $G(m)$  in [4]. Here we only note that  $G(m)$  and  $G\{m\}$  are connected by the relation  $G(m) = G\{m\} \oplus mG$ .

The graph  $G[H]$  can be expressed as the sum of the standard cartesian product  $G \times H$  and the graph  $G\{|V(H)|\}$ :

$$G[H] = (G \times H) \oplus G\{|V(H)|\}.$$

If  $G$  is the sum of a series of graphs:

$$G = F_1 \oplus F_2 \oplus \cdots \oplus F_k,$$

we can readily verify the following results:

$$\begin{aligned} G &= F_1\{m\} \oplus F_2\{m\} \oplus \cdots \oplus F_k\{m\}, \\ G \otimes H &= (F_1 \otimes H) \oplus (F_2 \otimes H) \oplus \cdots \oplus (F_k \otimes H). \end{aligned}$$

If each graph  $F_i$  is  $d$ -factorable, it is also clear that  $G$  is  $d$ -factorable, as it can be written as the sum of all the  $d$ -factors of the  $F_i$ .

The strong tensor product  $G \otimes' H$  is defined on the vertex set  $V(G) \times V(H)$  as

$$\begin{aligned} G \otimes' H &= (G \otimes H) \oplus (G \times \{v_1\} \cup G \times \{v_2\} \cup \cdots \cup G \times \{v_m\}), \\ &\text{where } V(H) = \{v_1, v_2, \dots, v_m\}. \end{aligned}$$

**3.0 Known results.** We first restate in our own words Kotzig's result for the cartesian product of regular graphs.

**3.1 THEOREM (Kotzig 1979, [3]):** *If  $G$  and  $H$  are two regular graphs for which at least one of the following conditions holds:*

- (i) *Both  $G$  and  $H$  contain 1-factor,*
- (ii)  *$G$  is 1-factorable,*

(iii)  $H$  is 1-factorable,  
 then the cartesian product  $G \times H$  1-factorable.

Kotzig also showed that these conditions, though being sufficient are not necessary. In particular he showed that for any cubic graph  $G$  and any cycle of length  $n, n$  greater than three, the cartesian product  $G \times C_n$  is 1-factorable [3, Theorem 7]. We shall in turn use this to show that our conditions for 1-factorability of  $G[H]$  are also not necessary.

Finally we shall require the 1-factorability of  $K_{2n}$ , the complete graph on an even number of vertices, and König's well known theorem that a regular bipartite graph is 1-factorable. These theorems can be found in Harar 's book [2 Theorems 9.1, 9.2].

**4.0 Lemma and main theorems.** We fist of all state a lemma concerning the graph  $G(2m)$ .

4.1 LEMMA. *If graph  $G$  is regular, then  $G\{2m\}$  is 1-factorable.*

*Proof.* In section 3 we mentioned that  $K_{2m}$  is 1-factorable. Using this result, let graphs  $F_1, F_2, \dots, F_{2m-1}$  be 1-factors of  $K_{2m}$ , which together make up a 1-factorisation:  $K_{2m} = F_1 \oplus F_2 \oplus \dots \oplus F_{2m-1}$ .

Now we have

$$G\{2m\} = G \otimes K_{2m} = (G \otimes F_1) \oplus (G \otimes F_2) \oplus \dots \oplus (G \otimes F_{2m-1})$$

and since  $F_i = mK_2$  ( $1 < i < 2m$ ), it follows that the tensor product  $G \times F$  can be written as  $G \otimes F_i = m(G \otimes K_2) = mG\{2\}$ .

But the graph  $G\{2\}$  is bipartite, since it has vertices on two levels  $G \times \{1\}$  and  $G \times \{2\}$ , and edges pass only between these two disjoint sets. It is also reregular since  $G$  is regular, and thus it is 1-factorable (cf. Section 3).

Since the tensor product  $G \otimes F_i$  is  $mG\{2\}$ , it is also 1-factorable for each  $i$ . This in turn means that the sum

$$(G \otimes F_1) \oplus (G \otimes F_2) \oplus \dots \oplus (G \otimes F_{2m-1}) = G\{2m\}$$

is 1-factorable.

The main theorem follows readily:

4.2 THEOREM. *If  $G$  and  $H$  are two regular graphs for which at least one of the following holds:*

- (i) both graphs  $G$  and  $H$  contain 1-factor,
- (ii)  $G$  is 1-factorable,
- (iii)  $H$  is 1 factorable,

*then the composition  $G[H]$  of  $G$  and  $H$  is 1-factorable.*

*Proof.* We use the identity  $G[H] = G \times H \oplus G\{|V(H)|\}$ .

By Theorem 3.1  $G \times H$  is 1-factorable in cases (i), (ii) and (iii).

In cases (i) and (iii)  $H$  has at least one 1-factor and thus the number of vertices  $V(H)$  is even. This means that  $G\{|V(H)|\}$  is 1-factorable by Lemma 4.1 and thus  $G[H]$  is 1-factorable.

There remains only case (ii) when  $G$  is 1-factorable. Let  $G = F_1 \oplus F_2 \oplus \cdots \oplus F_k$  be a 1-factorisation of  $G$ . Thus  $F_i$  is  $nK_2$ , where  $G$  has  $2n$  vertices. Now let  $H$  have  $m$  vertices, giving

$$G\{|V(H)|\} = G\{m\} = F_1\{m\} \oplus F_2\{m\} \oplus \cdots \oplus F_k\{m\}.$$

Considering the structure of the  $F_i$ , we have  $F_i\{m\} = n(K_2\{m\})$ .

The graph  $K_2\{m\}$  is a regular bipartite graph of degree  $m - 1$  and so is 1-factorable. This means that  $F_i\{m\}$  is 1-factorable and so in turn are  $G\{|V(H)|\}$  and  $G[H]$ . The theorem is proved.

That the conditions of this theorem are not necessary is demonstrated by the following theorem.

**4.3 THEOREM.** *Let  $G$  be a cubic graph and  $n$  greater than three. Then  $C_n[G]$  is 1-factorable.*

*Proof.*  $C_n \times G = G \times C_n$  is 1-factorable by Kotzig's theorem [3, Theorem 7] concerning the cartesian product of cubic graphs and cycles of length greater than three and  $C_n\{|V(G)|\}$  is 1-factorable by Lemma 4.1 since  $G$  has an even number of vertices.

If  $n$  is odd and  $G$  has no 1-factor, neither graph has a 1-factor and the conditions of Theorem 4.2 are certainly not satisfied. This counter-example is by no means unique. Other such graphs are for instance  $G(2m)$  for graph  $G$  cubic or regular of even degree but not 1-factorable [4].

Let us now consider the tensor products.

**4.4 THEOREM.** *If  $G$  and  $H$  are regular graphs at least one of which is 1-factorable, then the tensor product  $G \otimes H$  is 1-factorable.*

*Proof.* Since the tensor product is commutative, we can without loss of generality take  $G$  to be 1-factorable. Let

$$G = F_1 \text{ plus } F_2 \oplus \cdots \oplus F_k$$

be a 1-factorisation of  $G$ . Thus  $F_i = nK_2$ , where  $G$  has  $2n$  vertices. This gives

$$G \otimes H = (F_1 \otimes H) \oplus (F_2 \otimes H) \oplus \cdots \oplus (F_k \otimes H)$$

and  $F_2 \otimes H = n(K_2 \otimes H) = nH\{2\}$ . By Lemma 4.1  $H\{2\}$  is 1-factorable, this means that  $F_i \otimes H$  is 1-factorable and so also is  $G \otimes H$ .

For the strong tensor product we derive the following results.

**4.5 THEOREM.** *If  $G$  is a 1-factorable and  $H$  a regular graph, then the strong tensor product  $G \oplus' H$  is 1-factorable.*

*Proof.* We can write the strong tensor product  $G \otimes' H$  as  $G \otimes' H = (G \otimes H) \oplus nG$ , where  $n$  is the number of vertices of graph  $H$ . By Theorem 4.4, the tensor product  $G \otimes H$  is 1-factorable, if  $G$  is 1-factorable. That  $nG$  is 1-factorable is also immediate. Thus  $G \otimes' H$  is 1-factorable.

**5.0 Concluding remarks.** A question of interest is whether there exist simple necessary conditions for the different products of regular graphs to be 1-factorable. These are, however, likely to be difficult to find inasmuch as it is harder to disprove 1-factorability than to construct 1-factorisations for various classes of regular graphs.

#### REFERENCES

- [1] A. Bouchet, *Triangular imbeddings into surfaces of a join of equicardinal independent sets following a Eulerian graph*, Lecture notes in mathematics 642; *Theory and Applications of Graphs*, Springer-Verlag, New York, 1976, 86–115.
- [2] F. Harary, *Graph Theory*, Addison-Wesley; Reading, Massachusetts, 1969.
- [3] A. Kotzig, *1-factorizations of Cartesian Products of Regular Graphs*, *J. Graph Theory* **3** (1979), 23–34.
- [4] B. Mohar, T. Pisanski, *Edge-coloring of a family of regular graphs*, *Publ. Inst. Math. (Beograd) (N. S.)* **33 (47)** (1983), 157–162.
- [5] T. Pisanski, B. Mohar, J. Shawe-Taylor, *1-factorizations of compositions of regular graphs*, *Graph Theory Newsletter*, January 1981, Abstract No .4.

Department of Mathematics  
University of Ljubljana  
61000 Ljubljana, Jugoslavija

(Received 26 05 1982)